



# Pybox ligand-promoted copper(I)-catalyzed three-component tandem coupling-annulation of terminal alkynes, amines and *ortho*-alkynylaryl aldehydes

Min Yu<sup>a</sup>, Ying Wang<sup>a</sup>, Chao-Jun Li<sup>b,\*</sup>, Xiaoquan Yao<sup>a,\*</sup>

<sup>a</sup> Department of Applied Chemistry, School of Material Science and Engineering, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, PR China

<sup>b</sup> Department of Chemistry, McGill University, Montreal, QC, Canada H3A 2K6

## ARTICLE INFO

### Article history:

Received 27 July 2009

Revised 14 September 2009

Accepted 18 September 2009

Available online 24 September 2009

### Keywords:

Dihydroisoquinoline

Pybox ligand

Three-component tandem addition and cyclization

## ABSTRACT

Pybox ligand was utilized to promote a highly efficient, one-pot, copper(I)-catalyzed three-component tandem addition and cyclization of *ortho*-alkynylaryl aldehydes, primary amines, and terminal alkynes. Dihydroisoquinoline derivatives **2** were achieved with excellent to moderate yields. In some examples, water was found as an additive to activate the reaction with or without Pybox ligand. Compared with the reported methods, in which imine had to be pre-prepared and purified, the novel process provides a greener choice.

© 2009 Elsevier Ltd. All rights reserved.

Dihydroisoquinolines are common structural units of natural products and exhibit interesting biological activities such as antibiotic properties.<sup>1</sup> Among the methods to construct such structures, the nucleophilic addition to acetylenic imines, which were prepared from *ortho*-alkynylaryl aldehydes and primary amines, followed by amine–alkyne cyclization is the most useful.<sup>2</sup> However, in most of the cases, the key starting material, acetylenic imines had to be pre-prepared from *ortho*-alkynylaryl aldehyde (**1**) and amine (**4**), and then were utilized as pure compounds. To the best of our knowledge, there are only a few examples utilizing acetylenic imines *in situ* prepared.<sup>3</sup> Since the imines are easily decomposed in the presence of water, the following reactions were usually carried out in an anhydrous solvent. Recently, Wu and co-workers reported an ultrasound-assisted, copper-surfactant co-catalyzed, three-component reaction of *ortho*-alkynylaryl aldehydes, primary amines, and terminal alkynes.<sup>3c</sup> As an alternative choice, herein, we hope to report a copper(I)–Pybox complex-catalyzed, three-component tandem coupling-annulation of terminal alkynes, amines, and *ortho*-alkynylaryl aldehydes, in which 1-(arylethynyl)-1,2-dihydroisoquinolines (**2**) were achieved with excellent to moderate yields.<sup>4</sup>


Metal-mediated Grignard-type reactions represent some of the most common methodologies in C–C bond formations. The methods based on the catalytic reaction of C–H bonds provide a highly

atom-economic alternative, which avoids the utilization of stoichiometric amounts of metal and halides when compared to the classical methods.<sup>5</sup> During our earlier studies on the metal acetylides of group IB, such as Cu(I), Ag(I), and Au(I), we realized that the reactivities of the C–M bonds in metal acetylides could be adjusted by the introduction of N-donor and P-donor ligands, and in most cases, water was found to be helpful to cleave the C–M bonds.<sup>6</sup> Following this method, in current reaction, Pybox ligand was used with/without water to promote a copper(I)-catalyzed three-component tandem coupling-annulation.

2-(Phenylethynyl)benzaldehyde, aniline, and phenyl acetylene were selected as the prototype reaction. Referring to the reported method, in which the imine substrate had to be prepared previously, AgOTf was detected as the first attempt.<sup>2b</sup> It can be seen that under the literature conditions, no desired product **2a** was observed, while 2-phenylnaphthalene (**3**) was isolated as the major product with 50% of yield (Table 1, entry 1). While the loading of catalyst was increased to 10 mol %, **3** was still the only product. Considering the mechanism of copper-catalyzed formation of **3**, which resulted from a hetero-Diels–Alder reaction of **1a** and **5a**, and followed by rearrangement at higher temperature,<sup>7</sup> the reaction temperature was then decreased to 40 °C for 2 days, and 47% of **2a** was obtained without **3**. We tried to introduce water into the system. It was shown obviously that the yield of **2a** was decreased from 47% to 5%, and **3** became the major product (entry 4 vs entry 3). The above-mentioned results suggest that AgOTf was not an effective catalyst in the three-component tandem coupling-annulation reaction.

\* Corresponding authors. Tel./fax: +1 514 3988457 (C.-J.L.); tel.: +86 25 52112902; fax: +86 25 52112626 (X.Y.).

E-mail addresses: [cj.li@mcgill.ca](mailto:cj.li@mcgill.ca) (C.-J. Li), [yaoxq@nuaa.edu.cn](mailto:yaoxq@nuaa.edu.cn) (X. Yao).


**Table 1**Ag(I), Cu(I) and Cu(II)-catalyzed three-component tandem coupling-annulation of **1a**, **4a**, and **5a**<sup>a</sup>

Entry	Conditions	Yield <sup>b</sup> (%)	
		2a	3
1	AgOTf (5 mol %), DCE, 80 °C/6 h	0	50
2	AgOTf (10 mol %), DCE, 80 °C/9 h	0	48
3	AgOTf (10 mol %), DCE, 40 °C/2d	47	0
4	AgOTf (10 mol %), DCE/H <sub>2</sub> O(15:1), 40 °C/9 h	<5	32
5	CuOTf (10 mol %), DCE, 40 °C/9 h	61	0
6	CuOTf (10 mol %), CH <sub>2</sub> Cl <sub>2</sub> , 40 °C/9 h	53	0
7	CuOTf (10 mol %), CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O(15:1), 40 °C/9 h	68	0
8	CuOTf (5 mol %), CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O(15:1), 40 °C/2d	76	0
9	CuOTf (10 mol %), H <sub>2</sub> O, 40 °C/9 h	52	0
10	CuOTf (10 mol %), toluene, 40 °C/9 h	17	0
11	CuOTf (10 mol %), toluene/H <sub>2</sub> O(15:1), 40 °C/9 h	60	0
12	CuOTf (10 mol %), THF, 40 °C/9 h	33	0
13	CuOTf (10 mol %), THF/H <sub>2</sub> O(15:1), 40 °C/9 h	15	0
14	Cu(OTf) <sub>2</sub> (5 mol %), CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O(20:1), 40 °C/2d	82	0

<sup>a</sup> 2-(Phenylethynyl)benzaldehyde (0.2 mmol), aniline (0.24 mmol, 1.2 equiv), and phenylacetylene (0.3 mmol, 1.5 equiv) in 1.5 mL of selected solvent.<sup>b</sup> Isolated yield.

Copper(I) and copper(II) salts were then detected in the reaction (entries 5–14). It can be seen from Table 1 that the two catalysts can catalyze the reaction effectively with the promotion of water (entries 7, 8, and 14). However, compared with the reported method, with imine as a starting material,<sup>2b</sup> the yield is far from ideal. Considering the reported methods about copper, silver, and gold-catalyzed C–C bond formation reactions involved in the metal acetylides as key intermediates, in which ligands and/or water were believed to play the key role in adjusting the reactivities of M–C bond in metal acetylides,<sup>6</sup> a series of N-donor ligands in Figure 1 were detected for the reaction, and the results are listed in Table 2.

Several kinds of sp<sup>2</sup> N-donor ligands were scanned. Pybox **8** proved to be the best of choices, and up to 97% of **2a** was achieved

**Figure 1.** The ligands utilized in the reaction.**Table 2**Ligand-promoted, copper-catalyzed three-component tandem coupling-annulation of **1a**, **4a**, and **5a**<sup>a</sup>

Entry	Conditions	Yield <sup>b</sup> (%)	
		2a	3
1 <sup>c</sup>	CuOTf/ <b>6</b> , CH <sub>2</sub> Cl <sub>2</sub> , 40 °C/9 h	60	25
2 <sup>d</sup>	CuOTf/ <b>7</b> , CH <sub>2</sub> Cl <sub>2</sub> , 40 °C/9 h	56	19
3	CuOTf/ <b>8</b> , CH <sub>2</sub> Cl <sub>2</sub> , 40 °C/9 h	97	0
4 <sup>e</sup>	CuOTf/ <b>8</b> , CH <sub>2</sub> Cl <sub>2</sub> , 40 °C/9 h	89	0
5	CuOTf/ <b>8</b> , CH <sub>2</sub> Cl <sub>2</sub> , rt/2d	12	0
6	CuOTf/ <b>8</b> , CH <sub>2</sub> Cl <sub>2</sub> /H <sub>2</sub> O(15:1), 40 °C/9 h	60	0
7	CuOTf/ <b>8</b> , DCE, 40 °C/9 h	13	54
8	CuOTf/ <b>8</b> , DCE/H <sub>2</sub> O(15:1), 40 °C/9 h	20	30
9	CuOTf/ <b>8</b> , DCE/H <sub>2</sub> O(1:1), 40 °C/9 h	36	9
10	CuOTf/ <b>8</b> , toluene, 40 °C/9 h	94	0
11	CuOTf/ <b>8</b> , THF, 40 °C/9 h	72	0
12	CuOTf/ <b>8</b> , THF/H <sub>2</sub> O(15:1), 40 °C/9 h	92	0
13	CuOTf/ <b>8</b> , THF/H <sub>2</sub> O(1:1), 40 °C/9 h	35	0
14	Cu(OTf) <sub>2</sub> / <b>8</b> , CH <sub>2</sub> Cl <sub>2</sub> , 40 °C/9 h	84	4

<sup>a</sup> CuOTf (10 mol %), ligand (12 mol %), 2-(phenylethynyl)benzaldehyde (0.2 mmol), aniline (0.24 mmol, 1.2 equiv), and phenylacetylene (0.3 mmol, 1.5 equiv) in 1.5 mL of selected solvent.<sup>b</sup> Isolated yield.<sup>c</sup> CuOTf (10 mol %), ligand **6** (30 mol %).<sup>d</sup> CuOTf (10 mol %), ligand **7** (12 mol %).<sup>e</sup> Anhydrous MgSO<sub>4</sub> (100 mg) was added.


at 40 °C overnight in CH<sub>2</sub>Cl<sub>2</sub> (Table 2, entry 3). When desiccant, anhydrous MgSO<sub>4</sub>, was added in the reaction to remove the 1 equiv of water produced in situ, the yield decreased (entry 4 vs entry 3). However, with the introduction of additional water, the reaction became worse (entry 6). It seems that a proper amount of water might be of benefit to the reaction. Some other solvents were also tested, and with Pybox–CuOTf as catalyst, 94% and 92% of yields were observed in toluene and THF solutions, respectively. It should be mentioned that in the reaction with THF as a solvent, the addition of water improved the reaction significantly, while only 72% of **2a** was obtained in dried THF (entry 12 vs entry 11).

Subsequently, various acetylenic aldehydes were reacted with terminal alkynes and anilines under the standard reaction conditions (Table 3).<sup>8</sup> Obviously, the *ortho*-alkynylaryl aldehydes with the conjugated group on R<sup>1</sup> provide much better reactivities than the substrates with non-conjugated alkyl group (Table 3, entries 1–12 vs entries 13–15).

With (2-phenylethynyl)aldehyde (**1a**) as substrate, both electron-donating and electron-deficient arylacetylenes provide good to excellent yields for the desired products (entries 1–4), and a >99% of yield was achieved with 1-bromo-4-ethynylbenzene (**5b**) as the terminal alkyne (entry 2). On the other hand, aniline provides the best reactivity in comparison with other substituted aromatic amines (entry 1 vs entries 5–7). When cyclohexenyl group was introduced on R<sup>1</sup>, ~80% of yields were observed for different substituted arylacetylenes and anilines (entries 8–12).

However, when alkyl group was introduced on R<sup>1</sup>, the reactivities were decreased obviously, only moderate yields were obtained with *n*-butyl-substituted **1c** as a substrate (entries 13–15).

In order to further understand the effect of ligand and water on the reaction, copper phenylacetylide **9**, which was suggested as the

**Table 3**Pybox–CuOTf catalyzed three-component tandem coupling-annulation of **1**, **4**, and **5**<sup>a</sup>

Entry	1:R <sup>1</sup>	4:R <sup>2</sup>	5:R <sup>3</sup>	Product <b>2</b>	Yield <sup>b</sup> (%)
1	Ph– ( <b>1a</b> )	H ( <b>2a</b> )	H ( <b>5a</b> )	<b>2a</b>	97
2	Ph–	H	Br ( <b>5b</b> )	<b>2b</b>	>99
3	Ph–	H	Cl ( <b>5c</b> )	<b>2c</b>	87
4	Ph–	H	CH <sub>3</sub> ( <b>5d</b> )	<b>2d</b>	85
5	Ph–	CH <sub>3</sub> ( <b>2b</b> )	H	<b>2e</b>	77
6	Ph–	Br ( <b>2c</b> )	H	<b>2f</b>	93
7	Ph–	Br	Br	<b>2g</b>	86
8	Cyclohexenyl– ( <b>1b</b> )	H	H	<b>2h</b>	83
9	Cyclohexenyl–	H	CH <sub>3</sub>	<b>2i</b>	77
10	Cyclohexenyl–	H	Br	<b>2j</b>	76
11	Cyclohexenyl–	Br	H	<b>2k</b>	85
12	Cyclohexenyl–	CH <sub>3</sub>	H	<b>2l</b>	84
13	<i>n</i> -Bu– ( <b>1c</b> )	H	H	<b>2m</b>	40
14	<i>n</i> -Bu–	H	CH <sub>3</sub>	<b>2n</b>	49
15	<i>n</i> -Bu–	Br	H	<b>2o</b>	42

<sup>a</sup> Conditions: CuOTf (10 mol %) and **8** (12 mol %) were mixed in 1.5 mL of CH<sub>2</sub>Cl<sub>2</sub>, stirred at room temperature for 20 min, and then *ortho*-alkynylaryl aldehyde (0.2 mmol), aniline (0.24 mmol, 1.2 equiv), and arylacetylene (0.3 mmol, 1.5 equiv) were added. The mixture was then stirred at 40 °C overnight.

<sup>b</sup> Isolated yield.

key intermediate in the reaction, was prepared and utilized as a catalyst (or a catalyst precursor). It can be seen from Table 4 that both water and Pybox ligand improved the reactions significantly. Furthermore, the results in entries 2 and 5 also suggested that a proper amount of water is helpful to the reaction, while worse results were observed with pure water (Table 4, entry 2 vs entry 3, entry 5 vs entry 6). Therefore, we reasoned that the Pybox ligand might be weakening and activating the copper–carbon bond in the copper acetylide. Coordination of the electron-donating Pybox ligand increases the electron density on copper, resulting in the weakening of the copper–carbon bond, which was then cleaved with the promotion of water produced in situ from the aldehydes and anilines.

Thus, a tentative mechanism for the Pybox–Cu(I) complex-catalyzed tandem addition/cyclization is proposed: The reaction of terminal alkynes with Pybox–Cu(I) complex generates the cop-

per(I) acetylide species in the presence of anilines. The acetylide then reacts with imine formed in situ from aldehyde and aniline to give propargylic amine, followed by the amine–alkyne cyclization to give the vinylcopper intermediate; the carbon–copper bond is then quenched to give the final product **2** by protonolysis and regenerates the catalyst.<sup>9</sup>


In conclusion, with the help of Pybox ligand, dihydroisoquinoline derivatives **2** were achieved successfully via a highly efficient, one-pot, copper(I)-catalyzed three-component tandem addition and cyclization of *ortho*-alkynylaryl aldehydes, anilines, and terminal alkynes. Compared with the reported methods, in which the imine intermediate had to be pre-prepared and purified, the novel process provides a greener choice. The three-component tandem addition and cyclization with other nucleophiles based on this catalytic system are currently under investigation.

## Acknowledgment

We are grateful to the financial support from National Natural Science Foundation of China (20602018 to X.Y.).

## References and notes

- (a) Biber, B.; Muske, J.; Ritzan, M.; Graft, U. *J. Antibiot.* **1998**, *51*, 381; (b) Maruse, N.; Goto, M. *J. Antibiot.* **1998**, *51*, 545; (c) Wang, W.; Breining, T.; Li, T.; Milbum, R.; Attardo, G. *Tetrahedron Lett.* **1998**, *39*, 2459, and the references cited therein.
- For representative examples, see: (a) Obika, S.; Kono, H.; Yasui, Y.; Yanada, R.; Takemoto, Y. *J. Org. Chem.* **2007**, *72*, 4462, and the references cited therein; (b) Asao, N.; Yudha, S. S.; Nogami, T.; Yamamoto, Y. *Angew. Chem., Int. Ed.* **2005**, *44*, 5526; (c) Yanada, R.; Obika, S.; Kono, H.; Takemoto, Y. *Angew. Chem., Int. Ed.* **2006**, *45*, 3822; Some other examples involved in the nucleophilic addition to *ortho*-alkynylaryl aldehydes, see: (d) Asao, N.; Nogami, T.; Takahashi, K.; Yamamoto, Y. *J. Am. Chem. Soc.* **2002**, *124*, 764; (e) Patil, N. T.; Yamamoto, Y. *J. Org. Chem.* **2004**, *69*, 5139; (f) Nakamura, H.; Ohtaka, M.; Yamamoto, Y. *Tetrahedron Lett.* **2002**, *43*, 7631; (g) Asao, N.; Chan, C. S.; Takahashi, K.; Yamamoto, Y. *Tetrahedron* **2005**, *61*, 11322; (h) Patil, N. T.; Pahadi, N. K.; Yamamoto, Y. *J. Org. Chem.* **2005**, *70*, 10096; (i) Wei, L.-L.; Wei, L.-M.; Pan, W.-B.; Wu, M.-J. *Synlett* **2004**, 1497; Some examples for the electrophilic

**Table 4**Copper phenylacetylide **9** catalyzed three-component tandem coupling-annulation of **1a**, **4a**, and **5a**<sup>a</sup>

Entry	Conditions	Yield <sup>b</sup> (%)
1	<b>9</b> 10 mol %, CH <sub>2</sub> Cl <sub>2</sub> , 40 °C	32
2	<b>9</b> 10 mol %, H <sub>2</sub> O 1 equiv, CH <sub>2</sub> Cl <sub>2</sub> , 40 °C	56
3	<b>9</b> 10 mol %, H <sub>2</sub> O, 40 °C	26
4	<b>9</b> 10 mol %, <b>8</b> 12 mol %, CH <sub>2</sub> Cl <sub>2</sub> , 40 °C	54
5	<b>9</b> 10 mol %, <b>8</b> 12 mol %, H <sub>2</sub> O 1 equiv, CH <sub>2</sub> Cl <sub>2</sub> , 40 °C	81
6	<b>9</b> 10 mol %, <b>8</b> 12 mol %, H <sub>2</sub> O, 40 °C	15

<sup>a</sup> 2-(Phenylethynyl)benzaldehyde (0.2 mmol), aniline (0.24 mmol, 1.2 equiv), and phenylacetylene (0.3 mmol, 1.5 equiv) in 1.5 mL of selected solvent.

<sup>b</sup> Isolated yield.

- cyclization: (j) Barluenga, J.; Vazquez-Villa, H.; Ballesteros, A.; Gonzalez, J. M. *J. Am. Chem. Soc.* **2003**, *125*, 9028; (k) Yue, D.; Ca, N. D.; Larock, R. C. *Org. Lett.* **2004**, *6*, 1581.
3. (a) Ding, Q.; Wu, J. *Org. Lett.* **2007**, *9*, 4959; (b) Ding, Q.; Wang, B.; Wu, J. *Tetrahedron* **2007**, *63*, 12166; (c) Ye, Y.; Ding, Q.; Wu, J. *Tetrahedron* **2008**, *64*, 1378; (d) Ye, S.; Zhou, H.; Wu, J. *Tetrahedron* **2009**, *65*, 1294.
4. The CuOTf/Pybox catalytic system was utilized with TBHP as oxidant in a CDC reaction to synthesize the similar tetrahydroisoquinoline derivatives, see: Li, Z.; MacLeod, P. D.; Li, C.-J. *Tetrahedron: Asymmetry* **2006**, *17*, 590.
5. Recent reviews on C–H activations, see: (a) Dyker, G. *Angew. Chem., Int. Ed.* **1999**, *38*, 1698; (b) Chatani, N.; Asaumi, T.; Yorimitsu, S.; Ikeda, T.; Kakiuchi, F.; Murai, S. *J. Am. Chem. Soc.* **2001**, *123*, 10935; (c) Chen, H.; Schlecht, S.; Semple, T. C.; Hartwig, J. F. *Science* **2000**, *287*, 1995; (d) Goldman, A. S. *Nature* **1993**, *366*, 514; (e) Crabtree, R. H. *J. Organomet. Chem.* **2004**, *689*, 4083; (f) Jia, C.; Kitamura, T.; Fujiwara, Y. *Acc. Chem. Res.* **2001**, *34*, 633; For atom-economy, see: (g) Trost, B. M. *Science* **1991**, *254*, 1471; (h) Trost, B. M. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 259; (i) Anastas, P. T.; Warner, J. C. *Green Chemistry: Theory and Practice*; Oxford University Press: New York, 2000; About Grignard-type reactions utilizing C–H bonds directly in water, see: (j) Herreras, C. I.; Yao, X.; Li, Z.; Li, C.-J. *Chem. Rev.* **2007**, *107*, 2546.
6. (a) Yu, M.; Skouta, R.; Zhou, L.; Jiang, H.-F.; Yao, X.; Li, C.-J. *J. Org. Chem.* **2009**, *74*, 3378; (b) Yao, X.; Li, C.-J. *Org. Lett.* **2006**, *8*, 1953; (c) Yao, X.; Li, C.-J. *Org. Lett.* **2005**, *7*, 4395; (d) Wei, C.; Li, Z.; Li, C.-J. *Synlett* **2004**, 1472; (e) Wei, C.; Mague, J. T.; Li, C.-J. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5749; (f) Wei, C.; Li, C.-J. *J. Am. Chem. Soc.* **2002**, *124*, 5683; (g) Wei, C. M.; Li, C.-J. *J. Am. Chem. Soc.* **2003**, *125*, 9584; (h) Wei, C.; Li, Z.; Li, C.-J. *Org. Lett.* **2003**, *5*, 4473; (i) Li, Z.; Wei, C.; Chen, L.; Varma, R. S.; Li, C.-J. *Tetrahedron Lett.* **2004**, *45*, 2443; (j) Li, C.-J.; Wei, C. *Chem. Commun.* **2002**, 268; (k) Li, C.-J. *Acc. Chem. Res.* **2002**, *35*, 533; (l) Li, C.-J.; Wei, C. *Green Chem.* **2002**, *39*; For recent review about the ligand effects in IB metals-catalyzed reactions, see: (m) Gorin, D. J.; Sherry, B. D.; Toste, F. D. *Chem. Rev.* **2008**, *108*, 3351.
7. (a) Asao, N.; Nogami, T.; Lee, S.; Yamamoto, Y. *J. Am. Chem. Soc.* **2003**, *125*, 10921; (b) Asao, N.; Sato, K.; Menggenbater; Yamamoto, Y. *J. Org. Chem.* **2005**, *70*, 3682.
8. *General procedure*: Copper(I) trifluoromethanesulfonate benzene complex (0.0050 g, 10 mol % to copper(I)) and **8** (0.0072 g, 12 mol %) were mixed in 1.5 mL of CH<sub>2</sub>Cl<sub>2</sub>, stirred at room temperature for 20 min, and then, *ortho*-alkynylaryl aldehyde **1** (0.2 mmol), aniline **4** (0.24 mmol, 1.2 equiv), and arylacetylene **5** (0.3 mmol, 1.5 equiv) were added. The mixture was then stirred at 40 °C overnight. The reaction mixture was purified by flash column chromatography to give dihydroisoquinoline derivatives **2**.
9. See Refs. [6a–c](#) and [e,f](#).