


Selective C–H Olefination of Indolines (C5) and Tetrahydroquinolines (C6) by Pd/S,O-Ligand Catalysis

Wen-Liang Jia, Nick Westerveld, Kit Ming Wong, Thomas Morsch, Matthijs Hakkenes, Kananat Naksomboon, and M. Angeles Fernández-Ibáñez*

Van't Hoff Institute for Molecular Sciences, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

Supporting Information

ABSTRACT: Herein, we report a highly selective C–H olefination of directing-group-free indolines (C5) and tetrahydroquinolines (C6) by Pd/S,O-ligand catalysis. In the presence of the S,O-ligand, a wide range of challenging indolines, tetrahydroquinolines, and olefins was efficiently olefinated under mild reaction conditions. The synthetic potential of this methodology was demonstrated by the efficient olefination of several indoline-based natural products.




Indolines and tetrahydroquinolines (THQs) are ubiquitous structures in natural products and pharmaceuticals.¹ The development of new methodologies that permit the selective C–H functionalization of these structures could considerably widen the extent of current strategies for diversity-oriented synthesis in medicinal chemistry.² In this context, two main strategies to achieve efficient and selective C(sp²)–H functionalization reactions of indolines and THQs were employed. The first approach consists of the use of directing groups attached to the nitrogen atom, which leads to functionalized C7-indolines and C8-THQs (Figure 1a),³ while the second one provides C6-indolines and C7-THQs by using templates attached to the nitrogen atom (Figure 1b).⁴ In the particular case of indolines, the selective C5 functionalization was accomplished via (a) Ru(II)-catalyzed difluoromethylation,⁵ (b) Au(I)-catalyzed alkylation,⁶ and (c) Zn(II)-catalyzed Michael-type Friedel–Crafts alkylation (Figure 1c).⁷ In these examples, only alkyl groups are introduced, and the substrate scope is limited to neutral or electron-rich indolines. To the best of our knowledge, general strategies to selectively obtain C5-olefinated indolines and C6-olefinated THQs are still elusive.⁸ Herein, we report the first C5–H olefination of indolines and C6–H olefination of THQs by Pd/S,O-ligand catalysis (Figure 1d). The reaction in the presence of the S,O-ligand proceeds efficiently with a wide range of indolines, THQs, and olefins, providing the desired olefinated products with excellent selectivity and high yields.


Recently, we found out that the C–H olefination of a variety of aromatic compounds can be promoted by the presence of bidentate S,O-ligands.⁹ In particular, we reported the first general *para*-selective C–H olefination of aromatic amines. Thus, we hypothesized that a selective C–H olefination of indolines and THQs, which are ubiquitous moieties in natural products, could be achieved in the presence of our Pd/S,O-ligand catalyst.

First, we evaluated the reactivity of different *N*-protected (Me, Bn, Boc) indolines under conditions similar to the ones


(a) Functionalization of indolines (C7) and THQs (C8) using directing groups



(b) Functionalization of indolines (C6) and THQs (C7) using templates



(c) C5 alkylation of indolines



(d) This work: general method for the olefination of indolines (C5) and THQs (C6)






Figure 1. Metal-catalyzed selective C–H functionalization of indolines and THQs.

used in the C–H olefination of anilines (see Supporting Information, Table S1).^{9d} We observed that *N*-methyl indoline (1a) provided the highest yield (38%) and C5 selectivity and that indolines were less reactive than anilines as well as more sensitive to higher temperatures. Thus, different temperatures, reaction times, and stoichiometries were tested to obtain a compromise between reactivity and stability. Finally, the

Received: October 3, 2019

Scheme 1. C5 C–H Olefination of *N*-Methyl Indolines^a

After demonstrating the efficiency of our catalytic system in promoting the C–H olefination of indolines, we investigated the olefination of tetrahydroquinolines (Scheme 2). The reaction of *N*-methyl THQ **3a** under the optimal reaction conditions (for optimization of reaction conditions, see the Supporting Information, Tables S9–S11) provided the olefinated product **4a** in 73% isolated yield with perfect selectivity. In contrast with the reactivity observed for indolines bearing methyl groups in the aromatic ring, *N*-methyl 5-methyl THQ **3b** and *N*-methyl 7-methyl THQ **3c** were olefinated in 50 and 54% yield, respectively. The reaction of *N*-methyl 2,2,4,7-tetramethyl THQ (**3d**) furnished the olefinated product in 62% isolated yield. Then, we performed the reactions of more electron poor THQs. Under optimal conditions, *N*-methyl 7-chloro THQ (**3e**) and *N*-methyl 2,3-dihydro-1*H*-quinolin-4-one (**3f**) were olefinated in 53 and 56% yield, respectively, and with excellent selectivity. Reactions with spirotetrahydroquinolines **3g** and **3h** provided the olefinated products in 73 and 71% yield. Finally, we tested the reaction of *N*-methyl 8-chloro THQ (**3i**). As expected, this substrate gave only a trace amount of olefinated product, in line with the reactivity observed with *N,N*-dimethyl *ortho*-substituted anilines.^{9d} Our previous DFT calculations proved that the lack of reactivity of *ortho*-substituent anilines is due to the twist of the nitrogen atom out of the plane, deactivating the aniline. To prove that the same situation occurred in this case, we performed the reaction of the unprotected 8-chloro THQ


Scheme 2. C6 C–H Olefination of *N*-Methyl THQs^a

^aIsolated yields. Selectivities were determined by ¹H NMR analysis of the crude mixture. ^bReaction was performed at 50 °C. ^c1,4-Dioxane was used as the solvent. ^dReaction was performed at 80 °C.

3j, which provided the olefinated product in 32% isolated yield. Although the reaction was still not efficient, the higher yield obtained for the unprotected THQ in comparison with the protected one **3i** confirms that the same situation occurred. Indeed, the reaction of the unprotected 8-ethanone THQ (**3k**) furnished the olefinated product in 78% yield. Again, the key role of the S,O-ligand in the reaction was confirmed by comparing the results of the reaction with and without the ligand.

Next, we evaluated the scope of olefins, as shown in Scheme 3. The reaction of *N*-methyl 3,3-dimethyl indoline (**1k**) with several olefins, including methyl, phenyl, and cyclohexyl acrylates, furnished the olefinated products **5a–c** in high yields (72–76%) and selectivities. α -Methylene- γ -butyrolactone afforded compound **5d** in 79% yield as a mixture of **5d1**, **5d2**, and **5d3** in a ratio 6:12:1. Other activated olefins such as methyl vinyl ketone and vinylphosphonate were also used, providing the olefinated products **5e** and **5f** in 55 and 82% yields, respectively. We also tested the C–H olefination using the spiro-tetrahydroquinoline **3h**. The reaction using vinyl amide and vinyl sulfonate provided the corresponding olefinated products **5g** and **5h**, respectively, in synthetically useful yields (51–68%). To our delight, the challenging substrate styrene was also a suitable olefin for this reaction, providing the olefinated product **5i** in 51% yield.

In conclusion, we developed the first C5–H olefination of indolines and C6–H olefination of THQs by Pd/S,O-ligand catalysis. The reaction in the presence of the S,O-ligand proceeded efficiently with a wide range of indolines, THQs, and olefins, providing the desired olefinated products with excellent selectivity and high yields. Further applications and mechanistic studies are currently ongoing in our laboratory.

Scheme 3. Scope of Olefins^a

^aReaction conditions: (A) **1k** (0.5 mmol), olefin (0.25 mmol), Pd(OAc)₂ (10 mol %), S,O-ligand (10 mol %), PhCO₃tBu (1.0 equiv) in DCE (1.25 mL) at 60 °C for 16 h; (B) **3h** (0.25 mmol), olefin (0.375 mmol), Pd(OAc)₂ (10 mol %), S,O-ligand (10 mol %), PhCO₃tBu (1.0 equiv) in DCE (1.25 mL) at 40 °C for 16 h. Isolated yields. Selectivities were determined by ¹H NMR analysis of the crude mixture.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.9b03505.

General information, synthesis of indoline and quinoline substrates, reaction optimization for the C5 C–H olefination of indolines, general procedure for Pd-catalyzed C5 C–H olefination of indolines, reaction optimization for the C6 C–H olefination of tetrahydroquinolines, general procedure for Pd-catalyzed C6 C–H olefination of tetrahydroquinolines, general procedure for the evaluation of olefins, large scale reaction of Pd-catalyzed C5 C–H olefination of 1,3,3-trimethylindoline, ¹H NMR, ¹³C NMR, and ³¹P NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: m.a.fernandezibanez@uva.nl.

ORCID

M. Ángeles Fernández-Ibáñez: 0000-0002-7694-5911

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We acknowledge financial support from NWO through a VIDI grant (723.013.006). W.-L.J. gratefully acknowledges financial support from the China Scholarship Council (CSC) (File No. 201606180020).

■ REFERENCES

- (1) For selected reviews, see: (a) McCormick, J. L.; McKee, T. C.; Cardellina, J. H., II; Boyd, M. R. *J. Nat. Prod.* **1996**, *59*, 469. (b) Gul, W.; Hamann, M. T. *Life Sci.* **2005**, *78*, 442. (c) Sridharan, V.; Suryavanshi, P. A.; Menéndez, J. C. *Chem. Rev.* **2011**, *111*, 7157. (d) Ruiz-Sanchis, P.; Savina, S. A.; Albericio, F.; Álvarez, M. *Chem. - Eur. J.* **2011**, *17*, 1388. (e) Nammalwar, B.; Bunce, R. A. *Molecules* **2014**, *19*, 204. (f) Chung, P.-Y.; Bian, Z.-X.; Pun, H.-Y.; Chan, D.; Chan, A. S.-C.; Chui, C.-H.; Tang, J. C.-O.; Lam, K.-H. *Future Med. Chem.* **2015**, *7*, 947. (g) Heravi, M. M.; Rohani, S.; Zadsirjan, V.; Zahedi, N. *RSC Adv.* **2017**, *7*, 52852. (h) Muthukrishnan, I.; Sridharan, V.; Menéndez, J. C. *Chem. Rev.* **2019**, *119*, 5057.
- (2) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. *W. Chem. Soc. Rev.* **2016**, *45*, 546.
- (3) For selected examples, see: (a) Kalyani, D.; Deprez, N. R.; Desai, L. V.; Sanford, M. S. *J. Am. Chem. Soc.* **2005**, *127*, 7330. (b) Shi, Z.; Li, B.; Wan, X.; Cheng, J.; Fang, Z.; Cao, B.; Qin, C.; Wang, Y. *Angew. Chem., Int. Ed.* **2007**, *46*, 5554. (c) Nishikata, T.; Abela, A. R.; Huang, S.; Lipshutz, B. H. *J. Am. Chem. Soc.* **2010**, *132*, 4978. (d) Urones, B.; Arráyas, R. G.; Carretero, J. C. *Org. Lett.* **2013**, *15*, 1120. (e) Jiao, L.-Y.; Oestreich, M. *Org. Lett.* **2013**, *15*, 5374. (f) Neufeldt, S. R.; Seigerman, C. K.; Sanford, M. S. *Org. Lett.* **2013**, *15*, 2302. (g) Jiao, L.-Y.; Oestreich, M. *Chem. - Eur. J.* **2013**, *19*, 10845. (h) Pan, S.; Ryu, N.; Shibata, T. *Adv. Synth. Catal.* **2014**, *356*, 929.
- (4) (a) Yang, G.; Lindovska, P.; Zhu, D.; Kim, J.; Wang, P.; Tang, R.-Y.; Movassaghi, M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2014**, *136*, 10807. (b) Tang, R.; Li, G.; Yu, J.-Q. *Nature* **2014**, *507*, 215.
- (5) Yuan, C.; Zhu, L.; Chen, C.; Chen, X.; Yang, Y.; Lan, Y.; Zhao, Y. *Nat. Commun.* **2018**, *9*, 1189.
- (6) Zhang, W.; Xu, G.; Qiu, L.; Sun, J. *Org. Biomol. Chem.* **2018**, *16*, 3889.
- (7) Ertugrul, B.; Kilic, H.; Lafzi, F.; Saracoglu, N. *J. Org. Chem.* **2018**, *83*, 9018.
- (8) For selected examples of C6 functionalization of THQs, see: (a) Kumar, N.; Kumar, R.; Rao, L.; Muthineni, N.; Ramesh, T.; Babu, N.; Meshram, H. *Synthesis* **2017**, *49*, 3171. (b) Ramana, D. V.; Sudheer Kumar, K.; Srujana, E.; Chandrasekharam, M. *Eur. J. Org. Chem.* **2019**, *2019*, 742. For a review, see: (c) Sridharan, V.; Suryavanshi, P. A.; Menendez, J. C. *Chem. Rev.* **2011**, *111*, 7157.
- (9) (a) Naksomboon, K.; Valderas, C.; Gómez-Martínez, M.; Álvarez-Casao, Y.; Fernández-Ibáñez, M. Á. *ACS Catal.* **2017**, *7*, 6342. (b) Naksomboon, K.; Álvarez-Casao, Y.; Uiterweerd, M.; Westerveld, N.; Maciá, B.; Fernández-Ibáñez, M. Á. *Tetrahedron Lett.* **2018**, *59*, 379. (c) Álvarez-Casao, Y.; Fernández-Ibáñez, M. Á. *Eur. J. Org. Chem.* **2019**, *2019*, 1842. (d) Naksomboon, K.; Poater, J.; Bickelhaupt, F. M.; Fernández-Ibáñez, M. A. *J. Am. Chem. Soc.* **2019**, *141*, 6719. For other references using thioether ligands in C–H functionalization reactions, see: (e) Gorsline, B. J.; Wang, L.; Ren, P.; Carrow, B. P. *J. Am. Chem. Soc.* **2017**, *139*, 9605. (f) Zhuang, Z.; Yu, C.-B.; Chen, G.; Wu, Q.-F.; Hsiao, Y.; Joe, C. L.; Qiao, J. X.; Poss, M. A.; Yu, J.-Q. *J. Am. Chem. Soc.* **2018**, *140*, 10363. For a selected example using a sulfoxide ligand in C–H activation, see: Chen, M. S.; White, M. C. *J. Am. Chem. Soc.* **2014**, *126*, 1346.
- (10) The difference in reactivity can be explained by the higher instability of electron-rich indolines, which are also more prone to be oxidized to indoles. To avoid this oxidation, we performed the reaction of 1,7-dimethyl indoline using different oxidants such as AgOAc, Cu(OAc)₂, and O₂. However, we did not observe the desired olefinated product in any of these reactions (see [Supporting Information, Table S8](#)).