

Available online at www.sciencedirect.com



Tetrahedron Letters 45 (2004) 5493-5496

Tetrahedron Letters

## A formal asymmetric synthesis of both enantiomers of the *Erythrina* alkaloid 3-demethoxyerythratidinone

Steven M. Allin,<sup>a,\*</sup> Guy B. Streetley,<sup>a</sup> Martin Slater,<sup>b</sup> Stella L. James<sup>a,†</sup> and William P. Martin<sup>c</sup>

<sup>a</sup>Department of Chemistry, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK <sup>b</sup>BioFocus Discovery Limited, Chesterford Research Park, Saffron Walden, Essex CB10 1XL, UK <sup>c</sup>GlaxoSmithKline Pharmaceuticals, Gunnels Wood Road, Stevenage, Herts SG1 2NY, UK

Received 8 March 2004; revised 26 April 2004; accepted 7 May 2004

**Abstract**—A formal asymmetric synthesis of both enantiomers of the *Erythrina* alkaloid 3-demethoxyerythratidinone is reported through the application of a highly functionalised lactam template as an *N*-acyliminium precursor. © 2004 Elsevier Ltd. All rights reserved.



The genus *Erythrina* is common in tropical and subtropical regions and the alkaloids have been used in indigenous medicine.<sup>1</sup> Members of the *Erythrina* family display curare-like and hypnotic activity, and a variety of pharmacological effects are associated with the erythrinane skeleton including sedative, hypotensive, neuromuscular blocking and CNS activity.<sup>2a</sup> Consequently

<sup>†</sup> Present address: Charnwood Molecular Ltd, The Heritage Building, Beaumont Court, Prince William Road, Loughborough LE11 5GA, UK.

0040-4039/\$ - see front matter  $\odot 2004$  Elsevier Ltd. All rights reserved. doi:10.1016/j.tetlet.2004.05.060

there has been much interest in the synthesis of these alkaloids and derivatives (including pyrroloisoquinoline precursors) over recent years.<sup>2</sup>

3-Demethoxyerythratidinone, **1**, was isolated from *Erythrina lithosperma* in 1973 by Barton and co-workers.<sup>3</sup> Despite being structurally one of the simplest of the *Erythrina* class of alkaloid, it was not until 1984 that Tsuda reported the first total synthesis, in racemic form, of this natural product.<sup>4</sup> Almost a decade later this same group reported an asymmetric route to the enantiomeric form of the natural product, the so-called '*enantio-(-)-3-demethoxyerythratidinone*'.<sup>5</sup>

We have recently concerned ourselves with the development of novel, efficient and stereoselective routes to complex heterocyclic targets by employing *N*-acyliminium intermediates in cyclisation reactions.<sup>6</sup> In a recent publication<sup>6a</sup> we detailed a novel approach to an erythrinane-like ring system through the application of a Meyers chiral lactam template. We now wish to report the successful application of our methodology to construct a functionalised erythrinane ring system in a highly stereoselective manner, thus representing a formal asymmetric synthesis of both enantiomers of the natural product, 3-demethoxyerythratidinone.

Although the chiral lactams pioneered by Meyers have been widely utilised in asymmetric synthesis,<sup>7</sup> to the best of our knowledge, the present application, as a

<sup>\*</sup> Corresponding author. Tel.: +44-1-01509-222559; fax: +44-1-01509-223925; e-mail: s.m.allin@lboro.ac.uk



## Scheme 1.

precursor in an *N*-acyliminium mediated cyclisation reaction leading to erythrinane targets, represents a novel use of this popular chiral template. In order to access the more highly functionalised alkaloid skeleton we were required to prepare the protected keto-acid substrate **3**. This compound was readily accessed by enolate alkylation and subsequent ester group hydrolysis of the commercially available mono-protected diketone **2**, as shown in Scheme 1.

Our synthesis of the required lactam substrate 5 then followed similar methodology to that previously described by us,<sup>6</sup> with the required  $\beta$ -aminoalcohol 4 prepared in 97% yield by reduction of the commercially available amino acid, 3-(3,4-dimethoxyphenyl)-L-alanine, with LiBH<sub>4</sub> in the presence of Me<sub>3</sub>SiCl in THF for 24 h at room temperature. Condensation of substrates 3 and 4 under Dean-Stark conditions in toluene for 48 h gave a 63% yield of the desired lactam 5 as a single diastereoisomer (Scheme 2). The formation of a single product diastereoisomer of lactam 5 from the racemic keto acid 3 requires the epimerisation of the stereogenic centre adjacent to the ketone during the reaction, and this fact has been noted previously for similar substrates in the preparation of polycyclic lactams for use as N-acyliminium precursors.<sup>6a,8</sup>

With 5 in hand, we turned our attention to the proposed asymmetric cyclisation. On treating lactam 5 with 3 equiv of TiCl<sub>4</sub> as Lewis acid activator at low temperature in dichloromethane for 20 h we were pleased to isolate the tetracyclic product 6 in an excellent 92% yield. Perhaps not unexpectedly, the cyclisation under strong Lewis acid conditions had been accompanied by concomitant deprotection of the ketal protecting group. Reducing the number of equivalents of the Lewis acid did not result in selective formation of the corresponding cyclisation product with the ketal group intact, and instead an inseparable product mixture was obtained. Indeed, since the removal of the protecting group did not hinder the immediate progress of our work, we favoured the use of excess Lewis acid to achieve clean and complete deprotection at this stage. <sup>1</sup>H NMR analysis of the crude product mixture revealed the formation of a 10:1 mixture of product diastereoisomers.

The major diastereoisomer 6 was isolated by column chromatography. In our previous work on the synthesis of the related compound 7, prepared by analogous chemistry, we were able to determine the relative stereochemistry by X-ray crystallography. In this present case we were unable to obtain suitable crystals for X-ray, however the absence of an NOE between the protons at positions 4 and 13a supports the proposed structure; one would not expect a change in product stereochemistry to be favoured on simply introducing the remote protecting group in this current series. The stereochemical outcome of this cyclisation reaction can be rationalised using the same conformational model as previously proposed by our group for related cyclisations.<sup>6</sup> The presence of an asymmetric centre next to the iminium carbon may also act to influence the formation of the new chiral centre on cyclisation. If this were to be a contributing factor, one can appreciate that the aromatic ring approaches the planar acyliminium intermediate from the direction of least steric hindrance.

Removal of the pendant hydroxymethyl substituent (auxiliary) from the tetracyclic product **6** was achieved by application of a three-step procedure (Scheme 3). Dess–Martin periodinane oxidation of the primary alcohol proceeded in 93% yield to provide aldehyde **8**. We then employed our favoured Rh-catalysed decarbonylation protocol to access enamide **9** in 55% yield; a small amount of target compound **10** (6%) was also produced directly in this reaction. The auxiliary removal





**11**,  $[\alpha]_D = -26.1$  (*c* = 1.15, CHCl<sub>3</sub>)

Scheme 3. Reagents and conditions: (i) Dess–Martin periodinane, DCM; (ii) Rh(PPh<sub>3</sub>)<sub>2</sub>(CO)Cl, dppp, mesitylene,  $\Delta$ , 4 days; (iii) H<sub>2</sub>/10% Pd–C, EtOH; (iv) ethylene glycol, *p*TSA (cat.), toluene,  $\Delta$ .

sequence was completed by catalytic hydrogenation of the crude mixture from the previous step to furnish the amidoketone **10** in 79% yield.

The formal asymmetric synthesis of (+)-demethoxyerythratidinone, the natural enantiomer, simply required re-protection of the ketone functional group using ethylene glycol as shown in Scheme 3. Compound **11** has been converted by others to the natural product albeit in racemic form by a four-step sequence.<sup>4</sup>



The formal asymmetric synthesis of the unnatural enantiomer followed an identical synthetic route, beginning however with the enantiomeric form of the original amino acid substrate to access the required lactam substrate 12, which had the opposite optical rotation value to that obtained for lactam 5. Intermediate 13 gave an optical rotation of +39.6, compared to -36.9 for compound 10. The *enantio*-target 14, gave an optical rotation of +21.7, compared to -26.1 for compound 11.

To summarise, we report the first synthetic application of our recently developed asymmetric N-acyliminium cyclisation methodology. Our approach allows a facile and highly stereoselective formal asymmetric synthesis of both enantiomers of the alkaloid 3-demethoxyerythratidinone.

## Acknowledgements

Loughborough University, Biofocus Ltd (joint studentship to G.B.S) and GlaxoSmithKline Pharmaceuticals (joint studentship to S.L.J).

## **References and notes**

- Tanaka, H.; Tanaka, T.; Etoh, H.; Goto, S.; Terada, Y. *Heterocycles* 1999, 51, 2759–2764; Dyke, S. F.; Quessy, S. N. *The Alkaloids*; Rodrigo, R. G. A. Academic Press: New York, 1981; Vol. 18.
- 2. (a) Padwa, A.; Hennig, R.; Kappe, C. O.; Reger, T. S. J. Org. Chem. 1998, 63, 1144-1155, and references cited therein; (b) Rigby, J. H.; Deur, C.; Heeg, M. J. Tetrahedron Lett. 1999, 40, 6887–6890; (c) Rigby, J. H.; Hughes, R. C.; Heeg, M. J. J. Am. Chem. Soc. 1995, 117, 7834-7835; (d) Suda, Y. T.; Hosai, S.; Ishida, K.; Sangai, M. Chem. Pharm. Bull. 1994, 42, 204-213; (e) Lete, E.; Egiarte, A.; Sotomayor, N.; Vicente, T.; Villa, M.-J. Synlett 1993, 41-42; (f) Manteca, I.; Sotomayor, N.; Villa, M.-J.; Lete, E. Tetrahedron Lett. 1996, 37, 7841-7844; (g) Lee, Y. S.; Kang, D. W.; Lee, S. J.; Park, H. J. Org. Chem. 1995, 60, 7149-7152; (h) Lee, Y. S.; Kang, D. W.; Lee, S. J.; Park, H. Synth. Commun. 1995, 25, 1947–1956; (i) Lee, J. Y.; Lee, Y. S.; Chung, B. Y.; Park, H. Tetrahedron 1997, 53, 2449-2458; (j) Katritzky, A. R.; Mehta, S.; He, H.-Y. J. Org. Chem. 2001, 66, 148-152; (k) Garcia, E.; Arrasate, S.; Ardeo, A.; Lete, E.; Sotomayor, N. Tetrahedron Lett. 2001, 42, 1511-1513.
- Barton, D. H. R.; Gunatilaka, A. A. L.; Letcher, R. M.; Lobo, A. M. F. T.; Widdowson, D. A. J. Chem. Soc., Perkin Trans. 1 1973, 874–880.
- Tsuda, Y.; Nakai, A.; Ito, K.; Suzuki, F.; Haruna, M. *Heterocycles* 1984, 22, 1817–1820.
- Hosoi, S.; Ishida, K.; Tsuda, Y. Chem. Pharm. Bull. 1992, 40, 3115–3117.
- (a) Allin, S. M.; James, S. L.; Martin, W. P.; Smith, T. A. D. J. Org. Chem. 2002, 67, 9464–9467; (b) Allin, S. M.; Thomas,

C. I.; Allard, J. E.; Duncton, M.; Elsegood, M. R. J.; Edgar, M. *Tetrahedron Lett.* **2002**, *44*, 2335–2337; (c) Allin, S. M.; Vaidya, D. G.; James, S. L.; Allard, J. E.; Smith, T. A. D.; McKee, V.; Martin, W. P. *Tetrahedron Lett.* **2002**, *43*, 3661– 3663; (d) Allin, S. M.; James, S. L.; Martin, W. P.; Smith, T. A. D.; Elsegood, M. R. J. *J. Chem. Soc., Perkin Trans. 1* **2001**, 3029–3036; (e) Allin, S. M.; James, S. L.; Martin, W. P.; Smith, T. A. D. *Tetrahedron Lett.* **2001**, *41*, 3943–3946; (f) Allin, S. M.; Northfield, C. J.; Page, M. I.; Slawin, A. M. Z. *Tetrahedron Lett.* **1998**, *39*, 4905–4908.

- (a) Meyers, A. I.; Brengal, G. P. Chem. Commun. 1997, 1–8; (b) Groaning, M. D.; Meyers, A. I. Tetrahedron 2000, 56, 9843–9873.
- Ragan, J. A.; Claffey, M. C. *Heterocycles* 1995, *41*, 57–70; Ennis, M. D.; Hoffman, R. L.; Ghazal, N. B.; Old, D. W.; Mooney, P. A. J. Org. Chem. 1996, *61*, 5813–5817.