

generation of octanethiol was conducted in CH₂Cl₂ at room temperature in the presence of Et₃N (eq 1).




Table 1 shows various rhodium(I) and iridium(I) catalysts (5% based on thiols) that we examined for their ability to

Table 1. Screening of Catalysts for Reaction of Octanethiol with Dichloromethane^a

entry	catalyst	yield (%) ^b
1 ^c	[Rh(cod) ₂]BF ₄ /2 PPh ₃	33
2 ^c	[Rh(cod) ₂]BF ₄ /2 <i>n</i> -Bu ₃ P	3
3 ^c	[Rh(cod) ₂]BF ₄ /dppe	5
4 ^c	[Rh(cod) ₂]BF ₄ /dppb	1
5 ^c	[Rh(cod) ₂]BF ₄ /dcpe	3
6 ^c	[Rh(cod) ₂]BF ₄ /dppf	24
7 ^c	[Rh(cod) ₂]BF ₄ /Tol-BINAP	8
8 ^c	[Rh(cod) ₂]BF ₄ /XANTPHOS ^d	17
9 ^c	[Ir(cod) ₂]BF ₄ /2 PPh ₃	<1
10 ^c	[Rh(cod)Cl] ₂ /4 PPh ₃	<1
11	[Rh(cod) ₂]BF ₄ /2 PPh ₃	<1
12	RhCl(PPh ₃) ₃	39
13 ^e	RhCl(PPh ₃) ₃	81 ^f
14	none	0

^a Rh or Ir catalyst (0.010 mmol), phosphine (0–0.020 mmol), Et₃N (0.22 mmol), thiol (0.20 mmol), and CH₂Cl₂ (1.0 mL) were employed. ^b Determined by ¹H NMR. ^c The catalysts were treated with hydrogen (1 atm, rt, 0.5 h). ^d 9,9-Dimethyl-4,5-bis(diphenylphosphino)xanthene. ^e RhCl(PPh₃)₃ (0.025 mmol), Et₃N (0.5 mL), thiol (0.50 mmol), and CH₂Cl₂ (2.0 mL) were employed. Reaction time: 24 h. ^f Isolated yield.

facilitate this transformation. Among the phosphine ligands

(6) For recent reports concerning transition-metal-catalyzed synthesis of sulfur compounds, see: (a) Dong, C.; Alper, H. *Org. Lett.* **2004**, 6, 3489. (b) Savarin, C.; Srogl, J.; Liebeskind, L. S. *Org. Lett.* **2002**, 4, 4309. (c) Knapton, D. J.; Meyer, T. Y. *Org. Lett.* **2004**, 6, 687. (d) Hirai, T.; Kuniyasu, H.; Kambe, N. *Tetrahedron Lett.* **2005**, 46, 117. (e) Hirai, T.; Kuniyasu, H.; Kambe, N. *Chem. Lett.* **2004**, 33, 1148. (f) Sugoh, K.; Kuniyasu, H.; Sugae, T.; Ohtaka, A.; Takai, Y.; Tanaka, A.; Machino, C.; Kambe, N.; Kurosawa, H. *J. Am. Chem. Soc.* **2001**, 123, 5108. (g) Kuniyasu, H.; Ohtaka, A.; Nakazono, T.; Kinomoto, M.; Kurosawa, H. *J. Am. Chem. Soc.* **2000**, 122, 2375. (h) Arisawa, M.; Yamaguchi, M. *J. Am. Chem. Soc.* **2003**, 125, 6624. (i) Arisawa, M.; Suwa, A.; Fujimoto, K.; Yamaguchi, M. *Adv. Synth. Catal.* **2003**, 345, 560. (j) Arisawa, M.; Yamaguchi, M. *Org. Lett.* **2001**, 3, 763. (k) Kondo, T.; Baba, A.; Nishi, Y.; Mitsudo, T. *Tetrahedron Lett.* **2004**, 45, 1469. (l) Hua, R.; Takeda, H.; Onozawa, S.; Abe, Y.; Tanaka, M. *J. Am. Chem. Soc.* **2001**, 123, 2899. (m) Taniguchi, N. *J. Org. Chem.* **2004**, 69, 6904. (n) Taniguchi, N.; Onami, T. *J. Org. Chem.* **2004**, 69, 915. (o) Ananikov, V. P.; Beletskaya, I. P. *Org. Biomol. Chem.* **2004**, 2, 284. (p) Tanaka, K.; Ajiki, K. *Tetrahedron Lett.* **2004**, 45, 5677 and references therein.

(7) (a) Page, P. C. B.; Klair, S. S.; Brown, M. P.; Smith, C. S.; Maginn, S. J.; Mulley, S. *Tetrahedron* **1992**, 48, 5933. (b) Page, P. C. B.; Klair, S. S.; Brown, M. P.; Harding, M. M.; Smith, C. S.; Maginn, S. J.; Mulley, S. *Tetrahedron Lett.* **1988**, 29, 4477.

(8) Tanaka, K.; Ajiki, K. *Tetrahedron Lett.* **2004**, 45, 25.

examined, PPh₃ was the most effective (entries 1–8). The use of [Ir(cod)₂]BF₄ or [Rh(cod)Cl]₂ and the elimination of treatment of the catalyst with hydrogen engendered very low catalytic activity (entries 9–11). The highest catalytic activity was obtained using RhCl(PPh₃)₃ as a catalyst (entry 12). Finally, employing excess amounts of Et₃N furnished the desired methylenedithioether in 81% isolated yield (entry 13).⁹ No reaction was observed in the absence of rhodium catalyst (entry 14).

Table 2 shows the rhodium-catalyzed reactions of various thiols with polychloroalkanes, as investigated in polychloroalkane solvents.¹⁰ The reactions of both alkyl and aryl

Table 2. Rhodium-Catalyzed Reaction of Thiols with Polychloroalkanes^a

entry	thiol	polychloroalkane	product	yield (%) ^b
1	CH ₃ (CH ₂) ₇ SH	CH ₂ Cl ₂	$\begin{array}{c} \text{S(CH}_2)_7\text{CH}_3 \\ \text{S(CH}_2)_7\text{CH}_3 \end{array}$	81
2	CH ₃ (CH ₂) ₁₁ SH	CH ₂ Cl ₂	$\begin{array}{c} \text{S(CH}_2)_{11}\text{CH}_3 \\ \text{S(CH}_2)_{11}\text{CH}_3 \end{array}$	80
3	PhCH ₂ SH	CH ₂ Cl ₂	$\begin{array}{c} \text{SCH}_2\text{Ph} \\ \text{SCH}_2\text{Ph} \end{array}$	82
4	HO(CH ₂) ₁₁ SH	CH ₂ Cl ₂	$\begin{array}{c} \text{S(CH}_2)_{11}\text{OH} \\ \text{S(CH}_2)_{11}\text{OH} \end{array}$	63
5	$\text{MeO}_2\text{C}-\text{CH}(\text{NH}^+\text{Boc})-\text{CH}_2\text{SH}$	CH ₂ Cl ₂	$\left(\text{MeO}_2\text{C}-\text{CH}(\text{NH}^+\text{Boc})-\text{CH}_2\text{S} \right)_2$	73
6	<i>p</i> -TolSH	CH ₂ Cl ₂	$\begin{array}{c} \text{S}(p\text{-Tol}) \\ \text{S}(p\text{-Tol}) \end{array}$	99
7 ^c	CH ₃ (CH ₂) ₇ SH	CH ₂ ClCH ₂ Cl	$\begin{array}{c} \text{S(CH}_2)_7\text{CH}_3 \\ \text{S(CH}_2)_7\text{CH}_3 \end{array}$	84
8 ^c	<i>p</i> -TolSH	CH ₂ ClCH ₂ Cl	$\begin{array}{c} \text{S}(p\text{-Tol}) \\ \text{S}(p\text{-Tol}) \end{array}$	87
9 ^d	CH ₃ (CH ₂) ₁₁ SH	CHCl ₃	$\begin{array}{c} \text{S(CH}_2)_{11}\text{CH}_3 \\ \text{O} \\ \text{H} \end{array}$	69
10 ^d	CH ₃ (CH ₂) ₅ SH	CCl ₄	$\begin{array}{c} \text{S(CH}_2)_5\text{CH}_3 \\ \text{O} \\ \text{S(CH}_2)_5\text{CH}_3 \end{array}$	61

^a Reactions were conducted using RhCl(PPh₃)₃ (0.025 mmol), Et₃N (0.5 mL), thiol (0.50 mmol), and polychloroalkane (2.0 mL) at room temperature for 24 h. ^b Isolated yields based on thiols. ^c At 80 °C. ^d RhCl(PPh₃)₃ (0.10 mmol) and polychloroalkane (10 mL) were used.


thiols with CH₂Cl₂ in the presence of 5% RhCl(PPh₃)₃ proceeded at room temperature to furnish the corresponding formaldehyde dithioacetals in good yield (entries 1–6). A hydroxy-substituted alkanethiol and a cysteine derivative can

(9) Not only tertiary amines but also secondary and primary amines (*n*-Bu₂NH and *n*-BuNH₂) can be used for this reaction, although the reaction rate is low.

(10) In all entries, <5% conversions were observed in the absence of rhodium catalyst.


be used (entries 4 and 5).¹¹ The reaction of a vicinal dichloride ($\text{CH}_2\text{ClCH}_2\text{Cl}$) proceeded at 80 °C to furnish the corresponding ethylenedithioethers in good yield (entries 7 and 8). Importantly, although high catalyst loading and diluted condition were required, the reaction of CHCl_3 and CCl_4 also proceeded at room temperature to give a thioformate and a dithiocarbonic ester, respectively, presumably through hydrolysis of corresponding polythiomethanes by silica gel chromatography (entries 9 and 10). The reaction of CHCl_3 and CCl_4 serves as a convenient new method for preparation of a thioformate and a dithiocarbonic ester starting from a thiol.

Next, we investigated the rhodium-catalyzed reaction of thiols with alkyl halides. The reaction proceeded cleanly in Et_3N without using excess alkylating reagents when a reactive primary alkyl chloride (benzyl chloride, eq 2) or a primary alkyl bromide (dodecyl bromide, eq 3) was used as an alkylating reagent. No reaction was observed in the absence of rhodium catalyst.




The reaction of dodecanethiol with CDCl_3 was investigated to gain mechanistic insight into this reaction (eq 4). Deuterium of CDCl_3 was quantitatively incorporated into the formyl group. Furthermore, the reaction of benzyl mercaptan with (*R*)-(1-chloroethyl)benzene (78% ee) furnished the corresponding sulfide with complete inversion of configuration (78% ee, eq 5).

Scheme 1 depicts a plausible mechanism of this reaction. We believe that the rhodium(I) catalyst oxidatively inserts



into the thiol S–H bond, affording a rhodium(III) complex **A**. Elimination of HX with Et_3N furnishes rhodium(I) thiolate **B**, which reacted with alkyl halides in an $\text{S}_\text{N}2$ fashion, thereby furnishing sulfides and regenerating the rhodium(I) catalyst.

Scheme 1



In conclusion, we have established that $\text{RhCl}(\text{PPh}_3)_3$ catalyzes a reaction of thiols with polychloroalkanes in the presence of triethylamine. This reaction serves as a convenient new method to produce formaldehyde dithioacetals, ethylenedithioethers, thioformates, and dithiocarbonic esters under mild conditions.

Acknowledgment. This research was supported by Banyu Pharmaceutical Company Award in Synthetic Organic Chemistry, Japan. We thank Dr. M. Hirano for elemental analysis.

Supporting Information Available: Experimental procedures and compound characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0501673

(11) For synthesis of methylenedithioethers from cysteine derivatives using tetrabutylammonium fluoride hydrate, see: Ueki, M.; Ikeo, T.; Hokari, K.; Nakamura, K.; Saeki, A.; Komatsu, H. *Bull. Chem. Soc. Jpn.* **1999**, *72*, 829.