

m-Iodosylbenzoic acid, a tagged hypervalent iodine reagent for the iodo-functionalization of alkenes and alkynes

Mekhman S. Yusubov^{a,*}, Roza Ya. Yusubova^a, Andreas Kirschning^{b,*},
Joo Yeon Park^c, Ki-Whan Chi^{c,*}

^a The Siberian State Medical University, 2 Moskovsky Trakt, 634050 Tomsk, Russia and The Tomsk Polytechnic University, 30 Lenin Street, 634050 Tomsk, Russia

^b Institut für Organische Chemie und Zentrum für Biomolekulare Wirkstoffe (BMWZ), Leibniz Universität Hannover, Schneiderberg 1B, D-30167 Hannover, Germany

^c University of Ulsan, 680-749 Ulsan, Republic of Korea

Received 20 August 2007; revised 14 December 2007; accepted 21 December 2007
Available online 8 January 2008

Abstract

An efficient and facile method for the iodo-functionalization of alkenes **5** and alkynes **6** by using recyclable *m*-iodosylbenzoic acid (**2**) was developed. The final products can be easily isolated without any chromatographic purification by simple treatment of the crude mixture with an anionic exchange resin. Unreacted *m*-iodosylbenzoic acid and reduced *m*-iodobenzoic acid are effectively recovered from the resin by acidification with hydrochloric acid.

© 2008 Elsevier Ltd. All rights reserved.

Keywords: Alkenes; Alkynes; Hypervalent iodine; Iodination; *m*-Iodosobenzoic acid; Scavenger

Iodo-functionalization of alkenes and alkynes has been used as a pivotal transformation in organic synthesis.^{1a,b} In particular, hypervalent iodine compounds have been successfully applied for iodination, iodo-functionalization, and oxidation reactions due to their mild and selective oxidizing characteristics.^{2a–f} For example, Yusubov et al.^{2g} recently demonstrated an effective iodo-functionalization of alkenes by using (dichloroiodo)benzene. Broader application of hypervalent iodine compounds, however, has been restricted due to the tedious purification and recovery process necessary following reactions. To circumvent this problem, several recyclable hypervalent iodine reagents have been developed in the form of polymer-supported^{3a–y} or molecular species.^{4a–i} Nevertheless, multi-step syntheses and subsequent separation are required for the preparation of these recyclable hypervalent reagents.

We have recently developed a straightforward approach to simplify the work-up of iodine(III)-mediated reactions by using 3-(dichloroiodo)benzoic acid^{4c} (**1**) and *m*-iodosylbenzoic acid^{5a,b} (**2**) as recyclable reagents (Fig. 1). A detailed spectroscopic investigation of *m*-iodosylbenzoic acid and its sodium salt by Katritzky et al. supported the polymeric structure of these compounds.^{5a,c} Both hypervalent iodine reagents are prepared with ease and tagged with a carboxylic acid group so that reduced iodine by-products such as *m*-iodobenzoic acid (**3**) can be removed at the end of the reaction by simple treatment with an anion exchange resin (IRA 900; hydroxide form) or by the




Fig. 1.

* Corresponding authors.


E-mail address: andreas.kirschning@oci.uni-hannover.de (A. Kirschning).

was applied instead of **2**, only the ketal **9e** of α,α -diiodoacetophenone was obtained from phenylethyne (**6e**). In other reactions with (dichloroiodo)benzene, the tentative products of ketals **9a–d,f** decomposed once flash chromatography on silica gel was attempted. These results clearly demonstrate the usefulness and preference of tagged *m*-iodosylbenzoic acid (**2**) over other iodine(III) reagents.^{2g}


Reaction of pent-4-yn-1-ol (**6d**) with iodine and **2** in MeOH provided 2-(diiodomethyl)-2-methoxytetrahydrofuran (**9d**). It is plausible to suggest that the iodocyclization of **6d** firstly yields cyclized intermediate **10** followed by iodo-methoxylation of the exocyclic double bond to produce the final product **9d** (Scheme 2).

Based on the results^{5b} of iodine(III)-mediated iodination of arenes, we suggest that the methoxy or hydrated form of **2** oxidizes iodine to MeOI (in MeOH) or HOI (in H₂O) which serves as a reactive electrophilic intermediate (Scheme 3).

In conclusion, the inexpensive and easy-to-handle hypervalent iodine reagent *m*-iodosylbenzoic acid (**2**) was successfully applied for the iodo-methoxylation and iodo-hydroxylation of alkenes and alkynes under mild conditions. This tagged reagent and the aryl iodide by-product



Scheme 2.



Scheme 3.

are easily removed from the reaction mixture by simple treatment with an anion exchange resin. This methodology allows easier preparation and simplified purification of organoiodo compounds.

Acknowledgments

This work was funded by the Fonds der Chemischen Industrie. In addition, K.W.C., M.S.Y. and R.Y.Y. are grateful for the support of Brain Korea 21 program in Korea.

References and notes

- (a) Smith, M. B.; March, J. *Advanced Organic Chemistry*, 5th ed.; Wiley: New York, 2001; pp 970–1171; (b) Carey, K.; Sundberg, R. *Advanced Organic Chemistry, Part B*, 4th ed.; Plenum: New York, 2001; pp 191–248.
- (a) Wirth, T. *Hypervalent Iodine Chemistry*; Springer: Berlin, 2003; pp 1–274; (b) Varvoglis, A. *Hypervalent Iodine in Organic Synthesis*; Academic Press: London, 1997; pp 1–223; (c) Tohma, H.; Kita, Y. *Adv. Synth. Catal.* **2004**, *346*, 111–124; (d) Wirth, T. *Angew. Chem., Int. Ed.* **2005**, *44*, 3656–3665; (e) Zhdankin, V. V.; Stang, P. J. *J. Chem. Rev.* **2002**, *102*, 2523–2584; (f) Zhdankin, V. V. *Hypervalent Iodoarenes and Arene Iodonium Salts*. In *Science of Synthesis*, 2007; Vol. 31, Chapter 31.4.1; (g) Yusubov, M. S.; Yusubova, R.; Filimonov, V. D.; Chi, K.-W. *Synth. Commun.* **2004**, *34*, 443–450.
- (a) Togo, H.; Sakuratani, K. *Synlett* **2002**, 1966–1975; (b) Togo, H. *Eco Ind.* **2002**, *7*, 5–18; (c) Gibietis, J.; Puce, A.; Zicmanis, A. *Latvijas PSR Zinatnu Akademijas Vestis Kim. Ser.* **1986**, 600–606; (d) Ley, S. V.; Thomas, A. W.; Finch, H. *J. Chem. Soc., Perkin Trans. 1* **1999**, 669–772; (e) Tohma, H.; Takizawa, S.; Maegawa, T.; Kita, Y. *Angew. Chem., Int. Ed.* **2000**, *39*, 1306–1308; (f) Chen, D.-J.; Chen, Z.-C. *Synlett* **2000**, 1175–1177; (g) Tohma, H.; Morioka, H.; Takizawa, S.; Arisawa, M.; Kita, Y. *Tetrahedron* **2001**, *57*, 345–352; (h) Ficht, S.; Mulbauer, M.; Giannis, A. *Tetrahedron* **2001**, *57*, 4863–4866; (i) Tohma, H.; Morioka, H.; Harayama, Y.; Hashizume, M.; Kita, Y. *Tetrahedron Lett.* **2001**, *42*, 6899–6902; (j) Huang, X.; Zhu, Q.; Zhang, J. *J. Chem. Res. Synop.* **2001**, 480–481; (k) Sorg, G.; Mengei, A.; Jung, G.; Rademann, J. *Angew. Chem., Int. Ed.* **2001**, *40*, 4395–4397; (l) Chen, D.-J.; Cheng, D.-P.; Chen, Z.-C. *Synth. Commun.* **2001**, *31*, 3847–3850; (m) Kawamura, Y.; Maruyama, M.; Tokuoka, T.; Tsukayama, M. *Synthesis* **2002**, 2490–2496; (n) Cheng, D.-P.; Chen, Z.-C. *Synth. Commun.* **2002**, *32*, 2155–2159; (o) Tohma, H.; Maegawa, T.; Takizawa, S.; Kita, Y. *Adv. Synth. Catal.* **2002**, *344*, 328–337; (p) Kirschning, A.; Kunst, E.; Ries, M.; Rose, L.; Schöninger, A.

Table 2
Iodomethoxylation of alkynes **6** with *m*-iodosylbenzoic acid **2**^{14,15}

Alkyne 6	Product 9	Yield ^a (%)
 6a	 9a	83
 6b	 9b	99
 6c	 9c	78
 6d	 9d	76
 6e	 9e	90
 6f	 9f	84

^a Isolated yield of pure products.

- Wartchow, R. *Arkivoc* **2003**, 145–163; (q) Sakuratani, K.; Togo, H. *Synthesis* **2003**, 21–23; (r) Tohma, H.; Maegawa, T.; Kita, Y. *Synlett* **2003**, 723–725; (s) Sakuratani, K.; Togo, H. *Arkivoc* **2003**, 11–20; (t) Tohma, H.; Maegawa, T.; Kita, Y. *Arkivoc* **2003**, 62–70; (u) Cheng, D.-P.; Chen, Z.-C.; Zheng, Q.-G. *Synth. Commun.* **2003**, *33*, 2671–2676; (v) Chen, J.-M.; Lin, X.-J.; Huang, X. *J. Chem. Res.* **2004**, 43–44; (w) Lee, J.-W.; Ko, K. Y. *Bull. Korean Chem. Soc.* **2004**, *25*, 19–20; (x) Teduka, T.; Togo, H. *Synlett* **2005**, 923–926; (y) Kalberer, E. W.; Whitfield, S. R.; Sanford, M. S. *J. Mol. Catal. A: Chem.* **2006**, *251*, 108–113.
4. (a) Thottumkara, A. P.; Vinod, T. K. *Tetrahedron Lett.* **2002**, *43*, 569–572; (b) Tohma, H.; Maruyama, A.; Maeda, A.; Maegawa, T.; Dohi, T.; Shiro, M.; Morita, T.; Kita, Y. *Angew. Chem., Int. Ed.* **2004**, *42*, 3595–3598; (c) Yusubov, M. S.; Drygunova, L. A.; Zhdankin, V. V. *Synthesis* **2004**, 2289–2292; (d) Dohi, T.; Maruyama, A.; Yoshimura, M.; Morimoto, K.; Tohma, H.; Shiro, M.; Kita, Y. *Chem. Commun.* **2005**, 2205–2207; (e) Dohi, T.; Morimoto, K.; Takenaga, N.; Maruyama, A.; Kita, Y. *Chem. Pharm. Bull.* **2006**, *54*, 1608–1610; (f) Dohi, T. *J. Pharm. Soc. Jpn.* **2006**, *126*, 757–766; (g) Moroda, A.; Togo, H. *Tetrahedron* **2006**, *62*, 12408–12414; (h) Dohi, T.; Morimoto, K.; Takenaga, N.; Goto, A.; Maruyama, A.; Kiyono, Y.; Tohma, H.; Kita, Y. *J. Org. Chem.* **2007**, *72*, 109–116; (i) Justik, M. W. *Tetrahedron Lett.* **2007**, *48*, 3003–3007.
 5. (a) Yusubov, M. S.; Gilmkhanova, M. P.; Zhdankin, V. V.; Kirschning, A. *Synlett* **2007**, 563–566; (b) Yusubov, M. S.; Yusubova, R. Ya.; Chi, K.-W.; Park, J. Y.; Kirschning, A. *Beilstein J. Org. Chem.* **2007**, *3*, 19; (c) Katritzky, A. R.; Savage, G. P.; Gallos, J. K.; Durst, H. D. *J. Chem. Soc., Perkin Trans. 2* **1990**, 1515–1518.
 6. Amberlite IRA 900 (chloride form) was obtained from Supelco.
 7. Iranpoor, N.; Shekarriz, M. *Tetrahedron* **2000**, *56*, 5209–5211.
 8. Cerritelli, S.; Chiarini, M.; Cerichelli, G.; Capone, M.; Marsili, M. *Eur. J. Org. Chem.* **2004**, 623–630.
 9. Asensio, G.; Andreu, C.; Boix-Bernardini, C.; Mello, R.; Gonzalez-Nunez, M.-E. *Org. Lett.* **1999**, 2125–2128.
 10. Davidson, R. I.; Kropp, P. J. *J. Org. Chem.* **1982**, *47*, 1904–1909.
 11. Mahajan, V. A.; Shinde, P. D.; Gajare, A. S.; Karthikeyan, M.; Wakharkar, R. D. *Green Chem.* **2002**, 325–327.
 12. This transformation has been reported before: (a) Sato, T.; Tamura, K.; Nagayoshi, K. *Chem. Lett.* **1983**, *5*, 791–794; (b) Davidson, R. I.; Kropp, P. J. *J. Org. Chem.* **1982**, *47*, 1904–1909; (c) Schauble, J. H.; Trauffer, E. A.; Deshpande, P. P.; Evans, R. D. *Synthesis* **2005**, 1333–1339.
 13. (a) Janas, J. J.; Asirvatham, E. T.; McNelis, E. *Oxid. Commun.* **1985/86**, *8*, 65; (b) Bovonsombat, P.; McNelis, E. *Tetrahedron Lett.* **1992**, *33*, 4123–4126.
 14. **Typical procedure for iodomethoxylation.** Alkenes **5a–c,e** (0.3 mmol) were added to a mixture of iodine (25.4 mg, 0.10 mmol) and *m*-iodosylbenzoic acid (**2**) (29 mg, 0.11 mmol) in MeOH (0.5 mL) and the reaction mixture was stirred at room temperature for 1 h (the reactions were monitored by TLC). Then, dichloromethane (1.0 mL) was added and the resulting solution was cooled to 5 °C. IRA 900 (350–400 mg, hydrocarbonate form) was added and the mixture was stirred for 5 min. The resin was removed by filtration and the solution was concentrated under reduced pressure to afford pure methoxyiodides **7a–c,e** as judged by NMR-spectroscopy. *m*-Iodobenzoic acid (**3**) can be easily regenerated from the IRA 900 resin **4** by treatment with aqueous HCl and reoxidized to reagent **2** without additional purification as described in Ref. 5a. For the iodomethoxylation of alkenes **5d,g–k**, a molar ratio of alkene:**2**:iodine = 0.2:0.11:0.2 (mmol) was employed, instead. The iodohydroxylation was carried out accordingly except that MeCN–H₂O (5:1, 0.5 mL) was employed as solvent instead of MeOH.
Typical procedure for diiododimethoxylation. To a solution of iodine (28 mg, 0.11 mmol) and *m*-iodosylbenzoic acid (**2**) (29 mg, 0.11 mmol) in MeOH (0.5 mL) were added alkynes **6a–f** (0.15 mmol) in MeOH (0.5 mL), and the reaction mixture was stirred at room temperature for 1 h. Then, dichloromethane (1.0 mL) was added and the resulting solution was cooled to 5 °C. IRA 900 (350–400 mg, hydroxide form) was added and the mixture was stirred for 5 min in the dark. The resin was removed by filtration and the solution was concentrated under reduced pressure to afford pure diiodoketals **9a–f** as judged by NMR-spectroscopy.
 15. Spectral data for **7k**, **8k** and new compounds. **2-Iodo-3-methoxy-3-phenylpropan-1-ol (7k)**: Oil: ¹H NMR: δ 2.89 (1H, OH, s), 3.27 (3H, OCH₃, s), 3.81 (1H_a, CH₂, m), 3.95 (1H_b, CH₂, m), 4.34 (1H, CH–OMe, m), 4.52 (1H, CH–I, m); IR (neat) ν 3356 (OH), 1123 and 1085 (C–OH and C–O–C), 762 and 690 (C–I) cm^{−1}.
2-Iodo-1-phenylpropane-1,3-diol (8k): Colorless solid: mp 84–86 °C; ¹H NMR: δ 2.72 (1H, OH, s), 3.14 (1H, OH, s), 3.82 (1H_a, CH₂, m), 3.96 (1H_b, CH₂, m), 4.44 (1H, CH–OMe, d), 5.04 (1H, CH–I, m); IR (neat) ν 3380 (OH), 1074 (C–O), 780 and 684 (C–I) cm^{−1}.
3-Iodo-6-methoxycyclohex-1-ene (7e): Oil: ¹H NMR: δ 2.00–2.08 (4H, CH₂, m), 3.34 (3H, OCH₃, s), 3.99 (1H, CH–OMe, m), 4.43 (1H, CH–I, m), 5.72 (1H, =CH, m), 5.93 (1H, =CH, m); ¹³C NMR: δ 25.4 (CH₃), 30.0 (CH₂), 30.1 (CH₂), 57.0 (OCH₃), 80.6 (COMe), 124.1 (=C–COMe), 130.9 (=C–CI); IR (neat) ν 1125 (C–O–C), 926 (C=C), 644 (C–I) cm^{−1}. Anal. Calcd for C₇H₁₁IO: C, 35.32; H, 4.66. Found: C, 35.17; H, 4.73.
1,1-Diiodo-2,2-dimethoxyhexane (9a): Oil: ¹H NMR: δ 0.94 (3H, CH₃, t), 1.34–1.43 (4H, CH₂, m), 2.02 (2H, CH₂, t), 3.33 (6H, CH₃, s) 5.39 (1H, Cl₂H, s); ¹³C NMR: δ −15.10 (Cl₂), 12.2 (CH₃), 21.8 (CH₂), 26.4 (CH₂), 33.7 (CH₂), 50.4 (OCH₃), 98.8 (C(OMe)₂); IR (neat) ν 1075 and 1042 (C–O–C), 682 and 640 (C–I) cm^{−1}. Anal. Calcd for C₈H₁₆I₂O₂: C, 24.14; H, 4.05. Found: C, 23.89; H, 4.18.
4,4-Diiodo-5,5-dimethoxyoctane (9b): Oil: ¹H NMR: δ 0.97 (3H, CH₃, t), 1.05 (3H, CH₃, t), 1.54–1.62 (4H, CH₂, m), 2.03 (2H, CH₂–C(OMe)₂, t), 2.23 (2H, CH₂–Cl₂, t), 3.56 (6H, CH₃, s); IR (neat) ν 1095 and 1056 (C–O–C), 678 and 632 (C–I) cm^{−1}. Elemental analysis and ¹³C NMR could not be obtained for **9c** due to slow decomposition of the sample on storage.
(2,2-Diiodo-1,1-dimethoxyethyl)cyclohexane or 1,1-Diiodo-2,2-dimethoxyethyl-2-cyclohexylethane (9c): Colorless solid: mp 34–35 °C (decomp.); ¹H NMR: δ 1.12–1.26 (6H, CH₂, m), 1.66 (2H, CH₃, m), 1.77 (1H_a, CH₂, m), 1.89 (1H_b, CH₂, m), 2.28 (H, CH, m), 3.45 (6H, CH₃, s), 5.47 (1H, Cl₂, s); ¹³C NMR: δ −18.5 (CH₂), 26.3 (CH₂), 27.1 (CH₂), 29.2 (CH₂), 43.9 (CH), 50.6 (OCH₃), 51.8 (OCH₃), 97.1 (C(OMe)₂). Anal. Calcd for C₁₀H₁₈I₂O₂: C, 28.32; H, 4.28. Found: C, 28.70; H, 4.65.
2-Diiodomethyl-2-methoxytetrahydrofuran (9d): Oil: ¹H NMR: δ 2.05 (2H, CH₂, m), 2.35 (1H, CH₂, m), 2.44 (H, CH₂, m), 3.22 (6H, CH₃, s), 4.11 (2H, CH₂, m), 5.52 (1H, Cl₂H, s); ¹³C NMR: δ −20.1 (Cl₂), 25.8 (CH₂), 38.1 (CH₂), 48.2 (OCH₃), 71.5 (CH₂–O), 108.0 (O–C–OMe); IR (neat) ν 1050 and 1022 (C–O–C), 690 and 647 (C–I) cm^{−1}. Anal. Calcd for C₆H₁₀I₂O₂: C, 19.59; H, 2.74. Found: C, 19.44; H, 2.81.
1,1-Diiodo-2,2-dimethoxyethyl-2-phenylethane (9e): Colorless solid: mp 84–86 °C (decomp.); ¹H NMR: δ 3.32 (6H, OCH₃, s), 5.56 (1H, Cl₂H, s), 7.30–7.35 (3H_{arom.}, m) 7.62–7.67 (2H_{arom.}, m); ¹³C NMR: δ −19.4 (Cl₂), 49.0 (OCH₃), 97.4 (C(OMe)₂), 125.8, 127.3, 128.0, 133.7 (C_{arom.}); IR (KBr) ν 1089 and 1045 (C–O–C), 679 and 635 (C–I) cm^{−1}. Anal. Calcd for C₁₀H₁₂I₂O₂: C, 28.73; H, 2.89. Found: C, 28.86; H, 2.85.
2,2-Diiodo-1,1-dimethoxyethyl-1-phenylpropane (9f): Colorless solid: mp 65–66 °C (decomp.); ¹H NMR: δ 2.86 (3H, CH₃, s), 3.59 (6H, OCH₃, s), 7.37–7.40 (3H_{arom.}, m) 7.64–7.66 (2H_{arom.}, m); ¹³C NMR: δ 15.1 (CH₃), 45.3 (Cl₂), 53.89 (ICH₃), 102.3 (C(OMe)₂), 126.2, 128.0, 131.2, 132.5 (C_{arom.}); IR (KBr) ν 1068 and 1015 (C–O–C), 694 and 665 (C–I) cm^{−1}. Anal. Calcd for C₁₁H₁₄I₂O₂: C, 30.58; H, 3.27. Found: C, 30.71; H, 3.59.