Table I. Reactions of Vinylalanates withVarious Reagents in Ether

Vinylalanate derived from	Reagent	Product	Yield, $\%^{a,b}$
1-Hexyne		trans-2-Heptenoic acid	78
1 1100,010	HCHO ⁶	trans-2-Hepten-1-ol	73
	CH ₃ CHO	trans-3-Octen-2-ol	68 ^d
Cyclohexyl- acetylene	CO_2	trans-3-Cyclohexyl-2- propenoic acid	72
2-Butyne	CO_2	trans-2-Methyl-2-butenoic acid	76
3-Hexyne	CO2	trans-2-Ethyl-2-pentenoic acid	78

^a By isolation. ^b Yields are based on the amount of alkyne used. ^c Paraformaldehyde. ^d Yield 90% by glpc analysis.

Although the nature of the reactive organometallic intermediates has not been established, it is conceivable that the reactions proceed *via* the vinyllithium derivatives resulting from disproportionation of the vinylalanates. The configurational stability of vinyllithium

$$\begin{bmatrix} CH_{3} \\ H \\ R \end{bmatrix}^{-} C = C \begin{pmatrix} CH_{3} \\ H \\ H \end{pmatrix}^{-} Li^{+} \rightleftharpoons \frac{H}{R} C = C \begin{pmatrix} Li \\ H \end{pmatrix}^{-} Al(C_{4}H_{9})_{2}CH_{3}$$

compounds in solution, a requirement in view of the high stereospecifity of the above reactions, has been demonstrated by Seyferth and co-workers.⁵ We have observed that tetraalkylalanates do not react with carbon dioxide; however, their disproportionation into alkyllithium compounds and trialkylalanes should be thermodynamically unfavorable.

The synthesis of vinylalanes is not confined to the hydroalumination reaction. Thus trialkylalanes add to acetylene under mild conditions to yield *cis*-vinylalanes, whereas addition of the aluminum-carbon bond to 1-alkynes affords the corresponding disubstituted vinylalanes.² The availability of *cis*- and *trans*-mono-



substituted vinylalanes greatly increases the versatility of the above reactions.

The simplicity of the present procedure for the conversion of alkynes into α,β -unsaturated derivatives via vinylalanes is illustrated by the following example.

To 0.10 mole of 1-hexyne in 20 ml of *n*-heptane was added by means of a hypodermic syringe 0.10 mole of diisobutylaluminum hydride, while maintaining the temperature below 40°. After the initial exothermic reaction had subsided, the reaction mixture was heated for 2 hr at 50°. To the vinylalane formed was added at room temperature 0.10 mole of methyllithium in ether (60 ml). The solution was cooled to -30° , then carbonated by introducing a stream of carbon dioxide while maintaining the temperature at -30 to -10° . The reaction mixture was poured slowly into an iceconcentrated hydrochloric acid mixture and the carboxylic acid produced was extracted into ether. Distil-

(5) D. Seyferth, Record Chem. Progr. (Kresge-Hooker Sci. Lib.), 26, 86 (1965).

lation gave 10.0 g of *trans*-2-heptenoic acid (78%), bp 89° (0.3 mm), n^{22} D 1.4560, amide mp 124°.⁶

Acknowledgment. This work was supported by National Science Foundation Grants GP-3521 and GP-6633.

(6) P. Bruylants, Bull. Soc. Chim. Belges, 41, 333 (1932), reports for trans amide mp 125°.

(7) National Institutes of Health Predoctoral Fellow, 1965-1966.

George Zweifel, Roger B. Steele⁷ Department of Chemistry, University of California Davis, California 95616 Received April 12, 1967

Organonickel Compounds as Reagents for Selective Carbon-Carbon Bond Formation between Unlike Groups

Sir:

The extension of carbon chains by the coupling of two different groups using as reagents an organic halide, RHal, and an alkyl- or arylmetal derivative, R'Met, is seldom employed in synthetic practice if alternative operations are available. This is due in part to the fact that coupling products RR and R'R' are usually formed in substantial amounts¹ and in part to the intervention of a variety of other side reactions, *e.g.*, α - or β -elimination processes. We report here a new and promising method for the *selective combination of unlike groups* at carbon using organonickel reagents.

Reaction of a variety of allylic bromides with excess nickel carbonyl in dry benzene (50°, 2–3 hr) followed by removal of solvent and recrystallization from ether at $-70^{\circ 2}$ gives good yields of π -allylnickel(I) bromides, for example, **1a–c**, in 80–95% yield.^{3,4} These complexes are relatively inert toward alkyl halides in either hydro-



carbon solvents or ether-type solvents (e.g., tetrahydrofuran). In more *polar*, *coordinating media*, *e.g.*, dimethylformamide, N-methylpyrrolidone, or hexamethylphosphoramide, a facile reaction occurs between the complexes 1 and a *wide variety* of halides, with iodides generally being more reactive than (and preferable to) bromides.⁵ Equation 1 summarizes the over-all reaction, and Table I presents the results ob-

$$2R'Hal + 1 \longrightarrow 2R'CH_2C = CH_2$$
(1)

⁽¹⁾ For example, because of fast halogen-metal exchange prior to coupling. See D. E. Applequist and D. F. O'Brien, J. Am. Chem. Soc., **85**, 743 (1963).

⁽²⁾ All operations with organonickel compounds have been conducted with rigorous exclusion of oxygen using an argon atmosphere either with plastic glove bag or argon-line techniques.

⁽³⁾ The complex 1a, first prepared by E. O. Fischer and G. Bürger, Z. Naturforsch., 16b, 77 (1961), was originally obtained only in ca. 10% yield; however, much higher yields are easily realized if thermal or oxidative decomposition of the complexes is avoided.

⁽⁴⁾ These complexes can also be obtained efficiently from allylic bromides using bis(1,5-cyclooctadiene)nickel as reagent; see G. Wilke *et al.*, *Angew. Chem. Intern. Ed. Engl.*, **5**, 159 (1966). However, the synthesis from commercially available nickel carbonyl is usually more expeditious.

⁽⁵⁾ Chlorides are less generally useful because of their much lower reactivity.

Table I. Coupling of π -Methallylnickel(I) Bromide (1b) with Halides (Eq 1) in Dimethylformamide

Halide	Reaction time, hr (temp, °C)	Yield, %
Methyl iodide	10 (22)	90ª
Methyl bromide	19 (22)	90ª
Cyclohexyl iodide	3 (22)	91
4-Hydroxycyclohexyl iodide	10 (22)	880
t-Butyl iodide	24 (22)	25ª
Iodobenzene	1 (22)	98
Vinyl bromide	13 (22)	70ª
Benzyl bromide	6 (60)	91
3-Phenylpropyl bromide	46 (65)	92
Phenyl α -chloromethyl ether	18 (60)	50
p-Bromophenacyl bromide	0.5 (22)	75
Chloroacetone	2 (22)	46ª

^a These yields were determined by quantitative vpc analysis; others are yields of isolated pure products. ^b Mixture of cis and trans isomers which are converted by chromic acid oxidation to 4methallylcyclohexanone.

tained with 1b and a diverse collection of substrates.⁶ The reaction proceeds well not only with alkyl iodides and bromides but also with aryl and vinyl compounds. Further, the data indicate that the presence of carbonyl or hydroxyl functions in the halide partner need not interfere with coupling. Dihalides undergo disubstitution with the appropriate quantity of nickel(I) complex, as shown by the conversion of 1,6-diiodohexane with 1b (mole ratio 1:1.25) to 2,11-dimethyl-1,11-dodecadiene (95% yield) and of 1,4-dibromobenzene with 1b (mole ratio 1:1.25) to 1,4-dimethallylbenzene (97%). The latter substance is not readily accessible by conventional synthetic methods.

$$Br \longrightarrow CH_2 = CCH_2 \longrightarrow CH_2C = CH_2$$

The new method has been employed in direct and efficient syntheses of α -santalene (2)⁷ (88% yield) and epi- β -santalene (3)⁸ (90% yield) using the nickel(I) complex from α, α -dimethylallyl bromide as follows.⁹



epi- β -santalene

(6) The structures and products obtained by the reactions described herein have been verified by infrared, nmr, and mass spectrometric or elemental analysis. Purity was checked by vapor phase chromatography and quantitative nmr analysis

(7) E. J. Corey, S. W. Chow, and R. A. Scherrer, J. Am. Chem. Soc., **79**, 5773 (1957).

(8) E. J. Corey, R. Hartmann, and P. A. Vatakencherry, *ibid.*, 84, 2611 (1962). The iodocamphene used in the synthesis of 3 was prepared from the corresponding bromide by successive treatment with magnesium, mercuric chloride, and iodine. The bromocamphene was derived from π -bromocamphor by reduction (BH₄-) and dehydration with POCl₃-pyridine.

(9) The nickel(I) π complex from α, α -dimethylallyl bromide generally undergoes preferential coupling at the primary rather than at the

Substances of utility for further chain extension or cyclization can also be produced, for instance

Etooc
$$Ni_2$$
^{Br} + I(CH₂)₃Cl CH_2 = C(CH₂)₄Cl
 $|$ COOEt

In cases where substitution at C_1 and/or C_3 of the allyl group allows the possibility of geometrically isomeric coupling products, both *cis* and *trans* isomers usually result. For example, the π complex prepared from trans-geranyl bromide and nickel carbonyl affords with cyclohexyl iodide the coupling product 4 as a mixture of *cis-trans* isomers (40 and 60%).¹⁰

$$(CH_3)_2C = CHCH_2CH_2C = CHCH_2$$

 π -Allylnickel(I) complexes also can react with substances other than halides. The methallyl complex 1b reacts with benzaldehyde, acrolein, and cyclopentanone in dimethylformamide to yield 5, $R = C_6 H_5$ (85%), 5, R = CH=CH₂ (80\%), and 6 (50\%), respectively, and styrene oxide affords 7 (60%). In general, these reactions are considerably slower than those of 1b with iodides, and more drastic conditions were used (50°, ca. 24 hr).



The more general problem of effecting cross-coupling reactions of unlike *nonallylic* groups is currently the subject of a parallel investigation in this laboratory, not only in the nickel series but also with other transition metal derivatives, for example, organocopper¹¹⁻¹³ and organosilver^{14–17} compounds.

An especially crucial part of the coupling process described above is the step in which the new carboncarbon bond is formed and in particular the precursor for that step. Among the possibilities which presently appear as reasonable is the complex 8 which might be formed, for example, by the route¹⁸

tertiary terminal; with methyl iodide it affords pure 2-methyl-2-pentene (by vpc and nmr analysis) in ca. 90% yield.

(10) The interconversion of cis-trans forms of the allyl group could occur within the nickel complexes, e.g., via reversible $\sigma-\pi$ bonding changes. See (a) J. Powell, S. D. Robinson, and B. L. Shaw, Chem. Commun, 78 (1965); (b) K. G. Ramey and G. L. Statton, J. Am. Chem. Soc., 88, 4387 (1966). Alternatively, isomerization may involve equilibration of primary and tertiary allylic bromides via nickel(I) σ and π complexes.

(11) H. Gilman and J. M. Straley, Rec. Trav. Chim., 55, 821 (1936). (12) R. G. R. Bacon and H. A. O. Hill, Quart. Rev. (London), 19, 95

(1965). (13) C. E. Castro, et al., J. Org. Chem., 28, 2163, 3313 (1963); 31, 4071 (1966).

(14) E. A. Bickley and J. H. Gardner, ibid., 5, 126 (1940).

(15) G. M. Whitesides and C. P. Casey, J. Am. Chem. Soc., 88, 4541 (1966).

(16) G. Köbrich, H. Frölich, and W. Drischel, J. Organometal. Chem. (Amsterdam), 6, 194 (1966). (17) H. C. Brown and C. H. Snyder, J. Am. Chem. Soc., 83, 1002

(1961).

(18) Experiments in progress have shown that $bis-\pi$ -allylnickel(0) is unreactive toward bromobenzene and less reactive toward cyclohexyl iodide than is π -allylnickel(I) bromide (unpublished work with L. S. Hegedus and H. A. Kirst). Consequently, 8 is considered a more likely intermediate than analogs which are derived from reaction of nickel(0) complexes with RX.



L=e.g., DMF

The behavior of allylic halides toward π -allylnickel(I) derivatives, which contrasts sharply with that of nonallylic halides such as those appearing in Table I, will be detailed in due course.19

(19) This work was supported by National Science Foundation Grant GP-6527 X, and also by National Science Foundation Fellowships to M. F. S.

> E. J. Corey, Martin F. Semmelhack Department of Chemistry, Harvard University Cambridge, Massachusetts 02138 Received March 13, 1967

The Synthesis of Large-Ring 1.5-Dienes by Cyclization of Allylic Dibromides with Nickel Carbonyl

Sir:

The method for forming cycloolefins from allylic dihalides and nickel carbonyl which has recently been reported^{1,2} has now been examined in order to ascertain its scope with a series of dibromides of structure 1. The results which have been obtained suggest that this

BrCH₂CH=CH(CH₂)_nCH=CHCH₂Br

$$1$$

HOCH₂C=C(CH₂)_nC=CCH₂OH

cyclization process provides an unusually efficient route for the formation of large rings. In addition, because it leads to cyclic 1,5-dienes, it makes available a wide variety of cyclic structures which are not obtainable in a practical way via the acyloin reaction, currently the only commonly used general approach to large rings.

The dibromides 1, n = 2, 4, 6, 8, and 12, were synthesized in both cis, cis and trans, trans forms from the acetylenic diols 2, n = 2, 4, 6, 8, and 12, by selective reduction to the corresponding cis, cis-3 or trans,trans-ethylenic⁴ diols⁵ followed by reaction with phosphorus tribromide.6 The required acetylenic diols 2, n = 4, 6, and 8, were made by alkylation of a dibromide, $Br(CH_2)_n Br$, with the sodio derivative of propargyl alcohol tetrahydropyranyl ether⁷ in liquid ammonia-ether⁸ (n = 4, 6, 8) or in tetrahydrofuran-

(7) R. G. Jones and M. J. Mann, J. Am. Chem. Soc., 75, 4048 (1953).

dimethyl sulfoxide (n = 12), and the diol 2, n = 2, was obtained from 1,5-hexadiyne9 and formaldehyde.10

The cyclization experiments were conducted by slow addition (motor-driven syringe, 12-hr period of addition) of the allylic dibromide (1 mmole) in dimethylformamide (1 ml) to a solution of nickel carbonyl (5 mmoles) in dimethylformamide (ca. 40 ml) under argon at 50°. In the cases of the dibromides 1, n = 6, 8, and 12, the same cyclization product was obtained starting with either cis, cis or trans, trans isomer.



In each of these cases the cyclization product was principally (95-98%) the *trans.trans*-1.5-diene, but small amounts of an isomeric compound, probably the cis, trans isomer, could be detected by vapor phase chromatography.¹¹ The major product from 1, n =6. was identified as trans, trans-1,5-cyclododecadiene by spectroscopic comparison with the known substance;¹² the infrared spectrum manifested absorption at 10.4 μ due to CH=CH (*trans*) and no absorption at ca. 14.2 μ which would be expected for CH=CH (cis), and hydrogenation afforded cyclododecane, identical with an authentic sample. The principal products from 1, n = 8, and 1, n = 12, were identified by their infrared spectra (e.g., absorption at 10.4 μ but not at 14.2 μ), mass spectra, nmr spectra, and hydrogenation to cyclotetradecane, mp $54-55^{\circ}$, ¹³ and cyclooctadecane, mp $71-72^{\circ}$, ¹³ respectively. The predominant course of cyclization with the dibromides 1, n = 2, and 1, n =4, was formation of six-membered ring structures by 1-6 and 3-8 coupling, respectively; again the product composition was essentially independent of the geometry of the starting dibromide. With nickel carbonyl, 1, n = 2, afforded 4-vinylcyclohexene (42%) and cis,cis-1,5-cyclooctadiene (5%); the same products were also obtained using triphenylphosphinenickel tricarbonyl,¹⁴ but the relative amount of 1,5-cyclooctadiene was somewhat greater (ca. 20% of the mixture). In the case of dibromide 1, n = 4, with either nickel carbonyl or triphenylphosphinenickel tricarbonyl, only

- (10) F. Sondheimer, J. Am. Chem. Soc., 74, 4040 (1952).
- (11) Using a column packed with Carbowax 20M (10%) on Chromosorb P at 200°.
- (12) Reference spectra were furnished by Dr. Masaji Ohno, Toyo Rayon Co., Kamakura, Japan.
- (13) L. Ruzicka, M. Stoll, M. W. Huyser, and H. A. Boekenoogen. Helv. Chim. Acta, 13, 1152 (1930).
- (14) See ref 2 for an example of ligand control of product using this reagent.

⁽¹⁾ E. J. Corey and E. Hamanaka, J. Am. Chem. Soc., 86, 1641 (1964).

⁽²⁾ E. J. Corey and M. F. Semmelhack, Tetrahedron Letters, 6237 (1966).

⁽³⁾ Using Lindlar catalyst: see H. Lindlar and R. Dubuis, Org. Syn., 46, 89 (1966).

⁽⁴⁾ Using lithium aluminum hydride in tetrahydrofuran; see K. R. Bharucha and B. C. L. Weedon, J. Chem. Soc., 1584 (1953).

⁽⁵⁾ Satisfactory analytical and spectroscopic (infrared and nmr) data have been obtained for the new compounds reported herein. The stereochemical assignments to the cis, cis- and trans, trans-dibromides 1 and the corresponding diols rest not only on the methods of synthesis (previously shown to be stereospecific) but also on the occurrence of infrared absorption at 10.4 or 10.3 μ for the *trans,trans*-dihalides or diols and at 13.1-13.4 μ for all the *cis,cis*-dibromides. (6) J. M. Osbond, J. Chem. Soc., 5270 (1961).

⁽⁸⁾ H. Bader, L. C. Cross, I. Heilbron, and E. R. H. Jones, J. Chem. Soc., 619 (1949).
 (9) R. A. Raphael and F. Sondheimer, *ibid.*, 120 (1950).