

Published on Web 02/02/2008

Organocatalytic Enantioselective Mannich-Type Reaction of Phosphorus Ylides: Synthesis of Chiral *N*-Boc- β -Amino- α -methylene Carboxylic Esters

Yan Zhang,[†] Yan-Kai Liu,[†] Tai-Ran Kang,[†] Ze-Kai Hu,[†] and Ying-Chun Chen^{*,†,‡}

Key Laboratory of Drug-Targeting of Education Ministry and Department of Medicinal Chemistry, West China School of Pharmacy, Sichuan University, Chengdu 610041, China, and State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Chengdu, China

Received December 30, 2007; E-mail: ycchenhuaxi@yahoo.com.cn

 β -Amino carbonyl compounds containing an α -alkylidene group are highly densely functionalized materials which are widely applied in the synthesis of medicinal reagents and natural products.¹ They are usually prepared through the classic aza-Morita–Baylis– Hillman (MBH) reaction of activated imines and electron-deficient alkenes catalyzed by tertiary amines or phosphines.² Enormous efforts have been devoted to this valuable process including its asymmetric variant in the past decades, nevertheless, the aza-MBH reaction remains to be restricted in substrate scope and enantiocontrol.³ In addition, the nucleophilic catalysts typically utilized in aza-MBH reactions may induce product racemization.⁴ Here we would like to report the development of a novel but facile strategy to the enantioselective synthesis of *N*-Boc- β -amino- α -methylene carboxylic esters.⁵

Stabilized phosphorus ylides (P-ylides) have previously been shown to be good nucleophiles. Aside from the traditional Wittig reaction with carbonyl compounds, P-ylides also found uses in alkylation reactions.⁶ Nevertheless, their potential application as versatile synthons in asymmetric synthesis has not been addressed yet.⁷ We envisaged that P-ylides could act as competent nucleophilic species to attack appropriately activated imines, and the functionalized P-ylides with a chiral center would be generated after proton transfer. Following the reaction with formaldehyde should provide the desired β -amino- α -methylene carbonyl products (Scheme 1).

In light of the above consideration, the reaction of P-ylide 2a and N-Boc imine 3a was initially investigated in toluene at room temperature without any catalyst. Gratifyingly, the expected P-ylide 4a (R = Et) could be isolated as a relatively stable compound, and aza-MBH product 5a (R = Et) was cleanly obtained after reaction with formalin (Table 1, entry 1).⁸ Then we were inspired to explore the possible asymmetric Mannich-type reaction of 2a and 3a catalyzed by chiral Brønsted acids at 0 °C (see Figure 1).9 While low ee values were obtained with catalysis by various phosphoric acids¹⁰ after conversion to **5a** (entries 2-4),¹¹ thiourea **1d** also gave poor chiral induction (entry 5).^{12,13} Moderate ees were received when new thioureas 1e-g were applied (entries 6-8). To our delight, much better enantioselectivity was attained by employing the easily available bisthiourea 1h (entry 9).¹⁴ As a result, the concerted hydrogen bonding interaction of 2a and 3a with two thiourea functional groups might be responsible for the higher enantiocontrol in the Mannich-type reaction step.¹⁵ Ylides 2b and 2c, differing at the ester functionality, led to lower enantioselectivity (entries 10 and 11). Later the reaction was conducted at lower temperature to improve the enantioselectivity, and in general more reliable results could be obtained when 4a was previously isolated before quenching with HCHO.16 Superior results were observed in m-xylene (entry 12 vs 13). Bisthiourea 1i and 1j provided identical enantioselectivities compared to 1h (entries 14 and 15). In addition, the stabilized P-ylides containing a ketone carbonyl exhibited much

Scheme 1. A Novel Approach to Chiral β -Amino- α -methylene Carbonyl Compounds



Table 1. Screening Studies of Organocatalytic Mannich-type Reaction and Subsequent Synthesis of N-Boc- β -amino- α -methylene Carboxylic Esters^a

NHBoc

p				
Q	Bo	C		
\downarrow	+ _=Ń	1 (10 mol %)	Ö	NHBoc HCHO
I OR	Ph		L L	

∫	Ph 3a 2c R = <i>i</i> -Pr	solvent, 0 °C 4Å MS, 24 h	RO PPh ₃	rt, 24 h RO	Ph 5
entry	catalyst	2	solvent	yield ^b (%)	ee ^c (%)
$ \begin{array}{c} 1^{d} \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ 10 \\ 11 \\ 12^{f} \\ 13^{f} \\ 14^{f} \end{array} $	- 1a 1b 1c 1d 1e 1f 1g 1h 1h 1h 1h	2a 2a 2a 2a 2a 2a 2a 2a 2b 2c 2a 2a 2a 2a	toluene toluene toluene toluene toluene toluene toluene toluene toluene toluene m-xylene m-xylene	$\begin{array}{r} \mathbf{5a-53} \\ \mathbf{5a-62} \\ \mathbf{5a-75} \\ \mathbf{5a-60} \\ \mathbf{5a-84} \\ \mathbf{5a-80} \\ \mathbf{5a-69} \\ \mathbf{5a-62} \\ \mathbf{5a-62} \\ \mathbf{5a-62} \\ \mathbf{5a-62} \\ \mathbf{5b-77} \\ \mathbf{5c-51} \\ \mathbf{5a-89} \\ \mathbf{5a-87} \\ \mathbf{5a-87} \\ \mathbf{5a-87} \\ \mathbf{5a-71} \end{array}$	31 45 12 15 63 73 57 80 77 ^e 52 83 89 89
15 ^f	1j	2a	<i>m</i> -xylene	5a -87	89

^{*a*} Unless noted otherwise, reactions were performed with 0.1 mmol of **2**, 0.15 mmol of **3a**, 10 mol % of **1**, 40 mg 4 Å MS in 0.5 mL solvent at 0 °C for 24 h. Then excess HCHO was added and stirred at room temp for another 24 h. ^{*b*} Isolated yield for two steps. ^{*c*} Determined by chiral HPLC analysis. ^{*d*} At room temp for 48 h. ^{*e*} The absolute configuration was determined by converting **5b** to a known compound. ^{3c} f At –20 °C for 60 h, and then **4a** was isolated before treating with formalin in THF for 24 h.

lower reactivity with *N*-Boc imines under the current catalytic system. Attempts to synthesize β -substituted derivatives^{3g,5} with aldehydes other than HCHO failed owing to the steric hindrance of the chiral P-ylide **4a** and its instability to refluxing THF.^{6c} Such problems remain to be further explored.¹⁷

Consequently, the tandem Mannich-type/Wittig reaction for the synthesis of diverse chiral *N*-Boc- β -amino- α -methylene carboxylic esters was investigated with **2a** and a variety of *N*-Boc aldimines catalyzed by 10 mol % of bisthiourea **1h**. As demonstrated in Table 2, the reaction sequence showed substantial generality resulting in broad substrate scope. Excellent enantioselectivities with good

West China School of Pharmacy.

[‡] West China Hospital.



Figure 1. The structures of chiral Brønsted acid catalysts.



Ŭ	Boc 1) 1	h (10 mol %), մ MS _ 20 %	<i>m</i> -xylene					
$\iint OEt \qquad H \qquad 4A MS, -20^{\circ}C, 80 \text{ II} \qquad EtO \qquad R$								
PPh ₃	2)10	ormalin, THF, r	τ, 24 n	II				
2a	3			5				
entry	R	product	yield ^b (%)	ee ^c (%)				
1	Ph	5a	87	89				
2	p-F-Ph	5d	67	87				
3	p-Cl-Ph	5e	65	90				
4	m-Cl-Ph	5f	53	83				
5	o-Cl-Ph	5g	41	89				
6	p-Br-Ph	5h	80	94				
7	p-Me-Ph	5i	78	95				
8	m-Me-Ph	5j	84	93				
9	p-MeO-Ph	5k	84	91				
10	2-thienyl	51	84	68				
11^{d}	n-propyl	5m	70	57				
12	cyclohexyl	5n	52	91				
13	<i>i</i> -propyl	50	35 (63) ^e	96 (91) ^e				
14 ^f	Ph	5р	74	92				

^{*a*} At 0.1 mmol scale in 0.5 mL *m*-xylene, 2a/3/1h = 1:1.5:0.1. ^{*b*} Isolated yield for two steps. ^{*c*} Determined by HPLC analysis. The absolute configuration of the products was assigned by analogy to 5c. ^{*d*} At -40 °C for 84 h. ^{*e*} Data in parentheses is related to in situ formed imine at 4 °C for 60 h, see Supporting Information. ^{*f*} *N*-Cbz benzaldimine was used.

isolated yields were obtained for *N*-Boc aryl imines bearing various electron-withdrawing or -donating substitutions (entries 1–9), while modest ee was observed for imine with a 2-thienyl group (entry 10). Importantly α -enolizable alkyl imines could be successfully applied, and remarkable ee values were attained for branched substrates (entries 11–13). Moreover, the reaction with in situ formed alkyl imine in the presence of aqueous Cs₂CO₃ solution was promising,¹⁸ and better yield with high ee was gained (entry 13, data in parentheses). It should be stated that alkyl imines have not been effectively utilized in the normal asymmetric aza-MBH reaction.² In addition, good results could be achieved for *N*-Cbz benzaldimine (entry 14). It was noteworthy that catalyst **1h** could be recovered by FC and reused in the Mannich-type reaction without any effects on its efficacy (for **5a**, second use, 81% yield, 89% ee; third use, 85% yield, 90% ee).

In conclusion, we have presented the first asymmetric Mannichtype reaction of stabilized phosphorus ylides and Boc-protected aldimines by employing readily available and recyclable bisthiourea organocatalysts. Subsequent reaction with formaldehyde provides a facile access to chiral *N*-Boc- β -amino- α -methylene carboxylic esters in good to excellent enantioselectivities. This methodology presented herein may potentially open avenues for the application of phosphorus ylides in asymmetric synthesis. Currently mechanism exploration and reaction expansion of the related ylides are under way in our laboratory and will be reported in due course.

Acknowledgment. We are grateful for the financial support from NSFC (20502018), Education of Ministry (NCET-05-0781), and Sichuan Province Government (07ZQ026-027).

Supporting Information Available: Experimental procedures, structural proofs. This material is available free of charge via the Internet at http://pubs.acs.org.

References

- (a) Basavaiah, D.; Rao, A. J.; Satyanarayana, T. Chem. Rev. 2003, 103, 811. (b) Sorbetti, J. M.; Clary, K. N.; Rankic, D. A.; Wulff, J. E.; Parvez, M.; Back, T. G. J. Org. Chem. 2007, 72, 3326. (c) Nemoto, T.; Fukuyama, T.; Yamamoto, E.; Tamura, S.; Fukuda, T.; Matsumoto, T.; Akimoto, Y.; Hamada, Y. Org. Lett. 2007, 9, 927.
- (2) For recent reviews on aza-MBH reaction, see: (a) Shi, Y.-L.; Shi, M. Eur. J. Org. Chem. 2007, 2905. (b) Masson, G.; Housseman, C.; Zhu, J. Angew. Chem., Int. Ed. 2007, 46, 4614. (c) Basavaiah, D.; Rao, K. V.; Reddy, R. J. Chem. Soc. Rev. 2007, 36, 1581.
- (3) (a) Shi, M.; Xu, Y.-M. Angew. Chem., Int. Ed. 2002, 41, 4507. (b) Shi, M.; Chen, L.-H.; Li, C.-Q. J. Am. Chem. Soc. 2005, 127, 3790. (c) Shi, M.; Xu, Y.-M.; Shi, Y.-L. Chem.-Eur. J. 2005, 117, 1794. (d) Kawahara, S.; Nakano, A.; Esumi, T.; Iwabuchi, Y.; Hatakeyama, S. Org. Lett. 2003, 5, 3103. (e) Matsui, K.; Takizawa, S.; Sasai, H. J. Am. Chem. Soc. 2005, 127, 3680. (f) Gausepohl, R.; Buskens, P.; Kleinen, J.; Bruckmann, A.; Lehmann, C. W.; Klankermayer, J.; Leitner, W. Angew. Chem., Int. Ed. 2006, 45, 3689. (g) Utsumi, N.; Zhang, H.; Tanaka, F.; Barbas, C. F., III. Angew. Chem., Int. Ed. 2007, 46, 1878. (h) Ito, K.; Nishida, K.; Gotanda, T. Tetrahedron Lett. 2007, 48, 6147.
- (4) (a) Buskens, P.; Klankemayer, J.; Leitner, W. J. Am. Chem. Soc. 2005, 127, 16762.
- (5) For nontraditional approaches to access MBH or aza-MBH reaction products, see: (a) Reynolds, T. E.; Bharadwaj, A. R.; Scheidt, K. A. J. Am. Chem. Soc. 2006, 128, 15382. (b) Reynolds, T. E.; Scheidt, K. A. Angew. Chem., Int. Ed. 2007, 46, 7806. (c) Trost, B. M.; Chung, C. K. J. Am. Chem. Soc. 2006, 128, 10358. (d) Yamaguchi, A.; Aoyama, N.; Matsunaga, S.; Shibasaki, M. Org. Lett. 2007, 9, 3387. (e) Garcia Ruano, J. L.; Fernandez, I.; del Prado Catalina, M.; Hermoso, J. A.; Sanz-Aparicio, J.; Martinez-Ripoll, M. J. Org. Chem. 1998, 63, 7157.
- (6) (a) Asunskis, J.; Shechter, H. J. Org. Chem. 1968, 33, 1164. (b) von Strandtmann, M.; Cohen, M. P.; Puchalski, C.; Shavel, J., Jr. J. Org. Chem. 1968, 33, 4306. (c) Connor, D. T.; von Strandtmann, M. J. Org. Chem. 1973, 38, 1047.
- (7) Stable sulfur ylides have been applied in the organocatalytic asymmetric cyclopropanations, see: Kunz, R. K.; MacMillan, D. W. C. J. Am. Chem. Soc. 2005, 127, 3240.
- (8) High ee values but modest yields have been obtained in the aza-MBH reaction of acrylates and N-Nos imines, see: Raheem, I. T.; Jacobsen, E. N. Adv. Synth. Catal. 2005, 347, 1701.
- (9) For recent reviews on chiral Brønsted acid catalysis, see: (a) Akiyama, T. Chem. Rev. 2007, 107, 5744. (b) Doyle, A. G.; Jacobsen, E. N. Chem. Rev. 2007, 107, 5713.
- (10) For selected examples on phosphoric acid catalysis, see: (a) Akiyama, T.; Itoh, J.; Yokota, K.; Fuchibe, K. Angew. Chem., Int. Ed. 2004, 43, 1566. (b) Uraguchi, D.; Terada, M. J. Am. Chem. Soc. 2004, 126, 5356. (c) Uraguchi, D.; Sorimachi, K.; Terada, M. J. Am. Chem. Soc. 2005, 127, 9360. (d) Hoffmann, S.; Seayad, A. M.; List, B. Angew. Chem., Int. Ed. 2005, 44, 9360. (e) Storer, R. I.; Carrera, D. E.; Ni, Y.; MacMillan, D. W. C. J. Am. Chem. Soc. 2006, 128, 84.
- (11) Current attempts to separate ylide 4a on chiral HPLC columns were unsuccessful.
- (12) For selected examples on thiourea catalysis, see: (a) Sigman, M. S.; Jacobsen, E. N. J. Am. Chem. Soc. 1998, 120, 4901. (b) Okino, T.; Hoashi, Y.; Takemoto, Y. J. Am. Chem. Soc. 2003, 125, 12672. (c) Berkessel, A.; Cleemann, F.; Mukherjee, S. Angew. Chem., Int. Ed. 2005, 44, 7466. (d) Yamaoka, Y.; Miyabe, H.; Takemoto, Y. J. Am. Chem. Soc. 2007, 129, 6686. (e) Martin, N. J. A.; Ozores, L.; List, B. J. Am. Chem. Soc. 2007, 129, 8976. (f) Raheem, I. T.; Thiara, P. S.; Peterson, E. A.; Jacobsen, E. N. J. Am. Chem. Soc. 2007, 129, 13404.
- (13) For recent studies on thiourea catalysis from this group, see: (a) Liu, T.-Y.; Cui, H.-L.; Chai, Q.; Long, J.; Li, B.-J.; Wu, Y.; Ding, L.-S.; Chen, Y.-C. *Chem. Commun.* **2007**, 2228. (b) Liu, T.-Y.; Cui, H.-L.; Long, J.; Li, B.-J.; Wu, Y.; Ding, L.-S.; Chen, Y.-C. *J. Am. Chem. Soc.* **2007**, *129*, 1878.
- (14) (a) Sohtome, Y.; Tanatani, A.; Hashimoto, Y.; Nagasawa, K. *Tetrahedron Lett.* 2004, 45, 5589. (b) Berkessel, A.; Roland, K.; Neudorfl, J. M. Org. Lett. 2006, 8, 4195.
- (15) For selected examples in which additional hydrogen bonding interaction plays a crucial role in the stereocontrol, see: (a) Yoon, T. P.; Jacobsen, E. N. Angew. Chem., Int. Ed. 2005, 44, 466. (b) Herrera, R. P.; Sgarzani, V.; Bernardi, L.; Ricci, A. Angew. Chem., Int. Ed. 2005, 44, 6576. (c) Sibi, M. P.; Itoh, K. J. Am. Chem. Soc. 2007, 129, 8064. (d) Zeng, W.; Chen, G.-Y.; Zhou, Y.-G.; Li, Y.-X. J. Am. Chem. Soc. 2007, 129, 750. (e) Chem, W.; Du, W.; Duan, Y.-Z.; Wu, Y.; Yang, S.-Y.; Chen, Y.-C. Angew. Chem., Int. Ed. 2007, 46, 7667.
- (16) The possible background reaction of unconsumed P-ylide and imine in the quenching step would reduce the final ee.
- (17) Preliminary results indicate that less-hindered tributylphosphoranes are more reactive toward *N*-Boc imines yet lead to lower enantioselectivities.
- (18) Song, J.; Shih, H.-W.; Deng, L. Org. Lett. 2007, 9, 603.

JA7114844