

Communication

Asymmetric Radical Cyclopropanation of Alkenes with In Situ-Generated Donor-Substituted Diazo Reagents via Co(II)-Based Metalloradical Catalysis

Yong Wang, Xin Wen, Xin Cui, Lukasz Wojtas, and Peter Zhang

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.6b11336 • Publication Date (Web): 04 Jan 2017

Downloaded from <http://pubs.acs.org> on January 4, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

Asymmetric Radical Cyclopropanation of Alkenes with In Situ-Generated Donor-Substituted Diazo Reagents via Co(II)-Based Metalloradical Catalysis

Yong Wang,[§] Xin Wen,[§] Xin Cui,[†] Lukasz Wojtas[†] and X. Peter Zhang^{*§†}

[§]Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467, United States

[†]Department of Chemistry, University of South Florida, Tampa, Florida 33620, United States


Supporting Information Placeholder

ABSTRACT: Donor-substituted diazo reagents, generated in situ from sulfonyl hydrazones in the presence of base, can serve as suitable radical precursors for Co(II)-based metalloradical catalysis (MRC). The cobalt(II) complex of *D*₂-symmetric chiral porphyrin [Co(3,5-Di^tBu-Xu(2'-Naph)Pyrin)] is an efficient metalloradical catalyst that is capable of activating different *N*-arylsulfonyl hydrazones for asymmetric radical cyclopropanation of a broad range of alkenes, affording the corresponding cyclopropanes in high yields with effective control of both diastereoselectivity and enantioselectivity. This Co(II)-based metalloradical system represents the first catalytic protocol that can effectively utilize donor-type diazo reagents for asymmetric olefin cyclopropanation.

There has been lasting interest in devising strategies to control stereoselectivity, especially enantioselectivity, of radical reactions for applications in organic synthesis.¹ Among recent advances,^{1,2} metalloradical catalysis (MRC), which aims at the development of metalloradical-based systems for both catalytic initiation and selective control of radical processes, has led to the discovery of new catalytic pathways for stereoselective radical reactions.^{3,4} As stable metalloradicals, cobalt(II) complexes of *D*₂-symmetric chiral amidoporphyrins [Co(*D*₂-Por^{*})] have emerged as effective catalysts for asymmetric radical transformations through catalytic generation of metal-stabilized organic radicals, such as the fundamentally new α -metalloalkyl radicals and α -metalloaminy radicals, as the key intermediates.⁵ To date, Co(II)-based metalloradical catalysis (Co(II)-MRC) has exhibited capability of activating both acceptor- and acceptor/acceptor-substituted diazo reagents to generate corresponding α -Co(III)-alkyl radicals (also known as Co(III)-carbene radicals) as key intermediates for various C-centered radical processes, including radical cyclopropanation of alkenes.^{6,7} Studies suggest that potential hydrogen-bonding interactions between carbonyl groups of these types of diazo reagents and the amide units of the porphyrin ligands in the resulting α -Co(III)-alkyl radicals play an important role in enhancing reactivity and controlling stereoselectivity. It was unclear if Co(II)-MRC could also be applicable to other types of diazo reagents that lack a carbonyl group, such as donor-substituted diazo reagents. Specifically, we wondered whether [Co(*D*₂-Por^{*})] metalloradical catalysts were able to activate aryldiazomethanes to generate the corresponding α -Co(III)-benzyl radical intermediate **I** (Scheme 1). In the absence of carbonyl functionalities, what element of interaction could be explored for the effective control of enantioselectivity in the subsequent radical addition of the α -Co(III)-benzyl radical **I** to the olefin substrate? Furthermore, would the final step, 3-*exo-tet* radical cyclization of the resulting γ -Co(III)-alkyl radicals **II**, be diastereoselec-

tive? Additional questions arose from the intrinsic instability of donor-substituted diazomethanes, which are typically generated in situ from stable sulfonyl hydrazone precursors.⁸ Would Co(II)-MRC system be compatible with the basic conditions required for the in-situ generation protocol? If these questions could be answered positively, it would further expand the application of Co(II)-based radical cyclopropanation for the catalytic synthesis of enantioenriched 1,2-bisaryl and related cyclopropane compound **3**.


Scheme 1. Proposed Pathway for Asymmetric Radical Cyclopropanation with Donor-Substituted Diazo Reagents




Asymmetric olefin cyclopropanation with diazo reagents represents one of the most general methods for the construction of optically active three-membered carbocycles.⁹ While tremendous progress has been made with the use of several types of diazo reagents, asymmetric cyclopropanation with donor-substituted diazo reagents has been much less developed.¹⁰ As a novel alternative involving the further transfer of the initially formed electrophilic Rh₂-carbenes into nucleophilic chiral sulfur ylides, Aggarwal and coworkers developed an indirect approach for asymmetric cyclopropanation with sulfonyl hydrazones as diazo precursors using the combination of dirhodium carboxylates and chiral sulfides as the catalyst that works specifically for electron-deficient olefins.^{8c,10c} Nevertheless, it would be desirable to develop new catalytic systems that are capable of direct asymmetric cyclopropanation of diverse alkenes with donor-substituted diazo reagents.¹¹ In view of the radical pathway of MRC (Scheme 1), we hypothesized that the radical cyclopropanation approach could potentially address the aforementioned issues with donor-substituted diazo reagents because a neutral radical process is generally less sensitive to electronic effects. Herein, we wish to report the development of the first asymmetric catalytic system that is highly effective for direct cyclopropanation of alkenes with α -aryl diazomethanes. Under mild

conditions, the Co(II)-based metalloradical system is suitable for various alkenes, allowing for the efficient construction of cyclopropane rings with excellent control of both diastereoselectivity and enantioselectivity. The Co(II)-catalyzed cyclopropanation is further highlighted by its functional group tolerance, a feature that is closely related to the underlying radical mechanism.

Table 1. Asymmetric Cyclopropanation of Styrene with Tosylhydrazones by Metalloradical Catalysts [Co(*D*₂-Por*)]^a



entry	R	catalyst	product	yield (%) ^b	dr ^c	ee (%) ^d
1	H (1a)	[Co(PPP)]	3aa	46	88:12	-
2	H (1a)	[Co(P1)]	3aa	64	89:11	-42
3	H (1a)	[Co(P2)]	3aa	65	78:22	<5
4	H (1a)	[Co(P3)]	3aa	67	95:5	26
5	2-MeO (1b)	[Co(P3)]	3ba	78	95:5	99
6	2-Et (1c)	[Co(P3)]	3ca	87	85:15	33
7	4-MeO (1d)	[Co(P3)]	3da	23	93:7	23
8	3-MeO (1e)	[Co(P3)]	3ea	69	95:5	23




^a Carried out with **1x** (0.1 mmol) and **2a** (1.5 equiv.) in the presence of Cs₂CO₃ (2 equiv.) in methanol (0.6 mL). ^b Isolated yields. ^c Determined by ¹H NMR. ^d Determined by chiral HPLC for the major *trans* diastereomer.

Our study began with cyclopropanation of styrene (**2a**) with benzaldehyde tosylhydrazone (**1a**), which is known to generate α -phenyldiazomethane under basic conditions (Table 1). Even simple metalloradical catalyst [Co(PPP)] (PPP = 5,10,15,20-tetraphenylporphyrin) could catalyze the reaction in the presence of Cs₂CO₃ in methanol, forming the desired 1,2-diphenyl cyclopropane **3aa** in 46% yield (entry 1). This result signifies the compatibility of Co(II)-based metalloradical system with the use of both bases and polar solvents. The stereoselectivity of the reaction was then evaluated by using Co(II) complexes of *D*₂-symmetric chiral amidoporphyrins [Co(*D*₂-Por*)]. When the first-generation catalyst [Co(**P1**)] (**P1** = 3,5-Di^tBu-ChenPhyrin) was used,^{6f} **3aa** could form in higher yield with significant level of both enantioselectivity and diastereoselectivity (entry 2). To further improve the catalytic reaction, we then turned our attention to the second-generation Co(II)-based metalloradical catalysts.^{6b} While the phenyl-substituted [Co(**P2**)] (**P2** = 3,5-Di^tBu-QingPhyrin) was less selective, the naphthyl-substituted [Co(**P3**)] (**P3** = 3,5-Di^tBu-Xu(2'-Naph)Phyrin)) could effectively catalyze the formation of **3aa** with excellent diastereoselectivity but lower enantioselectivity (entry 4). It should be noted that comparable reactivity was observed with the use of preformed α -phenyldiazomethane (see Table S1 in SI). The ineffective asymmetric induction is likely attributed to the low-barrier rotation of the α -benzyl radical unit around the Co(III)-C σ bond in intermediate **I** (Scheme 1) that diminishes its approaching preference to the prochiral styrene substrate between the *re* and *si* faces. To restrict the rotation, we decided to use benzaldehyde tosylhydrazone derivative that contains an *ortho*-methoxy group

(**1b**). Considering MeO group is known as good hydrogen bond acceptors,¹² we postulated that the potential formation of N-H...O hydrogen bond between the amido group of the chiral ligand and the MeO group of tosylhydrazone **1b** (Figure S1 in SI) might rigidify the resulting intermediate **I**, leading to improved stereoselectivity. Indeed, the corresponding cyclopropane **3ba** was obtained in good yield (78%) with high diastereoselectivity (95:5 dr) and excellent enantioselectivity (99% ee) (entry 5; see Table S2 in SI for detailed solvent effect). Conversely, asymmetric induction of the cyclopropanation process was significantly reduced when the MeO group in **1b** was replaced by a sterically comparable Et group despite the high reactivity (entry 6). Moreover, low enantioselectivity was observed when the MeO group in **1b** was moved from *ortho*- to either *meta*- or *para*- positions of the phenyl ring although with similarly high diastereoselectivity (entries 7 and 8). These results seem support the hypothesized hydrogen-bonding interaction.

Table 2. Asymmetric Cyclopropanation of Different Olefins with Tosylhydrazone 1b Catalyzed by [Co(P3**)]^a**



entry	olefin	product	yield (%)	dr	ee (%)
1 ^b	2a	3ba	83%	95:5, ee: 99%	
2	2b	3bb	88%	95:5, ee: 94%	
3	2c	3bc	81%	96:4, ee: 93%	
4 ^c	2d	3bd	85%	94:6, ee: 94%	
5	2e	3be	85%	94:6, ee: 94%	
6	2f	3bf	89%	95:5, ee: 96%	
7 ^d	2g	3bg	61%	90:10, ee: 96%	
8	2h	3bh	90%	83:17, ee: 93%	
9	2i	3bi	42%	83:17, ee: 92%	
10	2j	3bj	71%	75:25, ee: 95%	
11	2k	3bk	69%	92:8, ee: 97%	
12	2l	3bl	68%	83:17, ee: 99%	
13	2m	3bm	77%	81:19, ee: 81%	
14	2n	3bn	50%	78:22, ee: 89%	
15	2o	3bo	49%	83:17, ee: 91%	
16	2p	3bp	41%	71:29, ee: 77%	
17 ^e	2q	3bq	42%	75:25, ee: 82%	

^a Carried out with **1b** (0.1 mmol) and **2x** (1.5 equiv.) in the presence of Cs₂CO₃ (2 equiv.) in methanol (0.6 mL); isolated yields; dr was determined by ¹H NMR; ee of the major *trans* diastereomer was determined by chiral HPLC. ^b Performed on 1.0 mmol scale. ^c Absolute configuration was determined by X-ray as (1*S*, 2*S*). ^d 5 mol % [Co(**P3**)]. ^e 5.0 equiv. **2q**.

Under the optimized conditions, the scope of [Co(**P3**)]/**1b**-based catalytic system was then evaluated by employing different alkenes (Table 2). Like styrene, its derivatives bearing substituents with different electronic properties, including electron-donating MeO and electron-withdrawing CF₃ groups, could be cyclopropanated in high yields with excellent diastereoselectivity and enantioselectivity (entries 1–3). Halogenated styrenes such as those with a Br atom at various positions were also shown to be suitable substrates for the catalytic process (entries 4–6). The relative and absolute configurations of cyclopropane **3bd** (entry 4) were established as *trans* and (1*S*, 2*S*), respectively (see Figure S2 in SI). As demonstrated for 2-bromostyrene (**2f**) together with 2,4,6-trimethylstyrene (**2g**), sterically hindered styrene derivatives could also undergo productive cyclopropanation (entries 6 and 7). Furthermore, this system was also applicable for 1,1-disubstituted olefins as demonstrated with α -substituted styrene derivatives **2h**, **2i** and **2j**, affording the desired cyclopropanes with enantioselective control of the newly-formed quaternary stereogenic centers, albeit with moderate diastereoselectivity (entries 8–10). Distinctly, the Co(II)-based radical process could tolerate functional groups as exemplified by the highly enantioselective cyclopropanation of 3-aminostyrene (entry 11). In addition to the expanded aromatic olefins such as 2-vinylnaphthalene (entry 12), conjugated dienes, including both aromatic diene **2m** and aliphatic diene **2n**, could also be employed as

substrates but with relatively lower stereoselectivities (entries 13 and 14). It was notable that both electron-rich and -deficient non-aromatic olefins, such as vinyl ether **2o**, acrylamide **2p** and vinyl ketone **2q** could be enantioselectively cyclopropanated as well (entries 15–17) with moderate control of stereoselectivities. The observed substrate scope and reactivity profile of [Co(**P3**)]-catalyzed cyclopropanation is in agreement with the underlying radical mechanism (Scheme 1). It was also noteworthy to mention that the one-pot protocol of the Co(II)-catalyzed cyclopropanation could be scaled up ten-fold as demonstrated with the high-yielding synthesis of cyclopropane **3ba** on 1.0 mmol scale without affecting the excellent stereoselectivity (entry 1).

Table 3. Asymmetric Cyclopropanation of Styrene with Various Sulfonyl Hydrazones Catalyzed by [Co(P3**)]^a**


entry	Ar	R ^b	product	temp. (°C)	yield (%) ^c	dr ^d	ee (%) ^e
1		Ts (1b)	3ba	40	78	95:5	99
2		Ts (1f)	3fa	40	91	95:5	94
3		Ts (1g)	3ga	RT	90	94:6	86
4		Ts (1g)	3ga	0	<10	-	-
5		TPS (1g')	3g'a	0	83	>99:1	93
6		Ts (1h)	3ha	RT	58	96:4	71
7		TPS (1h')	3h'a	0	75	>99:1	76
8		Ts (1i)	3ia	RT	85	96:4	88
9		TPS (1i')	3i'a	0	85	>99:1	93
10		Ts (1j)	3ja	RT	82	95:5	68
11		TPS (1j')	3j'a	0	81	>99:1	89

^a Carried out with **1x** (0.1 mmol) and **2a** (1.5 equiv.) in the presence of Cs₂CO₃ (2 equiv.) in methanol (0.6 mL). ^b Ts = 4-toluenesulfonyl; TPS = (2,4,6-triisopropyl)phenyl sulfonyl. ^c Isolated yields. ^d Determined by ¹H NMR. ^e Determined by chiral HPLC for the major *trans* diastereomer.

Guided by the postulated hydrogen-bonding interaction in the key intermediates **I**, we then sought to identify other benzaldehyde sulfonylhydrazone derivatives as effective radical precursors for asymmetric cyclopropanation (Table 3). Like arylhydrazone **1b** (entry 1), 2,5-dimethoxybenzaldehyde-derived tosylhydrazone (**1f**) could be also effectively activated by [Co(**P3**)] for cyclopropanation of styrene, affording cyclopropane **3fa** in 91% yield with 90% de and 94% ee (entry 2). Considering that fluorine atoms in organic fluorides have been demonstrated as a potential hydrogen bond acceptor,¹³ we explored the use of fluorine-containing arylhydrazones as diazo precursors. Remarkably, fluoroarene-based tosylhydrazones such as **1g–1j** were found to be even more active than **1b**, enabling the catalytic process even at room temperature (entries 3, 6, 8 and 10). The desired fluorine-containing bisaryl cyclopropanes **3ga–3ja** were obtained in up to 90% yield with 88–92% de and 68–88% ee. Attempts to further improve enantioselectivity with **1g** at lower temperature were unsuccessful due to ineffective generation of 2,6-difluorophenyl diazomethane at 0 °C (entry 4). Since the decomposition of the sterically bulky (2,4,6-triisopropylphenyl)sulfonyl hydrazone salt was shown to be considerably faster than tosylhydrazone salt,¹⁴ the low temperature asymmetric cyclopropanation was reexamined with the use of 2,6-difluorobenzaldehyde-derived (2,4,6-triisopropylphenyl)sulfonyl hydrazone **1g'**. As expected, the reaction could proceed effectively at 0 °C, affording **3ga** in 83% yield with considerably enhanced stereoselectivities (entry

5). Likewise, the reactions with related (2,4,6-triisopropylphenyl)sulfonyl hydrazones **1h'–1j'** could also be productively conducted at 0 °C, affording the cyclopropanes **3ha–3ja** with significantly improved diastereoselectivity and enantioselectivity (entries 7, 9 and 11). Under the same conditions, while *o*-Cl-benzaldehyde-derived sulfonylhydrazone gave a similar result to *o*-F-benzaldehyde-derived sulfonylhydrazone, *o*-Br- and *o*-I-benzaldehyde-derived sulfonylhydrazones proved to be less effective radical precursors for the catalytic process (see Scheme S1 in SI).

Scheme 2. Cyclopropanation of (*E*)- and (*Z*)- β -Deuterostyrenes to Probe Radical Reaction Mechanism



To probe the underlying stepwise radical mechanism of the Co(II)-catalyzed cyclopropanation, both (*E*)- and (*Z*)- β -deuterostyrenes were employed as substrates (Scheme 2). While a concerted mechanism is usually stereospecific, a stepwise radical mechanism (Scheme 1) would result in the formation of four possible diastereomers from either (*E*)- or (*Z*)- β -deuterostyrene: two *trans*-isotopomers **A** & **B** and two *cis*-isotopomers **C** & **D** due to the rotation of β -C–C bond in the γ -Co(III)-alkyl radical intermediate **II** before ring closure (Schemes S2 and S3 in SI). Using catalyst [Co(**P3**)], cyclopropanation of (*E*)- β -deuterostyrene with tosylhydrazone **1g** gave *trans*-**3ga** as the dominant product with a 93:7 ratio of isotopomers **A** and **B**. Under the same conditions, the reaction of (*Z*)- β -deuterostyrene resulted in the identical 93:7 ratio of isotopomeric distribution but in favoring **B** over **A**. The observation of *trans*-isotopomer **B** from (*E*)- β -deuterostyrene as well as *trans*-isotopomer **A** from (*Z*)- β -deuterostyrene is evidently a result of the rotation of β -C–C bond in intermediate **II**. When sterically less-hindered [Co(**P4**)] (**P4** = 3,5-Di^tBu-IbuPhyrin) was used as the catalyst,¹⁵ a significantly different isotopomeric ratio of *trans*-isotopomers **A** & **B** (from 93:7 to 80:20) was observed, suggesting easier rotation of the β -C–C bond in a less-crowded ligand environment (Schemes S4 and S5 in SI). Together, these observations convincingly support the proposed stepwise radical mechanism of the Co(II)-based cyclopropanation with *N*-arylsulfonyl hydrazones. Furthermore experiments were performed for direct detection of the α -Co(III)-benzyl radical species **I** (Scheme 1) by high-resolution mass spectrometry (HRMS).¹⁶ The exposure of [Co(**P4**)] to **1g** (10 equiv.) with base in methanol resulted in a mixture that was filtrated and analyzed by HRMS in the absence of any additives such as formic acids that commonly act as electron carriers for ESI ionization. As shown in Scheme S6 in SI, the obtained spectrum clearly revealed a signal corresponding to [Co(**P4**)](2,6-F₂C₆H₃CH)⁺ (*m/z* = 1361.6515), which resulted from the neutral α -Co(III)-benzyl radical **I** by the loss of one electron.

In summary, Co(II)-based metalloradical catalysis (MRC) has, for the first time, been successfully applied to donor-substituted diazo reagents, generated from *N*-arylsulfonyl hydrazones in the presence of base, for olefin cyclopropanation. Chiral metalloradical complex [Co(**P3**)] has been shown to be an effective catalyst for asymmetric radical cyclopropanation with α -aryldiazomethane precursors. The Co(II)-based cyclopropanation is applicable to a broad combination of *N*-arylsulfonyl hydrazones and alkenes, affording

the cyclopropane derivatives in high yields with excellent diastereoselectivity and enantioselectivity. In view of the underdevelopment of donor-substituted diazo reagents in asymmetric cyclopropanation, the successful development of this catalytic radical process will encourage further efforts in applying sulfonyl hydrazones as stable precursors for various asymmetric catalytic processes.

ASSOCIATED CONTENT

Supporting Information

Experimental details and analytical data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

* peter.zhang@bc.edu

ACKNOWLEDGMENT

We are grateful for financial support by NIH (R01-GM102554).

REFERENCES

- (1) For reviews on stereoselectivity of free radical transformations, see: (a) Yang, Y.-H.; Sibi, M. P. In *Encyclopedia of Radicals in Chemistry, Biology and Materials*; Chatgililoglu, C., Studer, A., Eds.; Wiley: Weinheim, 2012, p 655; (b) Curran, D. P.; Porter, N. A.; Giese, B. *Stereochemistry of radical reactions: concepts, guidelines, and synthetic applications*; John Wiley & Sons, 2008; (c) Zimmerman, J.; Sibi, M. P. *Top. Curr. Chem.* **2006**, 263, 107; (d) Bar, G.; Parsons, A. F. *Chem. Soc. Rev.* **2003**, 32, 251; (e) Zard, S. Z. *Radical reactions in organic synthesis*; Oxford University Press: Oxford, 2003.
- (2) For select examples of stereoselective radical processes via photoredox catalysis, see: (a) Kainz, Q. M.; Matier, C. D.; Bartoszewicz, A.; Zultanski, S. L.; Peters, J. C.; Fu, G. C. *Science* **2016**, 351, 681; (b) Brimouille, R.; Lenhart, D.; Maturi, M. M.; Bach, T. *Angew. Chem., Int. Ed.* **2015**, 54, 3872; (c) Huo, H. H.; Shen, X. D.; Wang, C. Y.; Zhang, L. L.; Rose, P.; Chen, L. A.; Harms, K.; Marsch, M.; Hilt, G.; Meggers, E. *Nature* **2014**, 515, 100; (d) Du, J. N.; Skubi, K. L.; Schultz, D. M.; Yoon, T. P. *Science* **2014**, 344, 392.
- (3) For reviews and highlights on Co(II)-MRC: (a) Studer, A.; Curran, D. P. *Angew. Chem., Int. Ed.* **2016**, 55, 58; (b) Pellissier, H.; Clavier, H. *Chem. Rev.* **2014**, 114, 2775; (c) Degennaro, L.; Trinchera, P.; Luisi, R. *Chem. Rev.* **2014**, 114, 7881; (d) Olivos Suarez, A. I.; Lyaskovskyy, V.; Reek, J. N.; van der Vlugt, J. I.; de Bruin, B. *Angew. Chem., Int. Ed.* **2013**, 52, 12510; (e) Doyle, M. P. *Angew. Chem., Int. Ed.* **2009**, 48, 850.
- (4) For Ti(III)-based MRC, see: (a) Gansäuer, A.; Hildebrandt, S.; Vogelsang, E.; Flowers Ii, R. A. *Dalton Trans.* **2016**, 45, 448; (b) RajanBabu, T. V.; Nugent, W. A. *J. Am. Chem. Soc.* **1994**, 116, 986; (c) Gansäuer, A.; Hildebrandt, S.; Michelmann, A.; Dahmen, T.; von Laufenberg, D.; Kube, C.; Fianu, G. D.; Flowers, R. A. *Angew. Chem., Int. Ed.* **2015**, 54, 7003; (d) Gansäuer, A.; Fleckhaus, A.; Lafont, M. A.; Okkel, A.; Kotsis, K.; Anoop, A.; Neese, F. *J. Am. Chem. Soc.* **2009**, 131, 16989; (e) Gansäuer, A.; Fan, C.-A.; Keller, F.; Keil, J. *J. Am. Chem. Soc.* **2007**, 129, 3484; (f) Gansäuer, A.; Rinker, B.; Pierobon, M.; Grimme, S.; Gerenkamp, M.; Mück-Lichtenfeld, C. *Angew. Chem., Int. Ed.* **2003**, 42, 3687.
- (5) For detailed experimental and theoretical studies on the radical mechanism of Co(II)-MRC: (a) Dzik, W. I.; Xu, X.; Zhang, X. P.; Reek, J. N. H.; de Bruin, B. *J. Am. Chem. Soc.* **2010**, 132, 10891; (b) Belof, J. L.; Cioce, C. R.; Xu, X.; Zhang, X. P.; Space, B.; Woodcock, H. L. *Organometallics* **2011**, 30, 2739; (c) Lu, H. J.; Dzik, W. I.; Xu, X.; Wojtas, L.; de Bruin, B.; Zhang, X. P. *J. Am. Chem. Soc.* **2011**, 133, 8518; (d) Xu, X.; Zhu, S. F.; Cui, X.; Wojtas, L.; Zhang, X. P. *Angew. Chem., Int. Ed.* **2013**, 52, 11857; (e) Xu, X.; Lu, H. J.; Ruppel, J. V.; Cui, X.; de Mesa, S. L.; Wojtas, L.; Zhang, X. P. *J. Am. Chem. Soc.* **2011**, 133, 15292; (f) Zhu, S. F.; Xu, X.; Perman, J. A.; Zhang, X. P. *J. Am. Chem. Soc.* **2010**, 132, 12796; (g) Zhu, S. F.; Ruppel, J. V.; Lu, H. J.; Wojtas, L.; Zhang, X. P. *J. Am. Chem. Soc.* **2008**, 130, 5042; (h) Chen, Y.; Ruppel, J. V.; Zhang, X. P. *J. Am. Chem. Soc.* **2007**, 129, 12074; (i) Chen, Y.; Fields, K. B.; Zhang, X. P. *J. Am. Chem. Soc.* **2004**, 126, 14718; (j) Fantauzzi, S.; Gallo, E.; Rose, E.; Raoul, N.; Caselli, A.; Issa, S.; Ragaini, F.; Cenini, S. *Organometallics* **2008**, 27, 6143.
- (7) For other catalytic processes involving α -Co(III)-alkyl radicals as intermediates, see: (a) Das, B. G.; Chirila, A.; Tromp, M.; Reek, J. N. H.; Bruin, B. d. *J. Am. Chem. Soc.* **2016**, 138, 8968; (b) Zhang, J.; Jiang, J.; Xu, D.; Luo, Q.; Wang, H.; Chen, J.; Li, H.; Wang, Y.; Wan, X. *Angew. Chem., Int. Ed.* **2015**, 54, 1231.
- (8) (a) Xiao, Q.; Zhang, Y.; Wang, J. *Acc. Chem. Res.* **2013**, 46, 236; (b) Barluenga, J.; Valdés, C. *Angew. Chem., Int. Ed.* **2011**, 50, 7486; (c) Fulton, J. R.; Aggarwal, V. K.; de Vicente, J. *Eur. J. Org. Chem.* **2005**, 1479.
- (9) (a) Intriieri, D.; Carminati, D. M.; Gallo, E. *Dalton Trans.* **2016**, 45, 15746; (b) Bartoli, G.; Bencivenni, G.; Dalpozzo, R. *Synthesis* **2014**, 46, 979; (c) Intriieri, D.; Caselli, A.; Gallo, E. *Eur. J. Inorg. Chem.* **2011**, 5071; (d) Lebel, H.; Marcoux, J.-F.; Molinaro, C.; Charette, A. B. *Chem. Rev.* **2003**, 103, 977; (e) Davies, H. M. L.; Antoulakis, E. G. *Org. React.* **2001**, 57, 1; (f) Doyle, M. P.; Forbes, D. C. *Chem. Rev.* **1998**, 98, 911.
- (10) For asymmetric cyclopropanation reactions with donor-type diazo reagents, see: (a) Goudreau, S. R.; Charette, A. B. *J. Am. Chem. Soc.* **2009**, 131, 15633; (b) Berkessel, A.; Kaiser, P.; Lex, J. *Chem. Eur. J.* **2003**, 9, 4746; (c) Aggarwal, V. K.; Alonso, E.; Fang, G.; Ferrara, M.; Hynd, G.; Porcelloni, M. *Angew. Chem., Int. Ed.* **2001**, 40, 1433; (d) Aggarwal, V. K.; Smith, H. W.; Jones, R. V. H.; Fieldhouse, R. *Chem. Commun.* **1997**, 1785.
- (11) For non-asymmetric cyclopropanation reactions with donor-type diazo reagents, see: (a) Verdecchia, M.; Tubaro, C.; Biffis, A. *Tetrahedron Lett.* **2011**, 52, 1136; (b) Dai, X.; Warren, T. H. *J. Am. Chem. Soc.* **2004**, 126, 10085; (c) Maas, G.; Seitz, J. *Tetrahedron Lett.* **2001**, 42, 6137; (d) Li, Y.; Huang, J.-S.; Zhou, Z.-Y.; Che, C.-M. *J. Am. Chem. Soc.* **2001**, 123, 4843; (e) Hamaker, C. G.; Djukic, J.-P.; Smith, D. A.; Woo, L. K. *Organometallics* **2001**, 20, 5189.
- (12) For non-asymmetric cyclopropanation using tosylhydrazones as donor-type diazo precursors, see: (a) Cyr, P.; Cote-Raiche, A.; Bronner, S. M. *Org. Lett.* **2016**, 18, 6448; (b) Reddy, A. R.; Hao, F.; Wu, K.; Zhou, C.-Y.; Che, C.-M. *Angew. Chem., Int. Ed.* **2016**, 55, 1810; (c) Barluenga, J.; Quiñones, N.; Tomás-Gamasa, M.; Cabal, M.-P. *Eur. J. Org. Chem.* **2012**, 2312; (d) Zhang, J.-L.; Hong Chan, P. W.; Che, C.-M. *Tetrahedron Lett.* **2003**, 44, 8733; (e) Adams, L. A.; Aggarwal, V. K.; Bonnert, R. V.; Bressel, B.; Cox, R. J.; Shepherd, J.; de Vicente, J.; Walter, M.; Whittingham, W. G.; Winn, C. L. *J. Org. Chem.* **2003**, 68, 9433; (f) Aggarwal, V. K.; de Vicente, J.; Bonnert, R. V. *Org. Lett.* **2001**, 3, 2785.
- (13) For intramolecular C-H alkylation using hydrazones as donor-type diazo precursors, see: (a) Soldi, C.; Lamb, K. N.; Squitieri, R. A.; Gonzalez-Lopez, M.; Di Maso, M. J.; Shaw, J. T. *J. Am. Chem. Soc.* **2014**, 136, 15142; (b) Reddy, A. R.; Zhou, C. Y.; Guo, Z.; Wei, J.; Che, C. M. *Angew. Chem., Int. Ed.* **2014**, 53, 14175.
- (14) (a) Palusiak, M.; Grabowski, S. J. *J. Mol. Struct.* **2002**, 642, 97; (b) Grabowski, S. J. *J. Phys. Org. Chem.* **2004**, 17, 18.
- (15) Champagne, P. A.; Desroches, J.; Paquin, J. F. *Synthesis* **2015**, 47, 306.
- (16) (a) Chamberlin, A. R.; Bond, F. T. *J. Org. Chem.* **1978**, 43, 154; (b) Cusack, N. J.; Reese, C. B.; Risius, A. C.; Roozpeikar, B. *Tetrahedron* **1976**, 32, 2157.
- (17) Ruppel, J. V.; Jones, J. E.; Huff, C. A.; Kamble, R. M.; Chen, Y.; Zhang, X. P. *Org. Lett.* **2008**, 10, 1995.
- (18) Schäfer, M.; Drayß, M.; Springer, A.; Zacharias, P.; Meerholz, K. *Eur. J. Org. Chem.* **2007**, 5162.

