

Note

Copper-promoted Double Oxidative C-H Amination Cascade for the Synthesis of Imidazo[1,5-a]quinolines

Zhong Li, Song-Song Wu, Zai-Gang Luo, Wei-Kang Liu, Cheng-Tao Feng, and Shi-Tang Ma

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.6b00569 • Publication Date (Web): 25 Apr 2016

Downloaded from <http://pubs.acs.org> on April 25, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

Copper-promoted Double Oxidative C-H Amination Cascade for the Synthesis of Imidazo[1,5-a]quinolines

Zhong Li [†], Song-Song Wu [†], Zai-Gang Luo [†], Wei-Kang Liu [†], Cheng-Tao Feng ^{†*} and Shi-Tang


Ma ^{‡*}

[†]School of Chemical Engineering, Anhui University of Science and Technology, Huainan, Anhui, 232001,
P. R. China

E-mail: fengct2010@163.com

[‡]College of Food and Drug, Anhui Science and Technology University, Fengyang, Anhui, 233100,
P. R. China

E-mail: nothingchina@126.com




ABSTRACT: A copper-promoted cascade reaction of 2-methylazaarenes and benzylamines has been developed *via* sequential double oxidative C(sp³)-H aminations. This protocol provides straightforward access to imidazo[1,5-*a*]quinoline derivatives without employing pre-functionalized substrates.

Imidazo[1,5-*a*]quinolines are an important class of fused N-heterocyclic compounds due to their wide applications in medicinal chemistry (e.g., NK1 receptor ligands).¹ A great deal of attention has been paid to synthesize imidazo[1,5-*a*]quinoline derivatives.² The most popular strategy for the synthesis of imidazo[1,5-*a*]quinolines is Vilsmeier-type cyclizations, which involved the use of N-2-pyridylmethanilamides as substrates.³ Another attractive tool is C-H amination based strategy. Zeng⁴ (Scheme 1a) and Xu⁵ (Scheme 1b) have respectively developed interesting copper catalyzed oxidative tandem cyclization reactions of N-heteroaryl aldehydes or ketones with alkylamines to construct

imidazo[1,5-*a*]quinolines. Recently, Wang has reported a novel protocol to synthesize imidazo[1,5-*a*]pyridines and -quinolines *via* sequential dual oxidative amination of C(sp³)-H bonds in the presence of stoichiometric N-iodobutanamide (Scheme 1c).^{6a} Later, the same reaction was performed employing CuI as catalyst by Adimurthy.^{6b} However, electron-withdrawing substituents on the carbon are necessary for these reactions. Moreover, the amination of 2-alkylazaarenes with aliphatic primary amines is difficult because of oxidative olefination reactions.⁷ The development of efficient methods for the synthesis of imidazo[1,5-*a*]quinolines from 2-methylquinolines without an activating group or pre-functionalization still remains a significant challenge.⁸

Scheme 1. C–H amination based strategies for the synthesis of imidazo[1,5-*a*]quinolines.




2-alkylazaarenes, which are inexpensive and broadly available, have emerged as a useful class of building blocks mainly in the form of a nucleophilic coupling partner for syntheses of various important organic compounds.⁹ The additions of 2-alkylazaarenes to a series of unsaturated bonds, including C=N double bond,^{10a} C=C double bond,^{10b} C=O double bond,^{10c} N=N double bond,^{10d} have been developed mostly to form acyclic compounds, while the cascade reactions of 2-alkylazaarenes to synthesize heterocyclic compounds have not been extensively studied. Metal catalyzed intermolecular direct

oxidative aminations of the sp^2 C–H bond have been well established in recent years.¹¹ As 2-methylquinoline can be isomerized to a nonaromatic enammine intermediate in the presence of transition metal,⁹ we hypothesized that metal catalyzed or promoted dual $C(sp^3)$ –H amination of 2-methylazaarenes with alkylamines might be doable to construct heterocyclic compounds. In continuation of our interest in the quinoline-based synthesis of heterocycles,¹² herein, we describe the direct synthesis of imidazo[1,5-a]quinoline derivatives from commercially accessible 2-methylazaarenes and benzylamines (Scheme 1d).

We began to study the dual $C(sp^3)$ –H amination reaction utilizing 2-methylquinoline **1a** and benzylamine **2a** as model substrates to optimize the reaction conditions (Table 1). When the reaction was performed in DMF in the presence of 1 equiv. of CuI and 2 equiv. of di-*t*-butyl peroxide (DTBP) for 12 h at 110°C, the desired 1-phenylimidazo[1,5-a]quinoline **3aa** was obtained in a 35% yield (Table 1, entry 1). The effect of solvents for this cascade reaction was examined. DMSO was found to be the optimal solvent to give **3aa** in a 57% yield (Table 1, entry 2). When other solvents (polar protic solvents or nonpolar solvents) were utilized, inferior results were obtained (Table 1, entries 3-6). A screening of oxidants, including DTBP, O₂, TBHP, I₂, K₂S₂O₈ and DDQ, showed that DTBP was still to be the best choice (Table 1, entries 7-11). Subsequently, we turned our attention to the effect of copper salts. Among the tested copper salts, such as CuCl, CuBr, Cu(OAc)₂·H₂O, CuI and CuI₂, CuI afforded **3aa** in the highest yield (Table 1, entries 12-15). The reaction did not proceed at all in the absence of a copper salt (Table 1, entry 16). When the amount of CuI was increased to two equivalents, a positive result was observed and the yield of **3aa** was improved to 68% (Table 1, entry 17). Recently, Zhang found that the combination of 1 equiv. of CuCl and 1 equiv. of Cu(OAc)₂·H₂O could mediate annulation of alkyl

ketones with α,β -unsaturated carboxylic acids to afford furan products.¹³ Inspired by this results, we carried out our reaction in the presence of 1 equiv. of CuI and 1 equiv. of Cu(OAc)₂·H₂O, the yield of **3aa** was increased to 83% (Table 1, entry18). When 2.0 equiv. of Cu(OAc)₂·H₂O was used, the yield of **3aa** was decreased to 34% (Table 1, entry19). Based on these results, the optimized reaction conditions of 100 mol% CuI, 100 mol% Cu(OAc)₂·H₂O, and 200 mol% DTBP in DMSO at 110°C were used for further investigation.

Table 1. Optimization of reaction conditions.^a



Entry	Cu salt	Oxidant	Solvent	Yield (%) ^b
1	CuI	DTBP	DMF	35
2	CuI	DTBP	DMSO	57
3	CuI	DTBP	DMA	41
4	CuI	DTBP	EtOH	trace
5	CuI	DTBP	H ₂ O	12
6	CuI	DTBP	PhMe	nd
7	CuI	TBHP	DMSO	36
8	CuI	K ₂ S ₂ O ₈	DMSO	trace
9	CuI	I ₂	DMSO	trace
10	CuI	DDQ	DMSO	trace
11	CuI	O ₂	DMSO	38
12	CuBr	DTBP	DMSO	26

















13	CuCl	DTBP	DMSO	40
14	Cu(OAc) ₂ ·H ₂ O	DTBP	DMSO	45
15	CuI ₂	DTBP	DMSO	17
16	-	DTBP	DMSO	nd
17 ^c	CuI	DTBP	DMSO	68
18 ^d	CuI/Cu(OAc) ₂ ·H ₂ O	DTBP	DMSO	83
19 ^e	Cu(OAc) ₂ ·H ₂ O	DTBP	DMSO	34

^aReaction conditions: **1a** (0.2 mmol), **2a** (0.6 mmol), Cu salt (0.2 mmol), DTBP (0.4 mmol), solvent (1.5 ml), 110 °C, 24 h. ^bIsolated yield. ^cTwo equivalents of CuI were used. ^dCuI (0.2 mmol) and Cu(OAc)₂·H₂O (0.2 mmol) were used. ^eTwo equivalents of Cu(OAc)₂·H₂O were used.

With the optimal conditions in hand, the substrate scope of the double oxidative C-H amination cascade was investigated. As showed in Table 2, a series of aliphatic amines **2** were employed to react with 2-methylquinoline **1a** under the optimized reaction conditions. Benzylamine derivatives with both electron-donating (4-Me, 4-OMe) and electron-withdrawing (4-CF₃) groups on the aromatic ring proceeded in this reaction smoothly to afford the desired products **3** in good to excellent yields. Generally, electron-withdrawing substituents on the aromatic ring have negative effect on the yield. Halogen substituted benzylamine derivatives, including fluorobenzylamines, chlorobenzylamines, or bromobenzylamine, were also well tolerated in this cascade reaction (Table 2, **3ah-3am**), which provides the possibility to allow additional coupling reactions. Interestingly, 2-chlorobenzylamines gave the corresponding product in 85% yield (Table 2, **3aj**). Steric effects had a little influence on the reaction since 2-methylbenzylamine gave similar yield than 4-methylbenzylamine (Table 2, **3ab** and **3ad**). Notably, the ring-fused and heterocyclic derived substrates also amenable to current reaction conditions

to deliver the corresponding products (**3an-3ap**) in good yields (41%-89%). Additionally, the present system is applicable to simple aliphatic amine, and the corresponding imidazo[1,5-*a*]quinoline (**3aq**) was obtained in moderate yield.


Table 2. Scope of Benzylamines^a

			
1a	2a-2p	3aa-3ap	
			
3aa 83%	3ab 92%	3ac 84%	3ad 88%
			
3ae 72%	3af 77%	3ag 51%	3ah 52%
			
3ai 68%	3aj 85%	3ak 71%	3al 73%
			
3am 60%	3an 41%	3ao 89%	3ap 58%
			
3aq 61%			

^aReaction conditions: **1a** (0.2 mmol), **2** (0.6 mmol), CuI (0.2 mmol), Cu(OAc)₂·H₂O (0.2 mmol), DTBP (0.4 mmol), DMSO (1.5 ml), 110 °C, 24 h. Isolated yield.

Next, we turned our attention to the scope of various 2-methylquinoline derivatives to prove the general applicability of the reaction and the results were summarized in Table 3. The reaction is sensitive to the electronic effect of the aromatic ring. 2-Methylquinoline derivatives with electron-donating groups such as -Me, -OMe and -OEt successfully gave the desired imidazo[1,5-*a*]quinolines in good yields (Table 3, **3ba-3fa**). Electron-withdrawing substituents such as -NO₂ (**3ka**) and -Ph (**3ga**) also had deleterious effect on this reaction. Halogen-substituted 2-methylquinolines (Table 3, **3ha-3ja**) were tolerated. Moreover, 4-methyl-2-phenylquinazoline also was found to be a suitable substrate and afforded the desired product (**3la**). However, the present system is not applicable for 2-methyl pyridine (Table 3, **3ma**), most of 2-methylquinoline remained unconsumed.


Table 3. Scope of 2-Methylquinnolines^a



^aReaction conditions: **1a** (0.2 mmol), **2a** (0.6 mmol), CuI (0.2 mmol), Cu(OAc)₂·H₂O (0.2 mmol), DTBP (0.4mmol), DMSO (1.5ml), 110 °C, 24h. Isolated yield.


To establish a possible mechanism of the copper-promoted oxidative C-H amination cascade, some control experiments were carried out. First of all, traces of 2-methylquinoline **1a** were converted into quinoline-2-carbaldehyde **4a** in the absence of **2a** under standard conditions, and most of 2-methylquinoline **1a** remained unconsumed (Scheme 2, eqn. 1). When quinoline-2-carbaldehyde **4a** was allowed to react with **2a**, the target product **3aa** was obtained in 57% yield (Scheme 1, eqn. 2). The reaction course through formation of aldehyde **4a** cannot be excluded. In order to confirm whether the reaction process was performed *via* a radical pathway, **1a** and **2a** were subjected to the standard conditions using TEMPO as radical scavenger (Scheme 2, eqn. 3). As expected, only 12% yield of target product **3aa** was obtained, which suggested that the reaction presumably proceeded through a radical formation pathway.

Scheme 2. Control Experiments for Mechanism



Based on results described above and literature reports,⁹ a plausible mechanism was proposed (Scheme 3). Initially, 2-methylquinoline **1a** is involved a hydrogen abstraction to form a carbon-centered radical **A**.

Scheme 3. Plausible Mechanism



ACS Paragon Plus Environment

can tolerate a broad range of functional groups. In particular, substrate pre-functionalization was not required in the reaction, which avoided the need of extra preparation steps for active precursors.

Experiment Section

General Remarks. Unless otherwise indicated, all commercial reagents and solvents were used without additional purification. ^1H NMR and ^{13}C NMR spectra were recorded at 400MHz and 100MHz, respectively, in CDCl_3 , using tetramethylsilane (TMS) as internal reference. HRMS were recorded using ESI-TOF techniques. Products were purified by flash chromatography on 200–300 mesh silica gels using petroleum ether/ethyl acetate as eluent.

General Procedure for the Synthesis of 3. CuI (38.1 mg, 0.2 mmol) and $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (39.9 mg, 0.2 mmol) was dissolved in a reaction tube using 1.5mL of DMSO, and then benzylamine **2a** (65.6 μL , 0.6 mmol), 2-methylquinoline **1a** (27.1 μL , 0.2 mmol) and DTBP (73.3 μL , 0.4 mmol) were added, sequentially. The mixture was stirred at 110 $^\circ\text{C}$ for 24 h, and the reaction process was monitored by TLC. After the reaction was completed, The reaction system was allowed to attain room temperature and extracted with ethyl acetate (3 $\times$ 20 ml), and then the organic layer was washed with brine (2 $\times$ 10 ml) and dried with anhydrous Na_2SO_4 . Subsequently, the solvent was removed under reduced pressure and the remaining crud product was purified by column chromatography over silica gel (PE/EtOAc=3:1) to afford 1-phenylimidazo[1,5-*a*]quinoline (**3aa**) (40.5 mg, 83%) as a yellow solid.

1-phenylimidazo[1,5-*a*]quinoline (**3aa**).^{2d}

Yellow solid (40.5 mg, 83%), melting point 113–115 $^\circ\text{C}$; ^1H -NMR (400MHz, CDCl_3) δ 7.67-7.65 (m, 3H), 7.55-7.52 (m, 5H), 7.34-7.28 (m, 2H), 7.19-7.16 (m, 1H), 7.03-7.01 (m, 1H); ^{13}C -NMR (100MHz, CDCl_3) δ 142.4, 133.8, 132.3, 130.6, 129.6, 129.3, 128.7, 128.6, 127.3, 125.6, 125.1, 122.4, 121.4, 117.4, 117.1.

HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{17}H_{13}N_2$ 245.1073, found 245.1076.

1-(p-tolyl)imidazo[1,5-a]quinoline (3ab).

Yellow solid (47.5 mg, 92%), melting point 121-125°C; 1H -NMR (400 MHz, $CDCl_3$) δ 7.63-7.54 (m, 5H), 7.35-7.28 (m, 4H), 7.22-7.18 (m, 1H), 7.02 (d, $J=9.3$ Hz, 1H), 2.49 (s, 3H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 142.6, 139.2, 132.5, 130.9, 130.5, 129.4, 128.5, 127.2, 125.6, 125.0, 122.3, 121.2, 117.4, 117.2, 21.5. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{18}H_{15}N_2$ 259.1237, found 259.1235.

1-(m-tolyl)imidazo[1,5-a]quinoline (3ac).

Yellow solid (43.3 mg, 84%), melting point 134-137°C; 1H -NMR (400 MHz, $CDCl_3$) δ 7.60-7.54 (m, 3H), 7.50 (s, 1H), 7.44-7.38 (m, 2H), 7.34-7.27 (m, 3H), 7.19-7.15 (m, 1H), 6.99 (d, $J=9.3$ Hz, 1H), 2.43 (s, 3H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 142.6, 138.6, 133.7, 132.4, 130.5, 130.2, 130.0, 128.5, 128.5, 127.2, 126.5, 125.6, 125.0, 122.3, 121.3, 117.4, 117.1, 21.4. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{18}H_{15}N_2$ 259.1234, found 259.1235.

1-(o-tolyl)imidazo[1,5-a]quinoline (3ad).

Yellow oil (45.4 mg, 88%); 1H -NMR (400 MHz, $CDCl_3$) δ 7.62-7.58 (m, 2H), 7.49-7.46 (m, 2H), 7.39-7.34 (m, 3H), 7.31-7.27 (m, 1H), 7.22 (d, $J=8.4$ Hz, 1H), 7.16-7.12 (m, 1H), 7.02 (d, $J=9.3$ Hz, 1H), 2.06 (s, 3H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 141.5, 138.4, 133.9, 132.6, 130.5, 130.4, 129.9, 129.8, 128.5, 127.8, 126.4, 125.3, 125.0, 122.0, 121.2, 117.2, 115.9, 19.57. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{18}H_{15}N_2$ 259.1233, found 259.1235.

1-(4-methoxyphenyl)imidazo[1,5-a]quinoline (3ae).

Yellow solid (39.5 mg, 72%), melting point 117-121°C; 1H -NMR (400 MHz, $CDCl_3$) δ 7.62-7.56 (m, 4H), 7.53 (s, 1H), 7.33-7.28 (m, 2H), 7.21-7.19 (m, 1H), 7.06 (d, $J=8.7$ Hz, 2H), 7.00 (d, $J=9.3$ Hz, 1H), 3.92 (s,

3H); ^{13}C -NMR (100 MHz, CDCl_3) δ 160.3, 142.3, 132.5, 130.9, 130.4, 128.5, 127.2, 126.1, 125.6, 125.0, 122.2, 121.2, 117.3, 117.2, 114.2, 55.3. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{15}\text{N}_2\text{O}$ 275.1182, found 275.1184.

1-(2-methoxyphenyl)imidazo[1,5-*a*]quinoline (3af).

Yellow solid (42.2 mg, 77%), melting point 118-122°C; ^1H -NMR (400 MHz, CDCl_3) δ 7.62-7.59 (m, 3H), 7.57-7.52 (m, 1H), 7.44 (d, $J=8.5\text{Hz}$, 1H), 7.36 (d, $J=9.3\text{Hz}$, 1H), 7.32-7.28 (m, 1H), 7.20-7.14 (m, 2H), 7.03 (d, $J=8.9\text{Hz}$, 2H), 3.59 (s, 3H); ^{13}C -NMR (100 MHz, CDCl_3) δ 158.1, 139.3, 133.0, 131.9, 131.1, 130.3, 128.2, 127.3, 125.2, 124.8, 123.3, 122.1, 121.2, 121.0, 117.2, 116.4, 111.0, 55.3. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{15}\text{N}_2\text{O}$ 275.1182, found 275.1184.

1-(4-(trifluoromethyl)phenyl)imidazo[1,5-*a*]quinoline (3ag).

Yellow solid (31.8 mg, 51%), melting point 152-156°C; ^1H -NMR (400 MHz, CDCl_3) δ 7.84-7.79 (m, 4H), 7.69 (d, $J=7.7\text{Hz}$, 1H), 7.60 (s, 1H), 7.52 (d, $J=8.4\text{Hz}$, 1H), 7.40-7.37 (m, 2H), 7.29-7.25 (m, 1H), 7.11 (d, $J=9.4\text{Hz}$, 1H); ^{13}C -NMR (100 MHz, CDCl_3) δ 140.7, 137.2, 132.0, 131.2, 131.1, 130.9, 129.8, 128.9, 128.0, 127.5, 125.7, 125.7, 125.6, 125.5, 125.3, 123.0, 122.6, 121.9, 117.3, 117.1. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{12}\text{N}_2\text{F}_3$ 313.0951, found 313.0953.

1-(4-chlorophenyl)imidazo[1,5-*a*]quinoline (3ah).

Yellow solid (29.0 mg, 52%), melting point 148-151°C; ^1H -NMR (400 MHz, CDCl_3) δ 7.64-7.61 (m, 3H), 7.56-7.51 (m, 4H), 7.37-7.35 (m, 2H), 7.28-7.23 (m, 1H), 7.06 (d, $J=9.3\text{Hz}$, 1H); ^{13}C -NMR (100 MHz, CDCl_3) δ 141.1, 135.3, 132.2, 132.2, 130.9, 130.8, 129.0, 128.7, 127.4, 125.6, 125.3, 122.7, 121.6, 117.3, 117.1. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2\text{Cl}$ 279.0689, found 279.0687.

1-(3-chlorophenyl)imidazo[1,5-*a*]quinoline (3ai).

Yellow solid (37.9 mg, 68%), melting point 149-152°C; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.03 (s, 1H), 7.63 (d, $J=6.9\text{Hz}$, 1H), 7.55-7.43 (m, 5H), 7.35-7.32 (m, 2H), 7.25-7.21 (m, 1H), 7.04 (d, $J=9.3\text{Hz}$, 1H); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 140.7, 135.4, 134.7, 132.1, 130.8, 129.9, 129.7, 129.4, 128.8, 127.7, 127.5, 125.6, 125.3, 122.7, 121.7, 117.3, 117.0. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2\text{Cl}$ 279.0688, found 279.0689.

1-(2-chlorophenyl)imidazo[1,5-*a*]quinoline (3aj).

Yellow solid (47.4 mg, 85%), melting point 121-124°C; $^1\text{H-NMR}$ (400MHz, CDCl_3) δ 7.66-7.46 (m, 6H), 7.40-7.32 (m, 2H), 7.28-7.19 (m, 2H), 7.08 (d, $J=9.3\text{Hz}$, 1H); $^{13}\text{C-NMR}$ (100MHz, CDCl_3) δ 139.0, 135.3, 133.6, 132.4, 132.4, 131.0, 130.3, 129.8, 128.6, 127.9, 127.3, 125.3, 125.2, 122.3, 121.6, 117.0, 115.9. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2\text{Cl}$ 279.0687, found 279.0689.

1-(3-fluorophenyl)imidazo[1,5-*a*]quinoline (3ak).

Yellow solid (37.2 mg, 71%), melting point 102-105°C; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.66-7.65 (m, 1H), 7.57-7.45 (m, 4H), 7.41-7.35 (m, 3H), 7.28-7.22 (m, 2H), 7.08-7.04 (m, 1H); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 164.0-161.5 (d, $^1J_{\text{C-F}}=246.2\text{ Hz}$), 140.9, 135.8-135.7 (d, $^3J_{\text{C-F}}=8.3\text{ Hz}$), 132.1, 130.8, 130.4-130.3 (d, $^3J_{\text{C-F}}=8.3\text{ Hz}$), 128.7, 127.5, 125.6, 125.4 (d, $^4J_{\text{C-F}}=3.0\text{ Hz}$), 125.3, 122.6, 121.7, 117.3, 117.1, 116.8-116.6 (d, $^2J_{\text{C-F}}=22.3\text{ Hz}$), 116.4-116.2 (d, $^2J_{\text{C-F}}=21.0\text{ Hz}$). HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2\text{F}$ 263.0982, found 263.0985.

1-(2-fluorophenyl)imidazo[1,5-*a*]quinoline (3al).

Yellow solid (38.3 mg, 73%), melting point 102-104°C; $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.72-7.68 (m, 1H), 7.64-7.61 (m, 2H), 7.56-7.55 (m, 1H), 7.49 (d, $J=8.2\text{Hz}$, 1H), 7.38-7.31 (m, 3H), 7.28-7.24 (m, 2H), 7.05 (d, $J=9.2\text{Hz}$, 1H); $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 161.9-159.4 (d, $^1J_{\text{C-F}}=248.2\text{ Hz}$), 136.3, 132.4,

132.1 (d, $^4J_{C-F}=1.6$ Hz), 131.5-131.4 (d, $^3J_{C-F}=8.0$ Hz), 130.7, 128.5, 127.7, 125.3, 125.2, 124.7 (d, $^3J_{C-F}=3.5$ Hz), 122.6, 122.4-122.2 (d, $^2J_{C-F}=15.2$ Hz), 121.7, 116.9, 116.1-115.9 (d, $^2J_{C-F}=20.9$ Hz), 116.0. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{17}H_{12}N_2F$ 263.0984, found 263.0985.

1-(4-bromophenyl)imidazo[1,5-*a*]quinoline (3am).

Yellow solid (38.8 mg, 60%), melting point 138-140°C; 1H -NMR (400 MHz, $CDCl_3$) δ 7.69-7.63 (m, 3H), 7.57-7.53 (m, 4H), 7.38-7.33 (m, 2H), 7.28-7.25 (m, 1H), 7.08-7.03 (m, 1H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 141.1, 137.9, 132.7, 132.0, 131.1, 129.6, 128.7, 127.4, 125.6, 125.3, 123.5, 122.7, 121.6, 117.3, 117.2. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{17}H_{12}N_2Br$ 323.0183, found 323.0184.

1-(benzo[d][1,3]dioxol-5-yl)imidazo[1,5-*a*]quinoline (3an).

Yellow solid (23.6 mg, 41%), melting point 115-119°C; 1H -NMR (400 MHz, $CDCl_3$) δ 7.63-7.61 (m, 2H), 7.52 (s, 1H), 7.35-7.32 (m, 2H), 7.28-7.23 (m, 1H), 7.14 (dd, $J_1=7.9$ Hz, $J_2=1.4$ Hz, 1H), 7.09-7.08 (m, 1H), 7.02 (d, $J=9.3$ Hz, 1H), 6.97 (d, $J=7.9$ Hz, 1H), 6.09 (s, 2H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 148.5, 147.9, 141.9, 132.4, 130.4, 128.6, 127.3, 125.6, 125.1, 123.7, 122.2, 121.3, 117.4, 117.1, 110.0, 108.7, 101.4. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{18}H_{13}N_2O_2$ 289.0975, found 289.0977.

1-(naphthalen-1-yl)imidazo[1,5-*a*]quinoline (3ao).

Yellow solid (52.3 mg, 89%), melting point 199-201°C; 1H -NMR (400 MHz, $CDCl_3$) δ 8.09 (d, $J=8.1$ Hz, 1H), 7.98 (d, $J=8.2$ Hz, 1H), 7.75 (d, $J=6.9$ Hz, 1H), 7.71 (s, 1H), 7.68-7.60 (m, 2H), 7.53-7.50 (m, 1H), 7.44 (d, $J=9.4$ Hz, 1H), 7.41-7.33 (m, 2H), 7.24-7.21 (m, 1H), 7.09 (d, $J=9.3$ Hz, 1H), 7.01 (d, $J=8.0$ Hz, 1H), 6.95-6.91 (m, 1H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 140.4, 133.7, 132.6, 132.3, 131.6, 130.3, 130.0, 128.8, 128.4, 127.6, 127.1, 126.4, 125.6, 125.5, 125.4, 124.9, 122.5, 121.5, 117.2, 117.0. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{21}H_{15}N_2$ 295.1234, found 295.1235.

1-(furan-2-yl)imidazo[1,5-*a*]quinoline (3ap).

Yellow solid (27.1 mg, 58%), melting point 90-94°C; ¹H-NMR (400 MHz, CDCl₃) δ 7.67-7.64 (m, 2H), 7.57 (s, 1H), 7.40-7.33 (m, 3H), 7.22-7.20 (m, 1H), 7.09 (d, *J*=9.3Hz, 1H), 6.88 (d, *J*=3.2Hz, 1H), 6.68-6.67 (m, 1H); ¹³C-NMR (100 MHz, CDCl₃) δ 145.2, 143.2, 132.4, 132.1, 131.0, 128.6, 128.1, 125.4, 122.7, 122.2, 116.8, 112.8, 111.8. HRMS(ESI) *m/z*: [M+H]⁺ calcd for C₁₅H₁₁N₂O 235.0870, found 235.0871.

1-ethylimidazo[1,5-*a*]quinoline (3aq).

Yellow oil (24.1mg, 61%), ¹H-NMR (400 MHz, CDCl₃) δ 8.07 (d, *J*=8.4Hz, 1H), 7.54-7.52 (m, 1H), 7.43-7.39 (m, 1H), 7.30-7.27 (m, 2H), 7.18-7.15 (m, 1H), 3.31 (q, *J*=7.3Hz, 2H), 1.49 (t, *J*=7.3Hz, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 144.1, 132.1, 129.3, 127.5, 126.6, 124.7, 123.7, 119.6, 119.5, 116.3, 115.5, 24.8, 10.7. HRMS(ESI) *m/z*: [M+H]⁺ calcd for C₁₃H₁₃N₂ 197.1079, found 197.1083.

7-methyl-1-phenylimidazo[1,5-*a*]quinoline (3ba).

Yellow solid (33.5 mg, 65%), melting point 117–120 °C; ¹H-NMR (400 MHz, CDCl₃) δ 7.66 (s, 2H), 7.53 (s, 4H), 7.42 (s, 2H), 7.34-7.31 (m, 1H), 7.02-6.97 (m, 2H), 2.40 (s, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 142.1, 134.8, 133.9, 130.5, 130.3, 129.6, 129.2, 128.7, 128.5, 128.4, 125.6, 122.3, 121.3, 117.2, 117.1, 20.8. HRMS(ESI) *m/z*: [M+H]⁺ calcd for C₁₈H₁₅N₂ 259.1236, found 259.1235.

6,8-dimethyl-1-phenylimidazo[1,5-*a*]quinoline (3ca).

Yellow solid (35.4 mg, 65%), melting point 104–107 °C; ¹H-NMR (400 MHz, CDCl₃) δ 7.55-7.53 (m, 2H), 7.43-7.39 (m, 4H), 7.22-7.17 (m, 1H), 7.11-7.08 (m, 2H), 6.90 (s, 1H), 2.47 (s, 3H), 2.04 (s, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 142.2, 136.8, 135.3, 134.0, 132.5, 130.6, 129.5, 129.1, 128.6, 127.8, 121.9, 121.7, 117.6, 116.2, 115.7, 21.6, 19.9. HRMS(ESI) *m/z*: [M+H]⁺ calcd for C₁₉H₁₇N₂ 273.1393,

found 273.1392.

7-methoxy-1-phenylimidazo[1,5-*a*]quinoline (3da).

Yellow oil (17.0mg, 31%); ¹H-NMR (400 MHz, CDCl₃) δ 7.66 (d, *J*=3.5Hz, 2H), 7.55-7.53 (m, 4H), 7.46 (d, *J*=9.2Hz, 1H), 7.35 (d, *J*=9.3Hz, 1H), 7.07 (s, 1H), 6.98 (d, *J*=9.3Hz, 1H), 6.78 (dd, *J*₁=9.1Hz, *J*₂=2.2Hz, 1H), 3.86 (s, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 156.5, 141.9, 133.8, 130.3, 129.6, 129.2, 128.7, 126.9, 126.6, 122.5, 121.1, 118.6, 117.6, 114.8, 110.8, 55.5. HRMS(ESI) *m/z*: [M + H]⁺ calcd for C₁₈H₁₅N₂O 275.1183, found 275.1184.

7-ethoxy-1-phenylimidazo[1,5-*a*]quinoline (3ea).

Yellow solid (24.2 mg, 42%), melting point 113-116 °C; ¹H-NMR (400 MHz, CDCl₃) δ 7.65 (s, 2H), 7.55-7.52 (m, 4H), 7.44 (d, *J*=9.1Hz, 1H), 7.33 (d, *J*=9.2Hz, 1H), 7.06 (s, 1H), 6.96 (d, *J*=9.2Hz, 1H), 6.77 (d, *J*=8.3Hz, 1H), 4.07 (q, *J*=6.3Hz, 2H), 1.43 (t, *J*=6.6Hz, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 155.9, 141.9, 133.8, 130.3, 129.6, 129.2, 128.7, 126.9, 126.5, 122.5, 121.2, 118.6, 117.5, 115.3, 111.6, 63.8, 14.8. HRMS(ESI) *m/z*: [M+H]⁺ calcd for C₁₉H₁₇N₂O 289.1342, found 289.1341.

9-methoxy-1-phenylimidazo[1,5-*a*]quinoline (3fa).^{2d}

Yellow solid (36.7 mg, 67%), melting point 115–117 °C; ¹H-NMR (400 MHz, CDCl₃) δ 7.47 (s, 1H), 7.43-7.41 (m, 2H), 7.28-7.15 (m, 5H), 7.09 (d, *J*=8.0Hz, 1H), 6.82 (d, *J*=9.2Hz, 1H), 6.67 (d, *J*=8.0Hz, 1H), 2.90 (s, 3H); ¹³C-NMR (100 MHz, CDCl₃) δ 149.1, 146.7, 136.4, 131.7, 128.5, 128.0, 127.5, 125.9, 125.0, 122.9, 122.4, 120.8, 119.5, 117.7, 109.8, 53.6. HRMS(ESI) *m/z*: [M+H]⁺ calcd for C₁₈H₁₅N₂O 275.1179, found 275.1177.

1,5-diphenylimidazo[1,5-*a*]quinoline (3ga).

Yellow solid (26.2 mg, 41%), melting point 159-161 °C; ¹H-NMR (400 MHz, CDCl₃) δ 7.72-7.71 (m,

2H), 7.67 (d, $J=8.1$ Hz, 2H), 7.60 (s, 1H), 7.55-7.47 (m, 8H), 7.32-7.20 (m, 3H); ^{13}C -NMR (100 MHz, CDCl_3) δ 142.3, 138.8, 133.8, 133.4, 132.5, 130.2, 129.6, 129.5, 129.2, 128.8, 128.6, 127.8, 127.4, 127.3, 125.3, 124.9, 122.9, 117.8, 117.0. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{23}\text{H}_{17}\text{N}_2$ 321.1393, found 321.1392.

7-chloro-1-phenylimidazo[1,5-*a*]quinoline (3ha).

Yellow solid (40.1 mg, 72%), melting point 151-155 °C; ^1H -NMR (400 MHz, CDCl_3) δ 7.65-7.55 (m, 7H), 7.46 (d, $J=9.0$ Hz, 1H), 7.39 (d, $J=9.3$ Hz, 1H), 7.14 (dd, $J_1=9.0$ Hz, $J_2=2.0$ Hz, 1H), 6.96 (d, $J=9.4$ Hz, 1H); ^{13}C -NMR (100 MHz, CDCl_3) δ 142.6, 133.4, 130.8, 130.5, 130.3, 129.5, 129.5, 128.9, 127.7, 127.2, 127.1, 123.0, 120.3, 118.7, 118.5. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2\text{Cl}$ 279.0688, found 279.0689.

7-bromo-1-phenylimidazo[1,5-*a*]quinoline (3ia).^{2d}

Yellow solid (34.2 mg, 53%), melting point 128-130 °C; ^1H -NMR (400 MHz, CDCl_3) δ 7.76 (d, $J=1.6$ Hz, 1H), 7.65-7.63 (m, 2H), 7.59-7.55 (m, 4H), 7.41-7.35 (m, 2H), 7.29-7.25 (m, 1H), 6.96-6.91 (m, 1H). ^{13}C -NMR (100 MHz, CDCl_3) δ 142.7, 133.4, 131.2, 130.8, 130.3, 130.1, 129.6, 129.5, 129.0, 127.5, 123.1, 120.2, 119.0, 118.5, 118.3. HRMS(ESI) m/z : $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2\text{Br}$ 323.0178, found 323.0175.

7-fluoro-1-phenylimidazo[1,5-*a*]quinoline (3ja).

Yellow solid (27.2 mg, 52%), melting point 113–116 °C; ^1H -NMR (400 MHz, CDCl_3) δ 7.56-7.53 (m, 2H), 7.50-7.39 (m, 5H), 7.30 (d, $J=9.3$ Hz, 1H), 7.22-7.18 (m, 1H), 6.87 (d, $J=9.4$ Hz, 1H), 6.83-6.78 (m, 1H); ^{13}C -NMR (100 MHz, CDCl_3) δ 160.7-158.2 (d, $^1J_{\text{C-F}}=244.1$ Hz), 142.4, 133.5, 130.3, 129.6, 129.5, 129.0, 128.8 (d, $^4J_{\text{C-F}}=2.1$ Hz), 127.4 (d, $^3J_{\text{C-F}}=8.4$ Hz), 123.0, 120.6 (d, $^4J_{\text{C-F}}=2.5$ Hz), 119.1 (d, $^3J_{\text{C-F}}=8.1$

Hz), 118.5, 114.9-114.7 (d, $^2J_{C-F}$ =23.5 Hz), 113.8-113.5 (d, $^2J_{C-F}$ =22.3 Hz). HRMS(ESI) m/z: $[M+H]^+$
calcd for $C_{17}H_{12}N_2F$ 263.0986, found 263.0985.

7-nitro-1-phenylimidazo[1,5-a]quinoline (3ka).

Yellow solid (32.9 mg, 57%), melting point 200-203 °C; 1H -NMR (400 MHz, $CDCl_3$) δ 8.53 (d, J =2.4Hz, 1H), 8.0 (dd, J_1 =9.3Hz, J_2 =2.5Hz, 1H), 7.66-7.49 (m, 7H), 7.5 (d, J =9.4Hz, 1H), 7.12 (d, J =9.4Hz, 1H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 144.2, 143.5, 136.0, 132.9, 130.4, 130.0, 129.4, 129.2, 126.0, 123.8, 123.6, 121.9, 120.5, 119.6, 118.0. HRMS(ESI) m/z: $[M+H]^+$ calcd for $C_{17}H_{12}N_3O_2$ 290.0931, found 290.0930.

3,5-diphenylimidazo[1,5-c]quinazoline (3la).⁸

Yellow solid (23.1 mg, 36%), melting point 142-145 °C; 1H -NMR (400 MHz, $CDCl_3$) δ 8.14-8.04 (m, 2H), 7.94-7.92 (m, 1H), 7.60-7.57 (m, 2H), 7.35 (d, J =7.4Hz, 2H), 7.28-7.02 (m, 8H); ^{13}C -NMR (100 MHz, $CDCl_3$) δ 146.2, 142.7, 138.1, 133.8, 131.5, 130.3, 129.9, 129.2, 128.6, 128.4, 128.2, 128.0, 127.7, 127.4, 121.7, 120.6, 119.3. HRMS (ESI) m/z: $[M+H]^+$ calcd for $C_{22}H_{16}N_3$ 322.1339, found 322.1338.

ASSOCIATED CONTENT

Supporting Information

1H NMR and ^{13}C NMR spectra for all the products. This material is available free of charge *via* the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

E-mail: fengct2010@163.com

E-mail: nothingchina@126.com

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS:

We gratefully thank the financial supports of the Natural Science Foundation of China (81403268, 21102003), Scientific Research Foundation of Anhui University of Science & Technology (QN201508) and Scientific Research Foundation for the Introduction of Young Teachers.

REFERENCES

- (1) For selected examples, see: (a) Kim, D. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 2129. (b) Kakehi, S. H.; Okumura, Y.; Itoh, K.; Kobayashi, K.; Aikawa, Y.; Misawa, K. *Chem. Pharm. Bull.* **2010**, *58*, 1502.
- (c) Malamas, M. S.; Ni, Y.; Erdei, J.; Stange, H.; Schindler, R.; Lankau, H. J.; Grunwald, C.; Fan, K. Y.; Parris, K.; Langen, B.; Egerland, U.; Hage, T.; Marquis, K. L.; Grauer, S.; Brennan, J.; Navarra, R.; Graf, R.; Harrison, B. L.; Robichaud, A.; Kronbach, T.; Pangalos, M. N.; Hoefgen, N.; Brandon, N. J. *J. Med. Chem.* **2011**, *54*, 7621.
- (2) (a) Kolar, P.; Tišler, M. *Z. Naturforsch.* **1991**, *46b*, 1110. (b) Benincori, T.; Brenna, E.; Sannicolo, F. J. *Chem. Soc., Perkin Trans. I*, **1993**, 675. (c) Crawforth, J. M.; Paoletti, M. *Tetrahedron Lett.* **2009**, *50*, 4916. (d) Wang, Q.; Zhang, S.; Guo, F. F.; Zhang, B. Q.; Hu, P.; Wang, Z. Y. *J. Org. Chem.* **2012**, *77*, 11161. (e) Huang, J. R.; Zhang, Q. R.; Qu, C. H.; Sun, X. H.; Dong, L.; Chen, Y. C. *Org. Lett.* **2013**, *15*, 1878. (f) Li, Y. J.; Chao, A.; Fleming, F. F. *Chem. Commun.* **2016**, *52*, 2111. (g) Joshi, A.; Mohan, D. C.; Adimurthy, S. *Org. Lett.* **2016**, *18*, 464.
- (3) Bower, J. D.; Ramage, G. R. *J. Chem. Soc.* **1955**, 2823.
- (4) Li, M. Y.; Xie, Y.; Ye, Y.; Zou, Y.; Jiang, H. F.; Zeng, W. *Org. Lett.* **2014**, *16*, 6232.

- (5) Wang, H. Q.; Xu, W. T.; Wang, Z. Q.; Yu, L. T.; Xu, K. *J. Org. Chem.* **2015**, *80*, 2431.
- (6) (a) Yan, Y. Z.; Zhang, Y. H.; Zha, Z. G.; Wang, Z. Y. *Org. Lett.* **2013**, *15*, 2274. (b) Mohan, D. C.; Rao, S. N.; Ravi, C.; Adimurthy, S. *Org. Biomol. Chem.* **2015**, *13*, 5602.
- (7) Gong, L.; Xing, L. J.; Xu, T.; Zhu, X. P.; Zhou, W.; Kang, N.; Wang, B. *Org. Biomol. Chem.* **2014**, *12*, 6557.
- (8) Very recently, a nice work described the synthesis of imidazo[1,5-*c*]quinazolines from 4-methylquinazolines and benzylamines by Li's group; see: Zhao, D.; Wang, T.; Shen, Q.; Li, J. X. *Chem. Commun.* **2014**, *50*, 4302. However, when Li's approach was applied to our model substrates (2-methylquinoline **1a** and benzylamine **2a**), the desired product **3aa** was obtained in 16% yield.
- (9) Vanjari, R.; Singh, K. N. *Chem. Soc. Rev.* **2015**, *44*, 8062.
- (10) For selected reviews, see: (a) Yan, Y. Z.; Xu, K.; Fang, Y.; Wang, Z. Y. *J. Org. Chem.* **2011**, *76*, 6849. (b) Komai, H.; Yoshino, T.; Matsunaga, S.; Kanai, M. *Org. Lett.* **2011**, *13*, 1706. (c) Niu, R.; Xiao, J.; Liang, T.; Li, X. *Org. Lett.* **2012**, *14*, 676. (d) Liu, J. Y.; Niu, H. Y.; Wu, S.; Qu, G. R.; Guo, H. M. *Chem. Commun.* **2012**, *48*, 9723.
- (11) For recent reviews on sp² C-H amination, see: (a) Muller, P.; Fruit, C. *Chem. Rev.* **2003**, *103*, 2905. (b) Davies, H. M. L.; Long, M. S. *Angew. Chem., Int. Ed.* **2005**, *44*, 3518. (c) Davies, H. M. L.; Manning, J. R. *Nature* **2008**, *451*, 417. (d) Collet, F.; Lescot, C.; Dauban, P. *Chem. Soc. Rev.* **2011**, *40*, 1926. (e) Cho, S. H.; Kim, J. Y.; Kwak, J.; Chang, S. *Chem. Soc. Rev.* **2011**, *40*, 5068. (f) Louillat, M. L.; Patureau, F. W. *Chem. Soc. Rev.* **2014**, *43*, 901.
- (12) (a) Feng, C. T.; Su, J. H.; Yan, Y. Z.; Guo, F. F.; Wang, Z. Y. *Org. Biomol. Chem.* **2013**, *11*, 6691. (b) Feng, C. T.; Yan, Y. Z.; Zhang, Z. L.; Xu, K.; Wang, Z. Y. *Org. Biomol. Chem.* **2014**, *12*, 4837. (c)

Feng, C. T.; Zhu, H. Z.; Li, Z.; Luo, Z. G.; Wu, S. S.; Ma, S. T. *Tetrahedron Lett.* **2016**, *57*, 800.

(13) Yang, Y. Z.; Yao, J. Z.; Zhang, Y. H. *Org. Lett.* **2013**, *15*, 3206.

(14) Liu, F.; Liu, K.; Yuan, X. T.; Li, C. Z. *J. Org. Chem.* **2007**, *72*, 10231.

(15) Xiao, F. H.; Chen, S. Q.; Chen, Y.; Huang, H. W.; Deng, G. J. *Chem. Commun.* **2015**, *51*, 652.