

Published on Web 02/10/2007

Catalytic Asymmetric Rearrangement of α, α -Disubstituted α -Siloxy Aldehydes to Optically Active Acyloins Using Axially Chiral Organoaluminum Lewis Acids

Takashi Ooi,[†] Kohsuke Ohmatsu, and Keiji Maruoka*

Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo, Kyoto 606-8502, Japan

Received May 2, 2006; E-mail: maruoka@kuchem.kyoto-u.ac.jp

The skeletal rearrangements involving 1,2-carbon-to-carbon migration are fundamental yet powerful methods for the structural reorganization of organic molecules through the consecutive or concurrent cleavage and formation of carbon-carbon bonds,¹ often making it feasible to construct otherwise inaccessible molecular frameworks. However, precise control of the stereochemical outcome together with the migratory aptitude is not a trivial task mainly because of the proneness to generate an intermediary carbocation under the conventional conditions. Thus, research for the elaboration of stereoselective variants had long been circumvented. The delivery of effective Lewis acids to this field resulted in an important progress, enabling the stereospecific reaction using chiral substrates, which generally establishes complete chirality transfer with either stoichiometric or catalytic amount of an appropriate promoter.² Nevertheless, such endeavors are still very limited, and in turn, examples of success in the enantioselective 1,2-rearrangement of achiral substrates by the use of chiral activator, especially in a catalytic quantity, have been extremely rare despite its conceptual and practical significance.3 In conjunction with our recent studies on the organoaluminum-promoted selective rearrangement of aminocarbonyl compounds,⁴ we here report our own approach toward this subject, that is, the development of skeletal rearrangement of α, α -dialkyl- α -siloxy aldehydes 1, which can be efficiently catalyzed by newly designed chiral organoaluminum Lewis acid 3 with high enantioselectivity, thereby offering a facile access to various optically active acyloins (Scheme 1).5

On the basis of our preliminary studies on the search for suitable chiral Lewis acid catalysts with α,α -dibenzyl- α -trimethylsiloxy aldehyde (**1a**, R² = Me) as a representative substrate,⁶ we designed a new axially chiral organoaluminum Lewis acid **3** that possesses the chiral environment created by the 2-[3,5-bis(trifluoromethyl)-phenyl]-substituted naphthyl moiety being extended over the coordination site of the aluminum, possibly enabling rigorous stereocontrol.

The requisite ligand **4** can be synthesized from (*S*)-BINOL in a seven-step sequence.⁷ Interestingly, the molecular structure of **4** visualized by the single-crystal X-ray diffraction analysis adopted homochiral (*S*,*S*) configuration (Scheme 1, cylinder model),^{7,8} and refluxing a solution of **4** in toluene for 24 h showed no conformational interconversion between (*S*,*S*) and (*S*,*R*) isomers. Thus, treatment of (*S*,*S*)-**4** (1.1 equiv) with Me₃Al in toluene at room temperature for 30 min generated in situ the stereochemically defined methylaluminum catalyst (*S*,*S*)-**3** (10 mol %),⁹ and subsequent reaction with **1a** (R² = Me) at -20 °C for 12 h resulted in clean formation of the desired α -siloxy ketone **2a** (R² = Me) in 96% yield with 77% ee (entry 1 in Table 1). This promising result prompted us to evaluate the effect of the trialkylsilyl group on the stereoselectivity, and α -triethylsiloxy linkage in **1a** was found to

Table 1.	Catalytic Asymmetric Rearrangement o	νf
α,α-Dialk	$/1-\alpha$ -siloxy Aldehydes 1 with (S,S) - 3 ^a	

	$\frac{R^{2}_{3}SiO}{R^{1}} \xrightarrow{R^{1}} CHO \xrightarrow{(S,S)} \frac{(S,S)}{2}$	3 (5-10 mol%) toluene 0 °C, 12 h	0 R ¹ 0 2	∠R ¹ SiR² ₃	
entry	R ¹	R^{2}_{3}	% yield ^b	% ee ^c	product
1	PhCH ₂ (1a)	Me ₃	96	77	2a
2	$PhCH_2$ (1a)	Et ₃	99	87	2a
3	$PhCH_2$ (1a)	t-BuMe ₂	99	80	2a
4	$PhCH_2$ (1a)	Et_3	96	87	2a
5	p -MeO $-C_6H_4CH_2$ (1b)	Et ₃	98	85	2b
6	$p-F-C_{6}H_{4}CH_{2}(1c)$	Et ₃	97	90	2c
7	$p-Cl-C_6H_4CH_2$ (1d)	Et ₃	94	86	2d
8	2-NaphCH ₂ (1e)	Et ₃	95	85	2e
9	<i>trans</i> -PhCH=CHCH ₂ (1f)	Et ₃	81	83	2f
10	$(CH_3)_2C = CHCH_2 (1g)$	Et ₃	84	80	2g
11	$(CH_3)_2 CHCH_2 (1h)$	Et_3	94	74	2h
12	<i>c</i> -Hex (1i)	Et_3	97	74	2i

^{*a*} The reaction was carried out with either 10 mol % (entries 1–3) or 5 mol % (entries 4–12) of (S,S)-3 in toluene at -20 °C for 12 h. ^{*b*} Isolated yield. ^{*c*} Enantiopurity was determined by HPLC analysis. The absolute configuration of **2a** was determined to be *S* by comparison of the optical rotation with a literature value after desilylation.⁴

Scheme 1



be optimal, leading to the quantitative production of the rearranged **2a** ($R^2 = Et$) with 87% ee under similar conditions (entry 2). Notably, the catalyst loading can be reduced to 5 mol % without significant loss of reactivity and selectivity (entry 4).

With the optimized conditions in hand, we conducted the experiments to probe the scope of this new catalytic asymmetric 1,2-rearrangement, and the selected examples are included in Table 1.¹⁰ Generally, the reaction proceeded smoothly at -20 °C under the influence of 5 mol % of (*S*,*S*)-**3**. A series of α -siloxy aldehydes bearing α -benzylic substituents of different electronic properties were tolerated (entries 5–8). The substrates having allylic α -sub-

[†] Current address: Department of Applied Chemistry, Graduate School of Engineering, Nagoya University, Chikusa, Nagoya 464-8603, Japan.

Table 2. Kinetic Resolution of Differently α, α -Disubstituted α-Siloxy Aldehydes 5^a

Et ₃ SiO R ^{1 ^} 5	< ^{R²} сно						
entry	5	time (h)	% yield of 6 ^b	% ee ^{c,d} (config)	recovery of 5 (%) ^b	% ee ^{c,e} (config)	s
1	5a	12	49 (6a)	86 (S)	51	84 (<i>R</i>)	39.5
2	5a	15	55 (6a)	79 (S)	45	92 (R)	22.7
3	5b	11	49 (6b)	86	51	85	44.2
4	5c	12	51 (6c)	85	49	83	22.8
5	5d	3	57 (6d)	63	43	90	15.6
6	5e	0.5	55 (6e)	77	44	88	17.5

^{*a*} The reaction was conducted with 5 mol % of (S,S)-3 in toluene at -20 °C for the given reaction time. ^b Isolated yield. The ratio of 6 to the product through the migration of R^2 was >20:1. ^{*c*} Determined by HPLC analysis. d The absolute configuration was determined by comparison of the optical rotation with a reported value after desilylation.¹⁶ ^e For assignment of the absolute configuration, see Scheme 2.

Scheme 2



stituents also underwent highly enantioselective rearrangement, featuring the advantage of this method to construct enantiomerically enriched acyloins not readily accessible by the previously known methodologies (entries 9 and 10).⁵ It should be emphasized that extremely facile migration of a simple alkyl group was realized with good enantioselectivity (entries 11 and 12).

To further expand the scope of the present system, we investigated the rearrangement of racemic, differently α, α -disubstituted α -siloxy aldehydes 5, which uncovered the impressive level of kinetic resolution (Table 2),¹¹ making it feasible not only to broaden the applicability but also to prepare optically active tertiary α -hydroxy aldehydes.¹² When **5a** was treated with 5 mol % of (S,S)-3 in toluene at -20 °C for 12 h, α -hydroxy ketone 6a was obtained almost exclusively in 49% yield with 86% ee (S), and 5a was recovered in 51% yield with 84% ee (entry 1, selectivity factor: s = 39.5).¹³ Performing the reaction for 15 h under otherwise similar conditions enabled the isolation of 5a in 45% yield with 92% ee (entry 2). The absolute configuration of the recovered 5a was determined to be R by conversion to the diol 7 (Scheme 2), a known compound whose asymmetric synthesis was dependent upon enzymatic resolution.¹⁴ This assignment confirms the highly stereospecific rearrangement of (S)-5a to (S)-6a.15 Other examples listed in Table 2 showed the migratory aptitude and stereoselectivity corresponding to the combination of the two different α -substituents in 5. These results demonstrate potential synthetic utility of our approach and also suggest that the origin of the asymmetric induction lies on the ability of 3 to discriminate the α -stereogenic center of the substrates.

In summary, we have invented an asymmetric rearrangement of α,α -dialkyl- α -siloxy aldehydes to α -siloxy ketones efficiently catalyzed by the newly designed, axially chiral organoaluminum Lewis acid under mild conditions, and a kinetic resolution of racemic, differently α, α -disubstituted substrates has also been achieved. This is the first example of catalytic, highly enantioselective 1,2-rearrangement of α -alkoxycarbonyl compounds and provides a unique tool for the synthesis of various acyloins and tertiary α -hydroxy aldehydes of high enantiomeric purities. Further intensive studies on the application of this approach are currently underway in our laboratory.

Acknowledgment. This work was partially supported by a Grant-in-Aid for Scientific Research on Priority Areas "Advanced Molecular Transformation of Carbon Resources" from the Ministry of Education, Culture, Sports, Science and Technology, Japan. K.O. is grateful to the Japan Society for the Promotion of Science for Young Scientists for a Research Fellowship.

Supporting Information Available: Representative experimental procedures and spectroscopic characterization of new compounds (PDF); the crystallographic data for (S,S)-4 (CIF). This material is available free of charge via the Internet at http://pubs.acs.org.

References

- For reviews, see: Comprehensive Organic Synthesis; Trost, B. M., Fleming, I., Eds.; Pergamon Press: New York, 1991; Vol. 3, Chapter 3, pp 705-1014.
- Pinacol-type rearrangement: (a) Suzuki, K.; Katayama, E.; Tsuchihashi, G. Tetrahedron Lett. 1983, 24, 4997. (b) Suzuki, K.; Tomooka, K.; Shimazaki, M.; Tsuchihashi, G. Tetrahedron Lett. 1985, 26, 4781 and references therein. Epoxy ether rearrangement: (c) Maruoka, K.; Hasegawa, M.; Yamamoto, H.; Suzuki, K.; Shimazaki, M.; Tsuchihashi, G. J. Gawa, M., Tahahoo, 1986, 108, 3827. (d) Suzuki, K.; Miyazawa, M.; Tsuchihashi, G. *Tetrahedron Lett.* 1987, 30, 3515. (e) Maruoka, K.; Ooi, T.; Nagahara, S.; Yamamoto, H. *Tetrahedron* 1991, 47, 6983. (f) Suda, K.; Kikkawa, T.; Nakajima, S.-i.; Takanami, T. J. Am. Chem. Soc. 2004, 126, 0554. 126, 9554.
- (3) (a) Brunner, H.; Stöhr, F. Eur. J. Org. Chem. 2000, 2777. (b) Trost, B. M.; Yasukata, T. J. Am. Chem. Soc. 2001, 123, 7162. (c) See also: Trost, M., Tasukada, T.J. Am. Chem. Soc. 2006, 125, 7102. (c) See also. 110
 B. M.; Xie, J. J. Am. Chem. Soc. 2006, 128, 6044.
 (4) Ooi, T.; Saito, A.; Maruoka, K. J. Am. Chem. Soc. 2003, 125, 3220.
- (5) For asymmetric synthesis of α -hydroxy ketones, see oxidation by chiral phase-transfer catalysis: (a) Masui, M.; Ando, A.; Shioiri, T. Tetrahedron Lett. 1988, 29, 2835. Oxidation with chiral oxaziridines: (b) Davis, F. A.; Sheppard, A. C.; Chen, B.-C.; Haque, M. S. J. Am. Chem. Soc. **1990**, 112, 6679. Dihydroxylation of enol ethers: (c) Hashiyama, T.; Morikawa, K.; Sharpless, K. B. J. Org. Chem. **1992**, 57, 5067. Benzoin condensation: (d) Enders, D.; Kallfass, U. Angew. Chem., Int. Ed. 2002, 41, 1743. Nitroso aldol synthesis: (e) Momiyama, N.; Yamamoto, H. J. Am. Chem. Soc. 2003, 125, 6038. (f) Córdova, A.; Sundén, H.; Bøgevig, A.; Johansson, M.; Himo, F. Chem.–Eur. J. 2004, 10, 3673. (g) Hayashi, Y.; Yamaguchi, J.; Sumiya, T.; Hibino, K.; Shoji, M. J. Org. Chem. 2004, 69, 5966. (h) Wang, W.; Wang, J.; Li, H.; Liao, L. Tetrahedron Lett. 2004, 45, 7235. (i) Momiyama, N.; Yamamoto, H. J. Am. Chem. Soc. 2005, 127, 1080. Cross silvl benzoin reaction: (j) Linghu, X.; Potnick, J. R.; Johnson, J. S. J. Am. Chem. Soc. 2004, 126, 3070.
- (6) For the results, see the Supporting Information.
- (7) For detail, see the Supporting Information.
- (8) 4 was obtained as two separable diastereomers in a (S,S)/(S,R) ratio of
- (9) For the generation of a similar cyclic methylaluminum complex from Me₃-Al and 2-hydroxy-2'-(trifluoromethanesulfonylamino)biphenyl, see: Ooi, T.; Ichikawa, H.; Maruoka, K. Angew. Chem., Int. Ed. 2001, 40, 3610. The mode of aggregation of 3 in solution is unclear at present.
- (10) Unfortunately, the present system is not effective for symmetrically α_{α} . diaryl-substituted a-siloxy aldehydes.
- For a recent review on nonenzymatic kinetic resolution, see: Vedejs, E.; Jure, M. Angew. Chem., Int. Ed. 2005, 44, 3974
- (12) Tertiary α -hydroxy aldehydes are versatile chiral building blocks in natural product syntheses. See, for example: (a) Cossy, J.; Bauer, D.; Bellosta, V. *Tetrahedron* **2002**, *58*, 5909. (b) Kowashi, S.; Ogamino, T.; Kamei, J.; Ishikawa, Y.; Nishiyama, S. *Tetrahedron Lett.* **2004**, *45*, 4393.
- (13) The s values in this communication were calculated based on conversion and ee of 5 assuming first-order dependence on 5. For discussion on the validity of calculated *s* values, see: Keith, J. M.; Larrow, J. F.; Jacobsen, E. N. *Adv. Synth. Catal.* 2001, *343*, 5.
 (14) (a) Hof, R. P.; Kellog, R. M. *Tetrahedron: Asymmetry* 1994, *5*, 565. (b) Hof, R. P.; Kellog, R. M. *J. Chem. Soc., Perkin Trans. 1* 1996, 2051.
- (15) Attempted treatment of 5a with (S,S)-3 (10 mol %) in toluene at -20 °C
- for 48 h furnished 6a quantitatively with 8% ee (S).
- (16) Giordani, A.; Carera, A.; Pinciroli, V.; Cozzi, P. Tetrahedron: Asymmetry 1997, 8, 253.

JA063051Q