

CHEMISTRY A European Journal



Accepted Article Title: Metal-Free, NIS-Induced Regioselective Iodophosphoryloxylation of Alkenes with P(O)-OH bonds Authors: Biquan Xiong, Shipan Xu, Yu Zhu, Lei Yao, Congshan Zhou, Yu Liu, Kw-Wen Tang, and Wai-Yeung Wong This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain

the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Chem. Eur. J. 10.1002/chem.202000575

Link to VoR: http://dx.doi.org/10.1002/chem.202000575

Supported by ACES



Metal-Free, NIS-Induced Regioselective Iodophosphoryloxylation of Alkenes with P(O)-OH bonds

Biquan Xiong,^{*, a,b} Shipan Xu, ^a Yu Zhu, ^a Lei Yao, ^a Congshan Zhou, ^a Yu Liu, ^a Ke-Wen Tang,^{*, a} Wai-Yeung Wong^{*, b}

Abstract: A simple and efficient method for the regioselective iodophosphoryloxylation of alkenes with P(O)-OH bonds has been established by using NIS (N-iodosuccinimide) as the iodination reagent under transition-metal-free conditions. The present protocol is compatible with different functional groups, and suitable for various alkenes and P(O)-OH compounds. A variety of functionalized β -iodo-1-ethyl phosphinic/phosphoric acid esters are obtained in good to excellent yields, which could be further transformed to diversified building blocks for the synthesis of bioactive compounds, pharmaceuticals and functional materials.

Introduction

Organophosphorus compounds have been recognized as the important intermediates in biological chemistry, asymmetric catalysis, and preparation of functional materials for a half century, due to their unique structural features and potential pharmacological activities.^{1,2} In particular, some β -iodo- α -alkyl phosphinic/phosphoric acid esters are significant building blocks present in a variety of biologically interesting natural products, pharmaceuticals, and synthetic intermediates of broad utility. Generally, these β-iodofunctionalized phosphinic/phosphoric acid esters are prepared either by the substitution of phosphoryl chlorides with β -iodo- α -ethanol compounds/direct iodination for the methyl group of 1-phenylethyl phosphinic or phosphoric acid esters or the direct cross coupling of (1-halo-2-iodo-alkyl) compounds with phosphinic/phosphoric acids and transition-metalcatalyzed [e.g., Fe, Cu, Pd] dehydrogenative coupling of β-iodo-αethanol derivatives with P(O)-H/P(O)-Cl bonds.³ However, these protocols suffer from certain disadvantages such as the use of a large excess of P(O)-H motifs and co-oxidants, poor regioselectivity, and low yields.

The direct dehydrogenative coupling of P(O)-H or P(O)-OH bonds with $C(sp^3)$ -H bonds of the methyl group in arenes has been achieved by Zhao and Xiong.^{5a-b} In 2015, Wei and Wang et al. established a novel and efficient procedure for the direct difunctionalization of alkenes with I₂O₅ and P(O)-H compounds.^{5c} Recently, Ishihara discovered a metal-free transesterification of phosphinates with alcohols via the catalysis of tetramethylammonium methyl carbonate.^{5d} Jacobsen et al. reported the intramolecular reaction of carboxylic acids with pendant olefins

 [a] Department of Chemistry and Chemical Engineering, Hunan Institute of Science and Technology, Yueyang, 414006, P.R.China. xiongbiquan@126.com (Dr. B. Xiong) tangkewen@sina.com (Prof. K. Tang)

[b] Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hung Hom, Hong Kong, P. R. China. wai-yeung.wong@polyu.edu.hk (Prof. W.-Y. Wong) Supporting information for this article is given via a link at the end of the document in the presence of a source of I^+ for the enantioselective generation of five- and six-membered lactones.^{4a} Since then, Fujioka et al. and Yeung et al. further demonstrated the enantioselective bromolactonization of 5-substituted 5-hexenoic acids catalyzed by the tris(imidazoline) and L-proline-derived S-alkyl thiocarbamate, respectively.4b,4c Feng et al. found a highly efficient catalytic chloroamination reaction of α,β -unsaturated γ -keto esters and chalcones via a chloronium-based mechanism to deliver antiregioselective vicinal chloroamines instead of the aziridinium intermediates delivered aminochlorides.^{4d} In 2012, Toste et al. invented a chiral anion phase-transfer system for enantioselective 6-exo-trig bromocyclization of styrenyl amides.^{4e} Later, Mukherjee et al. discovered the catalytic asymmetric iodoetherification of oximes with the help of a bifunctional thiourea catalyst. $^{\rm 4f}$ In addition, Denmark and Burk have systematically studied the Lewis base catalyzed bromo- and iodolactonization reactions and also explored the effects of catalyst structure on the rate and cyclization selectivity.^{4g} Although some elegant studies on the selective difunctionalization of carbon-carbon double bonds were achieved by these scientists,⁵ the use of P(O)-OH compounds as starting materials has not been reported. Herein, we demonstrate an efficient regioselective iodophosphoryloxylation of alkenes with P(O)-OH bonds by using NIS (N-iodosuccinimide) as the iodination reagent under transition-metal-free conditions (Scheme $1).^{6}$





Results and Discussion

At the first stage of our study, we attempted to use NIS as the activation reagent for the carbon-carbon double bond for the generation of 2-phenyliodiran-1-ium, and then introduced a P(O)-OH bond to realize the regioselective iodophosphoryloxylation of alkenes. To test our initial hypothesis, the reaction of styrene (1a) and diphenylphosphinic acid (2a) was investigated to delineate the reaction parameters.⁷ The reaction of 1a with 2a was carried out at room temperature in toluene under a N₂ atmosphere with the addition of iodine reagent NIS, and the corresponding addition

product of 2-iodo-1-phenylethyl diphenylphosphinate (3a) was generated in 7% yield. Besides toluene, other solvents, such as THF, DMF, CH₃CN, 1,4-dioxane and CH₂Cl₂ were further tested (Table 1, entries 2-6), and THF gave the product in a preferable yield of 78%. We then chose THF as the solvent for further optimization. When the reaction was operated under an air atmosphere, the expected product could only be produced in 66% yield (Table 1, entry 7). This phenomenon may be ascribed to the influence of water in air, which interrupted the phosphoryloxylation path of the reaction. Besides NIS, other iodine sources such as I₂, KI, NaI and tetrabutylammonium iodide (TBAI) showed negative results toward the reaction (Table 1, entries 8-11). The increase of reaction temperature within the 25-40 °C range is beneficial, but a further increase from 40 to 80 °C results in a little decrease of the product yield (Table 1, entries 12-14). Additionally, it is worth noting that the reaction would not produce any products in the absence of NIS (Table 1, entry 15). We further optimized the amount of NIS used, and the optimal condition was obtained at a "diphenyl phosphinic acid/styrene/NIS molar ratio" of 1:1:1.2 (Table 1, entries 16-18). Some other iodination systems were further screened for the reaction, and I₂ still showed negative results for the reaction even with the addition of an oxidant (H_2O_2). TBHP, DTBP, K₂S₂O₈). According to M. C. Mattos's method, TICA (triiodoisocyanuric acid) was synthesized in 86% yield, 7e which could promote the reaction with the generation of the desired product in 82% yield.

Table 1. Optimization of the reaction conditions.^a

	+ P	O h−P−OH iodine reage Ph	ent int	O P P Ph
	1a	2a	3	а
Entry	Solvent	Iodine reagent	Temp.	Yield of 3a ^b
1	toluene	NIS	r.t.	7%
2	THF	NIS	r.t.	78%
3	DMF	NIS	r.t.	49%
4	CH ₃ CN	NIS	r.t.	58%
5	1,4-dioxane	NIS	r.t.	23%
6	CH_2Cl_2	NIS	r.t.	65%
7	THF	NIS	r.t.	66% ^c
8	THF	I_2	r.t.	N.D.
9	THF	KI	r.t.	N.D.
10	THF	NaI	r.t.	N.D.
11	THF	TBAI	r.t.	N.D.
12	THF	NIS	40 °C	95%
13	THF	NIS	60 °C	93%
14	THF	NIS	80 °C	92%
15	THF	-	40 °C	N.D.
16	THF	NIS	40 °C	$22\%^{d}$
17	THF	NIS	40 °C	46% ^e
18	THF	NIS	40 °C	96% ^f
19	THF	I_2/H_2O_2	40 °C	N.D. ^g
20	THF	$I_2/TBHP$	40 °C	N.D. ^h
21	THF	I ₂ /DTBP	40 °C	N.D. ^{<i>i</i>}
22	THF	$I_2/K_2S_2O_8$	40 °C	N.D. ^j
23	THF	TICA	40 °C	82% ^k

١.

^a Reagents and conditions: styrene (1a, 0.5 mmol), diphenylphosphinic acid (2a, 0.5 mmol) and iodine reagent (1.0 eq) in solvent (1.0 mL), N₂ atmosphere, temp., 6 h. ^{b 31}P NMR yields. ^c Air conditions. ^d NIS (0.125 mmol, 25 mol%). ^e NIS (0.25 mmol, 50 mol%). ^f NIS (0.6 mmol, 120 mol%). ^g H₂O₂ (0.6 mmol, 25% solution in water). ^h TBHP (0.6 mmol, 70% solution in water). ⁱ DTBP (di-*t*-butyl peroxide, 0.6 mmol). ^j K₂S₂O₈ (0.6 mmol). ^k TICA (triiodoisocyanuric acid, 0.6 mmol).

As shown in **Table 2**, the regioselective iodophosphoryloxylation of alkenes with NIS and P(O)-OH bonds leading to β -iodophosphates can be applied to different kinds of alkenes. It is clear that styrene (**1a**), 1-methyl-4-vinylbenzene (**1b**), 1-bromo-4-vinylbenzene (**1c**), 1-bromo-3-vinylbenzene (**1d**), 1-

bromo-2-vinylbenzene (1e) and 1-fluoro-4-vinylbenzene (1f) can react efficiently with diphenylphosphinic acid (2a) under the optimized reaction conditions, affording the corresponding iodophosphoryloxylation products of 3a-3f in 85-95% isolated yields. For most cases, electron-donating or electron-withdrawing groups which are located on the aryl ring of styrenes do not change the yields of the products significantly. In addition, special alkenes such as allylbenzene (2g) and 1-allylnaphthalene (2h) could also afford the desired products of 3g and 3h in 58% and 45% yields, respectively.8 To our delight, when cyclic alkenes (2i-2k) were employed as the starting materials, 3i-3k were synthesized in 75-82% yield. In addition, straight-chain alkenes such as cinnamyl alcohol (11), and methyl cinnamate (1m) also showed positive results toward the reaction, and the expected products of 31 and 3m were generated in 86% and 90% yields, respectively. Furthermore, various kinds of alkenes containing ester group (10-1t) also exhibited high reactivities for the present reaction, affording the corresponding addition products in 63-94% yields. While 1methoxy-4-vinylbenzene (1u) and (E)-1-methoxy-4-(prop-1-en-1yl)benzene (1v) were used for the reaction, the desired iodophosphoryloxylation products can be synthesized in 91% and 80% ³¹P NMR yields, but unfortunately, these compounds could not be isolated from the reaction mixture via the short silicacolumn. It is deduced that these compounds might decompose during the purification process.

Table 2. Scope of alkenes. ^a



^{*a*} Reagents and conditions: alkene (**1a**, 0.5 mmol), diphenylphosphinic acid (**2a**, 0.5 mmol) and NIS (0.6 mmol) in THF (1.0 mL), at 40 °C under a N₂ atmosphere stirred for 6 h. ^{*b*} Isolated yields. ^{*c*} ³¹P NMR yield, light sensitive, cannot be separated and decomposed during passing through the silica gel column.

As depicted in **Table 3**, a range of substituted P(O)-OH compounds (**2b-2o**) were screened under the optimized reaction conditions with styrene. It is clear that symmetrically substituted

diarylphosphinic acid with electron-donating groups such as di-p-tolylphosphinic acid (**2b**), di-m-tolylphosphinic acid (**2c**), bis(3,5dimethylphenyl)phosphinic acid (**2d**) and bis(4methoxyphenyl)phosphinic acid (**2e**), exhibit high reactivity toward styrene, giving the corresponding products of **4a-4d** in 83-92% yields. For large functional groups in substituted diarylphosphinic acids, such as di(naphthalen-1-yl)phosphinic acid (**2f**) and di(naphthalen-2-yl)phosphinic acid (**2g**), they also showed positive results, affording the corresponding products in 80% and 78% yields.

When bis(3-fluorophenyl)phosphinic acid (2h) was used, the corresponding iodophosphoryloxylation product of 2-iodo-1phenylethyl bis(3-fluorophenyl)phosphinate (4g) could be produced in 68% yield. Moreover, asymmetric phosphonic acid, such as methyl(phenyl)phosphinic acid (2i), is also suitable for the reaction and yielded the product of 4h in 85% yield. Notably, different kinds of hydrogen phosphates such as diethyl hydrogen phosphate (2j), dibutyl hydrogen phosphate (2k), 2-ethylhexyl (2ethylpentyl) hydrogen phosphate (21), dibenzyl hydrogen phosphate (2m) and diphenyl hydrogen phosphate (2n) were also appropriate for this protocol to generate the corresponding products (4i-4m) in moderate to good yields. When (S)-(+)-1,1'-binaphthyl-2.2'-divl hydrogenphosphate (20) was used for the reaction, despite its low solubility and reactivity as a nucleophile, in this case it was quite effective with 53% yield for the synthesis of (11bS)-4-(2iodo-1-phenylethoxy)dinaphtho[2,1-d:1',2'-f][1,3,2] dioxaphosphepine 4-oxide (4n) as the desired product. The fact of substituent insensibility in alkenes to the reaction yields shows that substituents probably do not influence the rate determining step of the reaction. Furthermore, the difference in the yields of phosphates and phosphonates indicates that the phosphonates are more reactive than phosphates due to the pKa differences.





^{*a*} Reagents and conditions: styrene (1, 0.5 mmol), P(O)-OH compounds (0.5 mmol), NIS (0.6 mmol), THF (1.0 mL), under N₂ atmosphere, 40 $^{\circ}$ C , 6 h. ^{*b*} Isolated yields.

In order to demonstrate the practical application of this method, we further performed a large-scale reaction of styrene (1a, 20 mmol) with diphenyl phosphinic acid (2a, 20 mmol) in the presence of NIS which afforded 3a in 81% yield (7.257 g) (Scheme 2). In the solvent of DMF, 3a could react efficiently with

sodium azide to afford 2-azido-1-phenylethyl diphenylphosphinate (**5a**) in 96% yield.⁹ This compound provides great synthetic versatility to the phosphorus derivatives, allowing a variety of subsequent structural modification such as radical reaction for the synthesis of triazolyl-functionalized phosphinates with distinct biological activity. In addition, we further conducted the reaction of **3a** with water in the presence of silver nitrate (2.0 equiv.), and the corresponding hydroxylation product of 2-hydroxy-1-phenylethyl diphenylphosphinate (**5b**) was obtained in 99% yield.⁹



Scheme 2. Large-scale production and selective functionalization of 3a.



Scheme 3. Control experiments.

Control experiments were conducted for the reaction in order to gain the insight of the reaction mechanism. As depicted in Scheme 3, while the reactions were only performed with diphenyl phosphinic acid (2a) and NIS under the optimized reaction conditions, the desired coupling product was not detected after the reaction. Meanwhile, while we employed H₂O instead of diphenyl phosphinic acid in the reaction, the corresponding product of 2iodo-1-phenylethanol (6b) could be obtained in 76% yield.¹⁰ Besides diphenyl phosphinic acid, diphenyl phosphine chloride and diphenylphosphinic iodide were also tested for the reaction, but unfortunately, the desired adducts were not formed during the reaction. A competitive reaction between diphenyl phosphinic acid (2a) and water was operated under the optimized reaction conditions, and 2-iodo-1-phenylethyl diphenylphosphinate (3a) and 2-iodo-1-phenylethanol (6b) were generated in 72% and 21% vields, respectively. It is deduced that diphenyl phosphinic acid (2a) has a higher reactivity than water as a nucleophile.

WILEY-VCH



Scheme 4. Plausible mechanism.

A plausible mechanism for the present regioselective iodophosphoryloxylation reaction is proposed as illustrated in **Scheme 4**.¹¹ The *N*-iodosuccinimide (**B**) first undergoes the attack of styrene (**A**) leading to the formation of iodonium intermediate **D** with the release of pyrrolidine-2,5-dione anion (**C**). In the presence of P(O)-OH compound (**E**), which could easily operate the ringopening reaction with **D** to form the unstable corresponding oxonium intermediate (**F**). In the presence of pyrrolidine-2,5-dione anion (**C**), **F** could undergo the deprotonation process to give the iodophosphoryloxylation product **H** with the release of one molecule of pyrrolidine-2,5-dione (**G**).

Conclusion

In summary, we have developed an efficient NIS-induced regioselective iodophosphoryloxylation of unactivated alkenes with P(O)-OH bonds under transition metal-free conditions. A wide range of P(O)-OH compounds and alkenes bearing diverse functional groups were applicable for this protocol, providing the related β -iodo-1-ethyl phosphinic/phosphoric acid esters with good to excellent yields. To the best of our knowledge, it appears that the regioselective iodophosphoryloxylation of alkenes with P(O)-OH bonds has not been exploited previously, and the salient features of the reaction include its broad substrate scope, high step economy, and good chemoselectivity. The synthetic method also exhibits high potential for the construction of biologically active molecules, and organophosphorus compounds

Acknowledgement

This work was supported by National Natural Science Foundation of China (21606080), Natural Science Foundation of Hunan Province (2019JJ50203), Scientific Research Fund of Hunan Provincial Education Department (19A197) and Hunan Provincial Innovation Foundation for Postgraduate (CX2018B774). W.-Y.W. thanks the Hong Kong Polytechnic University (1-ZE1C) and the Endowed Professorship in Energy from Ms Clarea Au (847S) for the financial support.

Conflicts of interest

There are no conflicts to declare

Keywords: P(O)-OH bonds • Iodophosphoryloxylation • alkenes • transition metal-free

 a) H. Onouchi, T. Miyagawa, A. Furuko, K. Maeda, E. Yashima, J. Am. Chem. Soc. 2005, 127, 2960-2965; b) J. Yang, T. Q. Chen, L. B. Han, J. Am. Chem. Soc. 2015, 137, 1782-1785.

- [2] a) Y. Zhu, T. Q. Chen, S. Li, S. Shimada, L. B. H J. Am. Chem. Soc. 2016, 138, 5825-5828; b) V. Quint, F. Morlet-Savary, J. F. Lohier, J. Lalevée, A. C. Gaumont, J. Am. Chem. Soc. 2016, 138, 7436-7441.
- [3] a) P. Cheruku, J. Diesen, P. G. Andersson, J. Am. Chem. Soc. 2008, 130, 5595-5599; b) D. J. Fox, D. S. Pedersen, A. B. Petersen, S. Warren, Org. Biomol. Chem. 2006, 4, 3117-3119; c) L. Deng, Y. Wang, H. Mei, Y. Pan, J. L. Han, J. Org. Chem. 2019, 84, 949-956; d) M. Engman, P. Cheruku, P. Tolstoy, J. Bergquist, S. F. Völker, P. G. Anderssona, Adv. Synth. Catal. 2009, 351, 375-378.
- [4] a) G. E. Veitch, N. Eric Jacobsen, Angew. Chem. Int. Ed. 2010, 49, 7332-7335; b)
 K. Murai, T. Matsushita, A. Nakamura, S. Fukushima, M. Shimura, H. Fujioka, Angew. Chem. Int. Ed. 2010, 49, 9174-9177; c) X. Jiang, C. K. Tan, L. Zhou, Y.
 Y. Yeung, Angew. Chem. Int. Ed. 2012, 51, 7771-7775; d) Y. Cai, X. Liu, J. Jiang,
 W. Chen, L. Lin, X. Feng, J. Am. Chem. Soc. 2011, 133, 5636-5639; e) Y. M.
 Wang, J. Wu, C. Hoong, V. Rauniyar, F. D. Toste, J. Am. Chem. Soc. 2012, 134, 12928-12931; f) C. B. Tripathi, S. Mukherjee, Angew. Chem. Int. Ed. 2013, 52, 8450-8453; g) S. E. Denmark, M. T. Burk, PNAS. 2010, 107, 20655-20660.
- [5] a) L. Zhou, D. W. Tay, J. Chen, G. Y. C. Leung, Y. Y. Yeung, *Chem. Commun.* 2013, 49, 4412-4414; b) J. Chen, L. Zhou, Y.Y. Yeung, *Org. Biomol. Chem.* 2012, 10, 3808-3811; c) A. Alix, C. Lalli, P. Retailleau, G. Masson, *J. Am. Chem. Soc.* 2012, 134, 10389-10392; d) D. H. Paull, C. Fang, J. R. Donald, A. D. Pansick, S. F. Martin, *J. Am. Chem. Soc.* 2012, 134, 11128-11131.
- [6] a) J. Xu, P. Zhang, X. Li, Y. Gao, J. Wu, G. Tang, Y. Zhao, Adv. Synth. Catal. 2014, 356, 3331-3335; b) B. Xiong, G. Wang, C. Zhou, Y. Liu, P. Zhang, K. Tang, J. Org. Chem. 2018, 83, 993-999; c) C. Liu, M. Zhu, W. Wei, D. Yang, H. Cui, X. Liu, H. Wang, Org. Chem. Front. 2015, 2, 1356-1360; d) M. Hatano, Y. Tabata, Y. Yoshida, K. Toh, K. Yamashita, Y. Ogura, K. Ishihara, Green Chem. 2018, 20, 1193-1198.
- [7] a) R. Suresh, A. K. Simlandy, S. Mukherjee, Org. Lett. 2018, 20, 1300-1303; b)
 C. B. Tripathi, S. Mukherjee, Angew. Chem. Int. Ed. 2013, 52, 8450-8453; c) C.
 S. Brindle, C. S. Yeung, E. N. Jacobsen, Chem. Sci. 2013, 4, 2100-2104; d) J. E.
 Tungen, J. M. J. Nolsøe, T. V. Hansen, Org. Lett. 2012, 14, 5884-5887; e) R. S.
 Ribeiro, P. M. Esteves, M. C. Mattos, Tetrahedron Lett. 2007, 48, 8747-8751.
- [8] Due to the special intrinsic property of this type of compounds, a cyclization reaction would occur during the reaction with the generation of a cyclic phosphorus compounds, leading to the low yields of 3g and 3h. The study for the cyclization reaction is under way.
- [9] a) X. Yang, W. Liu, L. Li, W. Wei, C. J. Li, *Chem.-Eur. J.* 2016, *22*, 15252-15256;
 b) D. S. Rao, T. R. Reddy, K. Babachary, S. Kashyap, *Org. Biomol. Chem.* 2016, *14*, 7529-7543.
- [10] a) M. K. Agrawal, S. Adimurthy, B. Ganguly, P. K. Ghosh, *Tetrahedron*. 2009, 65, 2791-2797.
- [11] a) Q. Xue, J. Xie, P. Xu, K. Hu, Y. Cheng, C. Zhu, ACS Catal. 2013, 3, 1365-1368; b) R. Beltran, S. Nocquet-Thibault, F. Blanchard, R. H. Dodd, K. Cariou, Org. Biomol. Chem. 2016, 14, 8448-8451; c) H. Nakatsuji, Y. Sawamura, A. Sakakura, K. Ishihara, Angew. Chem. Int. Ed. 2014, 53, 6974-6977; d) A. N. French, S. Bissmire, T. Wirth, Chem. Soc. Rev. 2004, 33, 354-362.

WILEY-VCH

Entry for the Table of Contents



A simple and efficient method for the regioselective iodophosphoryloxylation of alkenes with P(O)-OH bonds has been established by using NIS (*N*-iodosuccinimide) as the iodination reagent under transition-metal-free and mild conditions. A wide range of P(O)-OH compounds and alkenes bearing diverse functional groups were applicable for this protocol. The synthetic method also exhibits high potential for the construction of biologically active molecules, and organophosphorus compounds.