

Communication

**Photoredox-Mediated #-Vinylolation of #-Amino Acids and N-Aryl Amines**

Adam Noble, and David W. C. MacMillan

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • DOI: 10.1021/ja506094d • Publication Date (Web): 14 Jul 2014

Downloaded from <http://pubs.acs.org> on July 17, 2014

**Just Accepted**

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



## Photoredox-Mediated $\alpha$ -Vinylolation of $\alpha$ -Amino Acids and *N*-Aryl Amines

Adam Noble and David W. C. MacMillan\*


Merck Center for Catalysis at Princeton University, Princeton, New Jersey 08544  
dmacmill@princeton.edu

RECEIVED DATE

**Abstract.** A new coupling protocol has been developed that allows the union of vinyl sulfones with photoredox-generated  $\alpha$ -amino radicals to provide allylic amines of broad diversity. Direct C–H vinylations of *N*-aryl tertiary amines, as well as decarboxylative vinylations of *N*-Boc  $\alpha$ -amino acids, proceed in high yield and with excellent olefin geometry control. The utility of this new allyl amine forming reaction has been demonstrated via the syntheses of several natural products and a number of established pharmacophores.

Visible light photoredox catalysis has recently emerged as a powerful activation strategy for the discovery and invention of new chemical transformations.<sup>1</sup> The capacity of metal-to-ligand charge transfer (MLCT) catalysts to simultaneously act as strong reductants and oxidants under the influence of low energy photon sources has enabled the design of many novel reaction mechanisms that can be chemoselectively triggered using visible light. One emerging application of photoredox catalysis is its use in the direct functionalization of unactivated  $sp^3$  C–H bonds,<sup>2</sup> an area of research that has become a fundamental goal of modern organic chemistry.<sup>3</sup> In this context, our laboratory has become interested in the activation of C–H bonds to generate  $\alpha$ -amino radicals, which are versatile intermediates that can participate in radical–radical couplings<sup>4</sup> to afford benzylic amines (Eq 1) or be trapped with radical acceptors to forge  $\alpha$ -*N*-alkylation adducts.<sup>5,6</sup> Generation of  $\alpha$ -amino radicals under photoredox conditions typically proceeds via single-electron oxidation of an amine and subsequent deprotonation of the resulting radical cation.<sup>4,5</sup> Recently, our group has also reported


**Scheme 1.** Proposed Mechanism for the Vinylolation Reaction.




an alternative strategy for  $\alpha$ -amino radical formation via the decarboxylation of *N*-*tert*-butoxycarbonyl (*N*-Boc)  $\alpha$ -amino acids,<sup>7,8</sup> a CO<sub>2</sub>-extrusion mechanism that has implications for the use of biomass feedstocks in conjugate additions and organometallic couplings.

Allylic amines have long been attractive targets for new reaction development due to their importance as (i) functionalized building blocks<sup>9</sup> and (ii) versatile intermediates in the production of medicinal agents.<sup>10</sup> Indeed, over the last three decades, numerous methods have been developed for the preparation of allyl amines, typically proceeding through the addition of vinyl metal reagents to C=N bonds<sup>11</sup> or the direct amination of allylic substrates.<sup>12</sup> Moreover, deprotonation-based methods for the direct  $\alpha$ -vinylolation of nitrogen centers have been reported via the  $\alpha$ -lithiation, transmetalation of tertiary carbamates, followed by metal-catalyzed cross-couplings with vinyl halides.<sup>3d,13</sup>

Taking inspiration from our previously developed photoredox arylation protocols (Eq 1),<sup>4,7</sup> we sought to extend the utility of  $\alpha$ -amino radicals to the direct synthesis of allylic amines from *N*-aryl amines or  $\alpha$ -amino acids. Specifically, we hypothesized that allyl amines should be accessible via exposure of  $\alpha$ -amino radicals to olefins that can generically participate in radical-based vinylolation mechanisms (i.e. incorporate leaving groups that are susceptible to single-electron  $\beta$ -elimination pathways). As an overarching goal of this work we hoped to demonstrate that photoredox-generated  $\alpha$ -amino radicals provide a complementary approach to allyl amine production in comparison to stoichiometric metal-based technologies. Herein, we describe the successful execution of these ideals and present



**Table 1.** Initial Studies Towards the C–H Vinylation Reaction.


entry	photocatalyst	base	solvent	yield <sup>a</sup>	<i>E:Z</i>
1	Ir(ppy) <sub>3</sub>	NaOAc	DMA	28%	21:79
2	Ir(ppy) <sub>3</sub>	NaOAc	toluene	57%	91:9
3	<b>11</b>	NaOAc	toluene	33%	>98:2
4	<b>1</b>	NaOAc	toluene	40%	95:5
5	<b>1</b>	CsOAc	toluene	81%	96:4
6 <sup>b</sup>	<b>1</b>	CsOAc	toluene	87%	97:3
7 <sup>b</sup>	<b>1</b>	CsOAc	DCE	91%	98:2
8 <sup>b</sup>	none	CsOAc	DCE	<5%	-
9 <sup>b,c</sup>	<b>1</b>	CsOAc	DCE	0%	-

<sup>a</sup>Determined by <sup>1</sup>H NMR analysis using an internal standard. <sup>b</sup>Performed with 3.0 equiv. base and at 0.1 M in solvent. <sup>c</sup>Performed in the absence of light. **1** = Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub>, **11** = Ir(ppy)<sub>2</sub>(dtbbpy)PF<sub>6</sub>.

the first direct C–H vinylation of *N*-aryl tertiary amines and the first decarboxylative olefination of *N*-Boc  $\alpha$ -amino acids using vinyl sulfones in the presence of iridium-based MLCT catalysts (Eq 2).


**Reaction Design.** Critical to our proposal of generating allylic amines from  $\alpha$ -amino radicals was the identification of an appropriate alkene SOMO-phile, an olefinic coupling partner that would be susceptible to radical addition yet could further participate in single-electron  $\beta$ -elimination to deliver the required vinylation adduct. In this context, Nozaki and coworkers, among others, have ingeniously demonstrated that unsaturated sulfones can generically react with alkyl radicals to produce a wide variety of olefin containing products.<sup>14,15</sup> For our purposes, we presumed that vinyl sulfones should be useful substrates for the proposed photoredox vinylation given that (i) they are electron-deficient olefins which should readily couple with nucleophilic  $\alpha$ -amino radicals and (ii) the radical C–C bond forming step should be rapidly followed by  $\beta$ -sulfone elimination to produce a sulfinyl radical, a species that should rapidly undergo single-electron reduction via the reduced state of the photocatalyst. Furthermore, with respect to substrate availability, we recognized that vinyl sulfones are easily prepared in one step from aldehydes using Horner–Wadsworth–Emmons reactions<sup>16a</sup> or via direct sulfonylation of olefins.<sup>16b</sup>














The specific mechanistic details of our proposed photoredox vinylation reaction are outlined in Scheme 1. It has been established that irradiation of Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub> [dF(CF<sub>3</sub>)ppy = 2-(2,4-difluorophenyl)-5-trifluoromethylpyridine, dtbbpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine] (**1**, depicted as Ir<sup>III</sup> in Scheme 1) with visible light will generate the excited state Ir<sup>III</sup> species **2**, which is a strong oxidant ( $E_{1/2}$  [<sup>\*</sup>Ir<sup>III/II</sup>] = +1.21 V vs. SCE in MeCN).<sup>17</sup> This photoexcited complex should readily undergo single-electron transfer (SET) with a tertiary amine **3** (e.g. *N*-phenylpyrrolidine (R = Ph),  $E_{1/2}^{\text{red}}$  = +0.70 V vs. SCE)<sup>18</sup> to form a radical cation that upon deprotonation will deliver the  $\alpha$ -amino radical **5** (for R = Ph). Similarly, the excited state Ir<sup>III</sup> species **2** could also undergo SET with the carboxylate formed by deprotonation of  $\alpha$ -amino acid **4** (e.g. Boc-Pro-OCs,  $E_{1/2}^{\text{red}}$  = +0.95 V vs. SCE)<sup>7a</sup> to generate a carboxyl radical, which upon loss of CO<sub>2</sub> would deliver  $\alpha$ -amino radical **5** (for R = Boc). Reaction of **5** with a vinyl sulfone would then generate the  $\beta$ -sulfonyl radical **6**, which after elimination of a sulfinyl radical would provide the allylic amine product. Completion of the

photocatalytic cycle would then be accomplished via reduction of the sulfinyl radical (For PhSO<sub>2</sub>/PhSO<sub>2</sub>Na,  $E_{1/2}^{\text{red}}$  = +0.50 V vs. SCE)<sup>19</sup> with Ir<sup>II</sup> **7** ( $E_{1/2}$  [Ir<sup>III/II</sup>] = -1.37 V vs. SCE)<sup>17</sup> to give a sulfinate anion while reconstituting the photocatalyst **1**.

With this mechanistic hypothesis in mind, we set out to explore the feasibility of our proposed vinylation reaction (Table 1). Our initial investigations began by exposure of *N*-phenylpyrrolidine (**8**) to (*E*)-(2-(phenylsulfonyl)vinyl)benzene (**9**), in the presence of Ir(ppy)<sub>3</sub>, NaOAc, and a 26 W household fluorescent light bulb. To our delight, the desired allylic amine product **10** was observed under these new photocatalytic conditions, albeit in modest yield and stereoselectivity (entry 1, 28%, 21:79 *E:Z*). The efficiency of this vinylation protocol was improved by changing to less polar solvents, with toluene providing higher yield and selectivity compared to *N,N*-dimethylacetamide (DMA) (entry 2, 57% yield, 91:9 *E:Z*). Moreover, evaluation of a number of different photocatalysts revealed that less reducing photocatalysts, such as Ir(ppy)<sub>2</sub>(dtbbpy)PF<sub>6</sub> (**11**) and Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub> (**1**), provided enhanced *E*-selectivities (entries 3 and 4,  $\geq 95:5$  *E:Z*).<sup>20,21</sup> The choice of base was also found to have a dramatic effect on the yield while maintaining the high *E*-selectivity, with CsOAc proving to be optimal (entry 5, 81% yield). Finally, the use of 1,2-dichloroethane (DCE) as the reaction medium was found to provide slightly higher yields than toluene, delivering allyl amine **10** in 91% yield and with a 98:2 *E:Z* ratio (entry 7). The required participation of both the photocatalyst and light were confirmed via control experiments (entries 8 and 9).

Having identified optimal reaction conditions, we next examined the scope of the C–H vinylation reaction with respect to the amine substrate. As shown in Table 2, this protocol is successful with a range of tertiary *N*-aryl amines (products **10** and **12–24**, 64–98% yield, 94:6 to >98:2 *E:Z*). A variety of cyclic amines are well tolerated, with five-, six-, and seven-membered rings giving their respective products in excellent yields (products **10**, **12**, and **13**, 84–98% yield,  $\geq 97:3$  *E:Z*). This vinylation protocol is relatively unaffected by the nature of the

**Table 2.** Direct C–H Vinylation: Amine Scope.<sup>a</sup>


product	yield	<i>E:Z</i>	product	yield	<i>E:Z</i>
( $\pm$ )- <b>10</b> 	84%	>98:2	( $\pm$ )- <b>19</b> 	79%	94:6
( $\pm$ )- <b>12</b> 	84%	97:3	( $\pm$ )- <b>20</b> 	74%	94:6
( $\pm$ )- <b>13</b> 	98%	>98:2	( $\pm$ )- <b>21</b> 	73%	>98:2
( $\pm$ )- <b>14</b> 	67%	>98:2	( $\pm$ )- <b>22</b> 	80%	>98:2
( $\pm$ )- <b>15</b> 	75%	>98:2	( $\pm$ )- <b>23</b> 	64%	>98:2
( $\pm$ )- <b>16</b> 	76%	>98:2	( $\pm$ )- <b>24</b> 	84%	>98:2
( $\pm$ )- <b>17</b> 	75%	>98:2			
( $\pm$ )- <b>18</b> 	65%	>98:2			

<sup>a</sup>All reactions performed with 0.50 mmol vinyl sulfone, 2.5 equiv. amine, 3.0 equiv. CsOAc, and 1.0 mol% photocatalyst **1** in DCE (5.0 mL). Yields are of isolated products. *E:Z* ratio determined by <sup>1</sup>H NMR analysis.

*N*-aryl group, with substrates possessing electron donating or withdrawing functionalities demonstrating good efficiency (products **14–18**). Heteroatom-containing cyclic amines were also found to be suitable (products **19** and **20**) along with acyclic substrates (products **21** and **22**). Vinylation of a number of pharmacologically important fused nitrogen heterocycles such as *N*-benzyl-protected indolines and tetrahydroquinolines was demonstrated to be possible, with excellent regioselectivity for the  $\alpha$ -methylene ring position over the benzylic  $\alpha$ -amino site (products **23** and **24**).

We next sought to establish the scope of the vinyl sulfone in this direct C–H olefination reaction. As shown in Table 3,  $\beta$ -sulfonyl styrenes were found to be excellent substrates regardless of the electronic nature of the aryl substituent (products **25–31**, 74–85% yield, 92:8 to >98:2 *E:Z*). Moreover, vinyl sulfones that incorporate heteroaromatic substituents were found to be viable, including electron deficient pyridines and electron rich indoles (products **32–34**, 68–83% yield, 93:7 to >98:2 *E:Z*). The reaction is not limited to styrenyl reagents, as exemplified by the use of a conjugated dienylyl sulfone to generate an  $\alpha$ -amino diene product (**35**, 60% yield). Trisubstituted alkene adducts could also be delivered via the use of  $\beta,\beta$ -disubstituted vinyl sulfones that incorporate alkyl–aryl and aryl–aryl groups (products **36** and **37**, 84% yield). Perhaps most notable, unsaturated sulfones bearing a  $\beta$ -heteroatom were also successful in this protocol, allowing direct access to enamine functionality (product **38**, 79% yield). Unfortunately, extension of this methodology to the synthesis of tetrasubstituted  $\alpha$ -amino olefins could not be realized as  $\alpha$ -substituted vinyl sulfones have proven to be non-reactive under our optimized conditions.

To further extend the utility of this photoredox-mediated vinylation reaction, we next turned our attention to the decarboxylative olefination of *N*-carbamoyl  $\alpha$ -amino acids. Importantly, the use of readily cleavable carbamate protecting groups provides a valuable synthetic handle for further elaboration. To our delight, we found that Ir(ppy)<sub>2</sub>(dtbbpy)PF<sub>6</sub> (**11**), in the presence of 26 W light and CsHCO<sub>3</sub>, rapidly promoted the decarboxylative coupling of Boc-Pro-OH with sulfone **9** to generate the desired allylic amine **39** in 76% yield and with excellent olefin geometry control (Table 4, 94:6 *E:Z*).<sup>22</sup> Intriguingly, we discovered that while this CO<sub>2</sub>-extrusion, vinylation mechanism is efficiently catalyzed by photocatalyst **1**, the use of photocatalyst **11** was necessary to achieve high levels of *E*-olefin selectivity.<sup>23</sup> As demonstrated in Table 4, these optimized conditions could be applied to a wide range of  $\alpha$ -amino acids, to generate allylic amines in high yield and olefin geometry control (products **39–45**, 68–77% yield, 92:8 to >98:2 *E:Z*). Modification of the  $\alpha$ -amino acid protecting group is well tolerated, allowing incorporation of different carbamate systems

**Table 3.** Direct C–H Vinylation: Vinyl Sulfone Scope.<sup>a</sup>

amine	vinyl sulfone	product	yield	<i>E:Z</i>	allylic amine	product	yield	<i>E:Z</i>			
			(±)- <b>25</b>	R = <i>p</i> -Me	80%	>98:2		(±)- <b>33</b>	R = <i>p</i> -Me	68%	82:18 <sup>b</sup> (>98:2)
			(±)- <b>26</b>	R = <i>p</i> -Cl	80%	>98:2		(±)- <b>34</b>	R = <i>p</i> -OMe	83%	93:7
			(±)- <b>27</b>	R = <i>p</i> -CO <sub>2</sub> Me	74%	93:7		(±)- <b>35</b>	R = <i>o</i> -F	82%	>98:2
			(±)- <b>28</b>	R = <i>p</i> -OMe	78%	97:3		(±)- <b>36</b>	R = <i>m</i> -CF <sub>3</sub>	85%	>98:2
			(±)- <b>29</b>	R = <i>o</i> -F	82%	>98:2		(±)- <b>37</b>	R = <i>m</i> -CF <sub>3</sub>	85%	>98:2
			(±)- <b>30</b>	R = <i>m</i> -CF <sub>3</sub>	85%	>98:2		(±)- <b>38</b>	R = <i>m</i> -CF <sub>3</sub>	85%	>98:2
			(±)- <b>31</b>	R = <i>m</i> -CF <sub>3</sub>	77%	92:8		(±)- <b>39</b>	R = Me	84%	>98:2
			(±)- <b>32</b>	R = Me	78%	>98:2		(±)- <b>40</b>	R = Ph	84%	-
			(±)- <b>33</b>	R = Ph	84%	-		(±)- <b>41</b>	R = NPhth	79%	15:85

<sup>a</sup>All reactions performed with 0.50 mmol vinyl sulfone, 2.5 equiv. amine, 3.0 equiv. CsOAc, and 1.0 mol% photocatalyst **1** in DCE (5.0 mL). Yields are of isolated products. *E:Z* ratio determined by <sup>1</sup>H NMR analysis. <sup>b</sup>Isomerization from 82:18 *E:Z* (crude ratio) to >98:2 *E:Z* occurred during the mildly acidic purification conditions. NPhth = Phthalimidoyl.


(products **39** and **40**, 76 and 75% yield, respectively). Unnatural cyclic amino acids are readily employed to create the corresponding  $\alpha$ -amino olefins (product **42**, 74% yield, >98:2 *E:Z*). Importantly, this decarboxylative vinylation may also be applied to a range of acyclic  $\alpha$ -amino acids that incorporate a variety of functional groups including aliphatic, aromatic, sulfide, and indolic functionalities (products **41** and **43–45**, 68–77% yield, >98:2 *E:Z*).

We next examined the scope of the vinyl sulfone coupling partner in this decarboxylative olefination (Table 5). It is clear that a range of vinyl sulfones can be successfully employed in this CO<sub>2</sub>-extrusion mechanism without loss in yield or olefin geometry control (entries 1–4, 69–84% yield, 94:6–97:3 *E:Z*). Again, unsaturated sulfones bearing electron-withdrawing functional groups react efficiently (entries 1 and 2, 79 and 84% yield, 94:6 *E:Z*), as do those possessing electron-donating groups (entry 3, 69% yield, 97:3 *E:Z*). Finally, heteroaryl rings are also

**Table 4.** Decarboxylative Vinylation: Amino Acid Scope.<sup>a</sup>

amino acid	product, yield ( <i>E:Z</i> )	amino acid	product, yield ( <i>E:Z</i> )	amino acid	product, yield ( <i>E:Z</i> )
Boc-Pro-OH (R = Boc)	(±)- <b>39</b> 76% (94:6)	Boc-Pip-OH	(±)- <b>42</b> 74% (>98:2)	Boc-Met-OH	(±)- <b>44</b> 70% (>98:2)
Cbz-Pro-OH (R = Cbz)	(±)- <b>40</b> 75% (92:8)				
	(±)- <b>41</b> 77% (>98:2)	Boc-Val-OH	(±)- <b>43</b> 73% (>98:2)	Boc-Trp-OH	(±)- <b>45</b> 68% (>98:2)


<sup>a</sup>All reactions performed with 0.50 mmol vinyl sulfone, 1.2 equiv. amino acid, 2.0 equiv. CsHCO<sub>3</sub>, and 0.5 mol% photocatalyst **11** at 50 °C in 1,4-dioxane (30 mL). Yields are of isolated products. *E:Z* ratio determined by <sup>1</sup>H NMR analysis.

**Table 5.** Decarboxylative Vinylation: Vinyl Sulfone Scope.<sup>a</sup>

<sup>a</sup>All reactions performed using the conditions described in Table 4. Yields are of isolated products. *E:Z* ratio determined by <sup>1</sup>H NMR analysis.

readily tolerated in this coupling protocol, as demonstrated by the synthesis of drug-like pyridinyl-pyrrolidine adduct **49** (entry 4, 69% yield, 95:5 *E:Z*).

With the scope of the C–H and decarboxylative vinylation reactions shown to be broad across a range of substrates, we next turned our attention to synthetic applications of this methodology. As shown in Figure 1, a variety of pharmacophores were readily accessible via an intramolecular Heck reaction (**50**), an acid-promoted cyclization (**51**), and an alkene reduction (**52**). Furthermore, the decarboxylative and C–H vinylation reactions were applied to enable the concise total syntheses of the natural products (±)-norruspoline (**53**)<sup>24</sup> and (±)-galipinine (**54**),<sup>25</sup> respectively (see Supporting Information for details).

**Figure 1.** Derivatization of Allylic Amine Products.

**Acknowledgement.** Financial support was provided by NIHGMS (R01 GM103558-03) and kind gifts from Merck and Amgen.

**Supporting Information Available.** Experimental procedures and spectral data are provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) For recent reviews, see (a) Yoon, T. P.; Ischay, M. A.; Du, J. *Nat. Chem.* **2010**, *2*, 527. (b) Narayanam, J. M. R.; Stephenson, C. R. J. *Chem. Soc. Rev.* **2011**, *40*, 102. (c) Tucker, J. W.; Stephenson, C. R. J. *J. Org. Chem.* **2012**, *77*, 1617. (d) Xuan, J.; Xiao, W.-J. *Angew. Chem. Int. Ed.* **2012**, *51*, 6828. (e) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. *Chem. Rev.* **2013**, *113*, 5322. (f) Xie, J.; Jin, H.; Xu, P.; Zhu, C. *Tetrahedron Lett.* **2014**, *55*, 36. (g) Schultz, D. M.; Yoon, T. P. *Science* **2014**, *343*, 1239176.

- (2) (a) Pirmot, M. T.; Rankic, D. A.; Martin, D. B. C.; MacMillan, D. W. C. *Science* **2013**, *339*, 1593. (b) Petronijević, F. R.; Nappi, M.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2013**, *135*, 18323. (c) Qvortrup, K.; Rankic, D. A.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 626. (d) Terrett, J. A.; Clift, M. D.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 6858. (e) Xia, J.-B.; Zhu, C.; Chen, C. *J. Am. Chem. Soc.* **2013**, *135*, 17494.
- (3) (a) Bergman, R. G. *Nature* **2007**, *446*, 391. (b) Godula, K.; Sames, D. *Science* **2006**, *312*, 67. (c) Labinger, J. A.; Bercew, J. E. *Nature* **2002**, *417*, 507. (d) Campos, K. R. *Chem. Soc. Rev.* **2007**, *36*, 1069.
- (4) McNally, A.; Prier, C. K.; MacMillan, D. W. C. *Science* **2011**, *334*, 1114.
- (5) (a) Kohls, P.; Jadhav, D.; Pandey, G.; Reiser, O. *Org. Lett.* **2012**, *14*, 672. (b) Miyake, Y.; Nakajima, K.; Nishibayashi, Y. *J. Am. Chem. Soc.* **2012**, *134*, 3338. (c) Miyake, Y.; Ashida, Y.; Nakajima, K.; Nishibayashi, Y. *Chem. Commun.* **2012**, *48*, 6966. (d) Zhu, S.; Das, A.; Bui, L.; Curran, D. P.; Rueping, M. *J. Am. Chem. Soc.* **2013**, *135*, 1823. (e) Espelt, L. R.; Wiensch, E. M.; Yoon, T. P. *J. Org. Chem.* **2013**, *78*, 4107. (f) Mizoguchi, H.; Oikawa, H.; Oguri, H. *Nat. Chem.* **2014**, *6*, 57. (g) Zhou, H.; Lu, P.; Gu, X.; Li, P. *Org. Lett.* **2013**, *15*, 5646. (h) Miyake, Y.; Nakajima, K.; Nishibayashi, Y. *Chem. Eur. J.* **2012**, *18*, 16473. (i) Singh, A.; Arora, A.; Weaver, J. D. *Org. Lett.* **2013**, *15*, 5390.
- (6) For reviews on functionalization of  $\alpha$ -amino C–H bonds using photoredox catalysis, see: (a) Shi, L.; Xia, W. *Chem. Soc. Rev.* **2012**, *41*, 7687. (b) Hu, J.; Wang, J.; Nguyen, T. H.; Zheng, N. *Beilstein J. Org. Chem.* **2013**, *9*, 1977.
- (7) (a) Zuo, Z.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 5257. (b) Zuo, Z.; Ahneman, D.; Chu, L.; Terrett, J.; Doyle, A. G.; MacMillan, D. W. C. *Science* **2014**, *10.1126/science.1255525*.
- (8) For a photoredox-mediated generation of  $\alpha$ -amino radicals from *N*-aryl  $\alpha$ -amino acids, see: Chen, L.; Chao, C. S.; Pan, Y.; Dong, S.; Teo, Y. C.; Wang, J.; Tan, C.-H. *Org. Biomol. Chem.* **2013**, *11*, 5922.
- (9) Nag, S.; Batra, S. *Tetrahedron* **2011**, *67*, 8959.
- (10) (a) Skoda, E. M.; Davis, G. C.; Wipf, P. *Org. Process Res. Dev.* **2012**, *16*, 26. (b) Stüttz, A. *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 320.
- (11) (a) Skucas, E.; Ngai, M.-Y.; Komanduri, V.; Krische, M. J. *Acc. Chem. Res.* **2007**, *40*, 1394. (b) Candeias, N. R.; Montalbano, F.; Cal, P. M. S. D.; Gois, P. M. P. *Chem. Rev.* **2010**, *110*, 6169.
- (12) (a) Cheikh, R. B.; Chaabouni, R.; Laurent, A.; Mison, P.; Nafti, A. *Synthesis* **1983**, 685. (b) Johannsen, M.; Jørgensen, K. A. *Chem. Rev.* **1998**, *98*, 1689. (c) Overman, L. E.; Carpenter, N. E. *Org. React.* **2005**, *66*, 1. (c) Collet, F.; Lescot, C.; Dauban, P. *Chem. Soc. Rev.* **2011**, *40*, 1926. (d) Ramirez, T. A.; Zhao, B.; Shi, Y. *Chem. Soc. Rev.* **2012**, *41*, 931.
- (13) For selected examples, see: (a) Dieter, R. K.; Li, S. J. *J. Org. Chem.* **1997**, *62*, 7726. (b) Dieter, R. K.; Oba, G.; Chandupatla, K. R.; Topping, C. M.; Lu, K.; Watson, R. T. *J. Org. Chem.* **2004**, *69*, 3076. (c) Beng, T. K.; Gawley, R. E. *Org. Lett.* **2011**, *13*, 394. (h) Barker, G.; McGrath, J. L.; Klapars, A.; Stead, D.; Zhou, G.; Campos, K. R.; O'Brien, P. *J. Org. Chem.* **2011**, *76*, 5936. (i) Millet, A.; Dailier, D.; Larini, P.; Baudoin, O. *Angew. Chem. Int. Ed.* **2014**, *53*, 2678.
- (14) For selected examples of the use of vinyl sulfones as radical vinylating agents, see: (a) Miyamoto, N.; Fukuoka, D.; Utimoto, K.; Nozaki, H. *Bull. Chem. Soc. Jpn.* **1974**, *47*, 503. (b) Russell, G. A.; Ngoviwatchai, P.; Tashtoush, H.; Paladmau, A.; Khanna, R. K. *J. Am. Chem. Soc.* **1988**, *110*, 3530. (c) Xiang, J.; Jiang, W.; Gong, J.; Fuchs, P. L. *J. Am. Chem. Soc.* **1997**, *119*, 4123. (d) Bertrand, F.; Quiclet-Sire, B.; Zard, S. Z. *Angew. Chem. Int. Ed.* **1999**, *38*, 1943. (e) Schaffner, A.-P.; Darmency, V.; Renaud, P. *Angew. Chem. Int. Ed.* **2006**, *45*, 5847. (f) Poittevin, C.; Liautard, V.; Beniazza, R.; Robert, F.; Landais, Y. *Org. Lett.* **2013**, *15*, 2814.
- (15) After submission of this article, a related photochemical vinylation was reported: Amaoka, Y.; Nagatomo, M.; Watanabe, M.; Tao, K.; Kamijo, S.; Inoue, M. *Chem. Sci.* **2014**, doi: 10.1039/c4sc01631a.
- (16) (a) Lee, W. J.; Oh, D. Y. *Synth. Commun.* **1989**, *19*, 2209. (b) Sawangphon, T.; Katrun, P.; Chaisiwamongkhon, K.; Pohmakotr, M.; Reutrakul, V.; Jaipetch, T.; Soorukram, D.; Kuhakarn, C. *Synth. Commun.* **2013**, *43*, 1692.
- (17) Lowry, M. S.; Goldsmith, J. I.; Slinker, J. D.; Rohl, R.; Pascal Jr., R. A.; Malliaras, G. G.; Bernhard, S. *Chem. Mater.* **2005**, *17*, 5712.
- (18) Liu, W.; Ma, Y.; Yin, Y.; Zhao, Y. *Bull. Chem. Soc. Jpn.* **2004**, *79*, 577.
- (19) Persson, B. *Acta Chem. Scand.* **1977**, *31B*, 88.
- (20) The lower *E*-selectivities with Ir(ppy)<sub>3</sub> were found to be a result of post reaction isomerization. Post-reaction isomerization was also promoted by photocatalysts **1** and **11**, although this only became apparent after prolonged reaction times. For a related isomerization process, see: Singh, K.; Staig, S. J.; Weaver, J. D. *J. Am. Chem. Soc.* **2014**, *136*, 5275.
- (21) For discussions regarding the stereoselectivity of radical additions to vinyl sulfones see references 14b and 14c.
- (22) See Supporting Information for optimization studies.
- (23) Using photocatalyst **1** instead of **11** gave comparable yields of **39** but with an *E:Z* ratio of 1:1. Using photocatalyst **1** in combination with KOAc as the base led to improved *Z*-selectivity ( $\geq 3:1$  *Z:E*).
- (24) Roessler, F.; Ganzinger, D.; John, S.; Schöpp, E.; Hesse, M. *Helv. Chim. Acta* **1978**, *61*, 1200.
- (25) (a) Jacquemond-Collet, I.; Hannedouche, S.; Fabre, N.; Fouraste, I.; Moulis, C. *Phytochemistry* **1999**, *51*, 1167. (b) Jacquemond-Collet, I.; Benoit-Vical, F.; Mustofa; Valentin, A.; Stanislas, E.; Mallié, M.; Fourasté, I. *Planta Med.* **2002**, *68*, 68.

## Graphical Abstract

