



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Preparation of N-Arylsulfonyl Imines from Sulfonamides and Aryl Aldehydes Using Magnesium Oxide as a Heterogeneous and Reusable Catalyst Under Solvent-Free Conditions

Alireza Hasaninejad ^a, Abdolkarim Zare ^b, Ahmad Reza Moosavi Zare ^c, Abolfath Parhami ^c, Hashem Sharghi ^c & Ali Khalafi-Nezhad ^c

^a Department of Chemistry, Faculty of Sciences, Persian Gulf University, Bushehr, Iran

^b Department of Chemistry, Payame Nour University (PNU), Iran

^c Department of Chemistry, College of Sciences, Shiraz University, Shiraz, Iran

Published online: 07 Oct 2008.

To cite this article: Alireza Hasaninejad, Abdolkarim Zare, Ahmad Reza Moosavi Zare, Abolfath Parhami, Hashem Sharghi & Ali Khalafi-Nezhad (2008) Preparation of N-Arylsulfonyl Imines from Sulfonamides and Aryl Aldehydes Using Magnesium Oxide as a Heterogeneous and Reusable Catalyst Under Solvent-Free Conditions, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 183:11, 2769-2776

To link to this article: <http://dx.doi.org/10.1080/10426500802002802>

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Preparation of *N*-Arylsulfonyl Imines from Sulfonamides and Aryl Aldehydes Using Magnesium Oxide as a Heterogeneous and Reusable Catalyst Under Solvent-Free Conditions

Alireza Hasaninejad,¹ Abdolkarim Zare,² Ahmad Reza Moosavi Zare,³ Abolfath Parhami,³ Hashem Sharghi,³ and Ali Khalafi-Nezhad³

¹Department of Chemistry, Faculty of Sciences, Persian Gulf University, Bushehr, Iran

²Department of Chemistry, Payame Nour University (PNU), Iran

³Department of Chemistry, College of Sciences, Shiraz University, Shiraz, Iran

*An efficient solvent-free procedure for the preparation of *N*-arylsulfonyl imines from sulfonamides and aryl aldehydes in the presence of a catalytic amount of magnesium oxide and tetrabutylammonium bromide (TBAB) under microwave irradiation is described. The advantages of this method are good to high yields, short reaction times, low cost, and matching with green chemistry protocols.*

Keywords Aldehyde; magnesium oxide; microwave; *N*-arylsulfonyl imine; solvent-free; sulfonamide

INTRODUCTION

N-Sulfonyl imines are versatile intermediates in organic synthesis.¹ They are excellent substrates in aza Diels-Alder reactions,² nucleophilic additions³ and reductions,⁴ as well as in radical⁵ or ene reactions.⁶ They have been also used in the synthesis of aziridines.⁷ Several synthetic methods for the preparation of *N*-sulfonyl imines have been reported so far.^{8–21} Most of them involve the condensation of sulfonamides

Received 25 November 2007; accepted 19 February 2008.

We thank the Persian Gulf University, the Payame Nour University, and the Shiraz University Research Councils for financial support of this work.


Address correspondence to Alireza Hasaninejad, Department of Chemistry, Faculty of Sciences, Persian Gulf University, Bushehr 75169, Iran. E-mail: ahassaninejad@yahoo.com or Abdolkarim Zare, Department of Chemistry, Payame Nour University (PNU), Iran. E-mail: abdolkarimzare@yahoo.com

with aldehydes in the presence of strong Lewis acids. In many cases, strongly acidic conditions are not compatible with other functionalities present in a given substrate. Some methods need two-steps procedures or they are expensive methods and generate toxic byproducts. Moreover, some methods require the preparation of the starting materials such as oxime, sulfinylimine and/or aziridine.^{13,14,17} Because of the limitations of the above methods, development of a new method for the synthesis of *N*-sulfonyl imines under basic conditions in one step would be desirable.

Recently, mineral oxides have proved to be useful to chemists in the laboratory as well as in industry due to the effective activation of adsorbed compounds and the enhancement of reaction rate, selectivity, easier workup, recyclability of the supports and the eco-friendly reaction conditions.²² Magnesium oxide is one of these oxides: it is inexpensive, reusable, commercially available and an environmentally benign catalyst. Magnesium oxide has been extensively studied as a promising catalyst,²³ modifier, or promoter,²⁴ and catalyst support in several transformations such as Michael addition and Knoevenagel condensation,²⁵ transformation of ethanol to butanol²⁶ and self-condensation of propanol.²⁷

Solvent-free organic reactions have been applied as a useful protocol in organic synthesis. Solvent-free reactions under thermal or microwave conditions often lead to shorter reaction times, increased yields, and easier workup. They match with green chemistry protocols and may enhance the regio- and stereoselectivity of reactions.²⁸

Considering the above aspects, and along with our previous work on the synthesis of *N*-sulfonyl imines published in this journal,⁹ as well as in extension of our previous studies on solvent-free organic syntheses,^{22a,29} we report here a clean, facile and rapid method for the preparation of *N*-arylsulfonyl imines from sulfonamides and aryl aldehydes in the presence of a catalytic amount of magnesium oxide and tetrabutylammonium bromide (TBAB) under microwave irradiation (Scheme 1).



SCHEME 1

TABLE I The Influence of Various Basic Catalysts on the Reaction of Benzenesulfonamide with Benzaldehyde in the Presence of TBAB Under Microwave Irradiation (300 W, max. 110°C)

Catalyst	Time (min)	Yield ^a (%)
MgO	8	87
CaO	8	74
Cs ₂ CO ₃	8	61
K ₂ CO ₃	8	68
<i>t</i> -BuOK	8	53
DABCO	12	65
NBu ₃	12	57
1-Methyl-imidazole	12	60

^aIsolated yield.

RESULTS AND DISCUSSION

To determine the optimal reaction conditions we have studied the reaction of benzenesulfonamide with benzaldehyde as a model reaction to provide compound **1** (Scheme 1 and Table I). At first, the effect of various basic catalysts was examined to evaluate their capabilities. The results are summarized in Table I. As shown in Table I higher yields and shorter reaction times were achieved when MgO was used. Therefore, MgO was chosen as catalyst for all the other reactions.

In order to select the appropriate microwave power the model reaction was examined at different microwave powers (100–600 W) with controlled temperature (max. 110°C) in the presence of MgO and TBAB. The best results were observed at 300 W of microwave power.

In another study, the role of TBAB was evaluated. The absence of TBAB in the reaction medium afforded low yields even when the reaction time or the microwave power was increased. Thus, the presence of TBAB in the reaction mixture is critically significant. TBAB absorbs the microwave irradiation, generates in situ heat and increases the temperature above its melting point (100–103°C). In addition TBAB creates a homogeneous medium similar to that of an ionic liquid.^{28a}

To investigate the versatility, as well as the capacity of our method, the reaction of sulfonamides with various structurally diverse aromatic aldehydes were examined (Table II). As indicated in Table II, the reactions proceeded efficiently and the *N*-arylsulfonyl imines were obtained in good to high yields in short reaction times.

It has been observed that the electronic properties of the aromatic ring of aldehydes affected the reaction. The results showed that halogen

TABLE II Preparation of *N*-Arylsulfonyl Imines from Sulfonamides and Aldehydes in the Presence of Catalytic Amounts of MgO and TBAB Under Microwave Irradiation (300 W, max. 110°C)





































Aldehyde	Product	M. p. (Lit.)	Time (min)	Yield ^a (%)
		77–80 (78–81) ^[18]	8	87
		123–124 (125–126) ^[8]	15	64
		205–207 (208–209) ^[19]	12	85
		206–207 (209–210) ^[11]	15	79
		130–132 (131–132) ^[20]	12	84
		113–114 (111) ^[11]	15	62
		164–165 (162–164) ^[21]	15	56
		108–109 (109) ^[16]	12	80
		126–128 (126–128) ^[9]	20	60
		172–173 (172–173) ^[8]	15	77
		181–183 (182–185) ^[18]	15	78
		88–90 (91–92) ^[16]	20	54
		112–114 (111–112) ^[8]	15	76
		118–120 (121–123) ^[18]	15	75

TABLE II Preparation of *N*-Arylsulfonyl Imines from Sulfonamides and Aldehydes in the Presence of Catalytic Amounts of MgO and TBAB Under Microwave Irradiation (300 W, max. 110°C) (*Continued*)

Aldehyde	Product	M. p. (Lit.)	Time (min)	Yield ^a (%)
		111–114 (114–117) ^[18]	17	71
		126–128 (128–129) ^[8]	15	73
		102–104 (101–102) ^[8]	20	57
		172–173 (172–173) ^[9]	20	59

^aIsolated yield.

atoms as well as electron-donating substituents had no significant effect on the yields (Table II). However, aryl aldehydes possessing electron-withdrawing or sterically demanding substituents generally need longer reaction times, and the yields are lower (Table II). It should be mentioned that the yield of *N*-arylsulfonyl imines resulting from the reaction of benzenesulfonamide with aldehydes was higher than that starting from 4-methyl benzenesulfonamide.

The interesting behavior of magnesium oxide lies in the fact that it can be reused after simple washing with ethyl acetate, thus rendering the process more economic. The yields of compound **1** (model compound) when MgO was used for the 2nd, 3rd, 4th, and 5th time were almost as high as when it was used for the first time.

In conclusion, the procedure presented here provides a clean, rapid and efficient solvent-free route to *N*-arylsulfonyl imines *via* the reaction of sulfonamides with aryl aldehydes using cheap and safe reagents.

EXPERIMENTAL

All chemicals were obtained from Merck or Fluka. The compounds were identified by the comparison of their melting points and/or ¹H and ¹³C NMR data with those of authentic samples. All reactions were carried out using a CEM MARS 5TM microwave oven. The ¹H (250 MHz) and ¹³C (62.5 MHz) NMR spectra were obtained with a Bruker Avance DPX-250

spectrometer. Melting points were recorded with a Büchi B-545 apparatus in open capillary tubes and are uncorrected.

General Procedure for the Preparation of *N*-Arylsulfonyl Imines from Sulfonamides and Aryl Aldehydes Under Solvent-Free Conditions

To a well ground mixture of the sulfonamide (2 mmol), MgO (0.020 g, 0.5 mmol) and TBAB (0.129 g, 0.4 mmol) in a microwave vessel was added the respective aldehyde (2.6 mmol),³⁰ and the compounds were carefully mixed with a tiny spatula. The mixture was irradiated in the microwave oven at 300 W for the times reported in Table II. The microwave oven was programmed to give a maximum internal temperature to 110°C. The reaction mixture was cooled to room temperature, suspended in ethyl acetate (50 mL), filtered, and the filtrate was washed with water (2 × 50 mL) and dried over MgSO₄. The filtrate was concentrated to 5 mL, *n*-hexane (20–30 mL) was added, and the solution was allowed to stand at room temperature for 5–6 h. During this time crystals formed, which were collected by filtration, washed with *n*-hexane (20 mL) and dried.

REFERENCES

- [1] (a) S. M. Weinreb, *Top. Curr. Chem.*, **190**, 131 (1997); (b) M. Gohain, *Synlett*, 2097 (2003); (c) R. Bloch, *Chem. Rev.*, **98**, 1407 (1998).
- [2] (a) J. Sisko and S. M. Weinreb, *Tetrahedron Lett.*, **30**, 3037 (1989); (b) D. L. Boger, W. L. Corbett, T. T. Curran, and A. M. Kasper, *J. Am. Chem. Soc.*, **113**, 1713 (1991); (c) O. Garcia-Mancheno, R. Gomez-Arrayas, and J. C. Carretero, *J. Am. Chem. Soc.*, **126**, 456 (2004) and references cited therein.
- [3] (a) M. J. Melnick, A. J. Freyes, and S. M. Weinreb, *Tetrahedron Lett.*, **29**, 3891 (1988); (b) K. Yamada, H. Fujihara, Y. Yamamoto, Y. Miwa, T. Taba, and K. Tomioka, *Org. Lett.*, 3509 (2002); (c) V. Aggarwal, E. Alonso, M. Ferrar, and S. E. Spey, *J. Org. Chem.*, **67**, 2335 (2002); (d) T. Soeta, K. Nagai, H. Fujihara, M. Kuriyama, and K. Tomioka, *J. Org. Chem.*, **68**, 9723 (2003); (e) P. Wipf, C. Kendall, and C. R. Stephenson, *J. Am. Chem. Soc.*, **125**, 761 (2003); (f) M. Gunter and H.-J. Gais, *J. Org. Chem.*, **68**, 8037 (2003). (g) H. -K. Yim and H. N. C. Wong, *J. Org. Chem.*, **69**, 2892 (2004).
- [4] (a) M. Hojo, C. Murakami, A. Fujii, and A. Hosomi, *Tetrahedron Lett.*, **40**, 911 (1999); (b) H. Nisikori, R. Yoshihara, and A. Hosomi, *Synlett*, 561 (2003).
- [5] K. -I. Yamada, H. Fujihara, Y. Yamamoto, Y. Miwa, T. Taga, and K. Tomioka, *Org. Lett.*, 3509 (2002).
- [6] M. Yamanaka, A. Nishida, and M. Nakagwa, *Org. Lett.*, 159 (2002).
- [7] (a) R. Hori, T. Aoayama, and T. Shoiri, *Tetrahedron Lett.*, **41**, 9455 (2000); (b) L. G. Arini, A. Sinclair, P. Szeto, and R. A. Stockan, *Tetrahedron Lett.*, **45**, 1589 (2004).
- [8] B. M. Trost and M. Christopher, *J. Org. Chem.*, **56**, 6468 (1991).
- [9] A. Hasaninejad and H. Sharghi, *Phosphorus, Sulfur, and Silicon*, **182**, 873 (2007).

- [10] K. Y. Lee, C. G. Lee, and J. N. Kim, *Tetrahedron Lett.*, **44**, 1231 (2003) and references cited therein.
- [11] E. Ziegler and W. Ruef, *Z. Naturforsch.*, **30B**, 946 (1975).
- [12] G. D. Artman, A. Bartolozzi, R. W. Franck, and S. M. Weinreb, *Synlett*, 232 (2001).
- [13] D. L. Boger and W. L. Corbett, *J. Org. Chem.*, **57**, 4777 (1992).
- [14] J. Wolfe and J. E. Ney, *Org. Lett.*, **5**, 4607 (2003).
- [15] (a) W. B. Jennings and C. J. Lovely, *Tetrahedron*, **47**, 5561 (1991); (b) G. I. Georg, G. C. B. Harriman, and S. A. Peterson, *J. Org. Chem.*, **60**, 7366 (1995); (c) R. N. Ram and A. A. Khan, *Synth. Commun.*, **31**, 841 (2001); (d) T. Jin, G. Feng, M. Yang, and T. Li, *Synth. Commun.*, **34**, 1277 (2004).
- [16] F. A. Davis, J. Lamendola, U. Nadir, E. W. Kluger, T. C. Sedergran, T. W. Panunto, R. Billmers, R. Jenkins, I. J. Turchi, W. H. Watson, J. S. Chen, and M. Kimura, *J. Am. Chem. Soc.*, **102**, 2000 (1980).
- [17] J. L. G. Ruano, J. Aleman, M. Belen Cid, and A. Parra, *Org. Lett.*, 179 (2005).
- [18] A. Vass, J. Dudas, and R. S. Varma, *Tetrahedron Lett.*, **40**, 4951 (1999).
- [19] V. I. Naddaka, K. V. Avanesyan, M. L. Cherkinskaya, and V. I. Minkin, *Zh. Org. Khim.*, **24**, 603 (1988).
- [20] Z. I. Kuplennik, Z. N. Belaya, and A. M. Pinchuk, *Zh. Obshch. Khim.*, **51**, 2711 (1981).
- [21] F. A. Davis and S. G. La, *J. Org. Chem.*, **53**, 5004 (1988).
- [22] (a) A. Zare, A. Hasaninejad, A. Khalafi-Nezhad, A. R. Moosavi Zare, A. Parhami, and G. R. Nejabat, *Arkivoc*, **1**, 58 (2007); (b) M. Hosseini Sarvari and H. Sharghi, *Tetrahedron*, **61**, 10903 (2005); (c) M. Hosseini Sarvari, *Synthesis*, 787 (2005); (d) M. Hosseini Sarvari and H. Sharghi, *J. Org. Chem.*, **69**, 6953 (2004); (e) Y. J. Kim and R. S. Varma, *Tetrahedron Lett.*, **45**, 7205 (2004); (f) R. S. Varma, *Tetrahedron*, **58**, 1235 (2002); (g) H. Sharghi and M. Hosseini, *Synthesis*, 1057 (2002); (h) M. Gupta, S. Paul, R. Gupta, and A. Loupy, *Tetrahedron Lett.*, **46**, 4957 (2005); (i) F. Tamaddon, M. A. Amrollahi, and L. Sharafat, *Tetrahedron Lett.*, **46**, 7841 (2005).
- [23] M. A. Aramendia, V. Borau, C. Jiménez, J. M. Marinas, J. R. Ruiz, and F. J. Urbano, *Appl. Catal. A*, 207 (2003).
- [24] P. Kovacheva, K. Arishtirova, and S. Vassilev, *Appl. Catal. A*, 3915 (2001).
- [25] C. Xu, J. K. Bartley, D. I. Enache, D. W. Knight, and G. J. Hutchings, *Synthesis*, 3468 (2005).
- [26] A. S. Ndou, N. Plint, and N. J. Coville, *Appl. Catal. A*, 337 (2003).
- [27] A. S. Ndou and N. J. Coville, *Appl. Catal. A*, 103 (2004).
- [28] (a) A. Loupy, *Microwave in Organic Synthesis*, Wiley-VCH, Weinheim, 2004; (b) K. Tanaka, *Solvent-free Organic Synthesis*, (Wiley-VCH, Weinheim, 2004); (c) R. S. Varma, *Advances in Green Chemistry: Chemical Synthesis Using Microwave Irradiation*, Astra Zeneca Research Foundation, Kavitha Printers: Bangalore, India, 2002; (d) A. Shockravi, H. Sharghi, H. Valizadeh, and M. M. Heravi, *Phosphorus, Sulfur, and Silicon*, **177**, 2555 (2002); (e) A. Ramazani, L. Yousefi, E. Ahmadi, and A. Souldozi, *Phosphorus, Sulfur, and Silicon*, **179**, 1459 (2004); (f) M. Zolfigol, F. Shirini, K. Zamani, E. Ghofrani, and S. Ebrahimi, *Phosphorus, Sulfur, and Silicon*, **179**, 2177 (2004); (g) X. Wang, J. Liu, and Z. Li, *Phosphorus, Sulfur and Silicon*, **181**, 627 (2006); (h) M. Zolfigol, P. Salehi, and M. Shiri, *Phosphorus, Sulfur and Silicon*, **179**, 2273 (2004); (i) A. Javidan and A. Banan, *Phosphorus, Sulfur, and Silicon*, **179**, 2253 (2004); (j) A. Shaabani, A. Bazgir, and S. Arab-Ameri, *Phosphorus, Sulfur, and Silicon*, **179**, 2169 (2004); (k) A. Khalafi-Nezhad, A. Zare, A. Parhami, M. N. Soltani Rad, and G. R. Nejabat, *Can. J. Chem.*, **84**, 979 (2006).
- [29] (a) A. Khalafi-Nezhad, A. Zare, A. Parhami, M. N. Soltani Rad, and G. R. Nejabat, *Phosphorus, Sulfur, and Silicon*, **182**, 657 (2007); (b) A. Khalafi-Nezhad, A. Zare,

- A. Parhami, M. N. Soltani Rad, and G. Reza Nejabat, *Synth. Commun.*, **36**, 3549 (2006); (c) A. Khalafi-Nezhad, A. Zare, M. N. Soltani Rad, A. Mokhtari, and A. Parhami, *Synthesis*, 419 (2005); (d) A. Khalafi-Nezhad, A. Parhami, M. N. Soltani Rad, and A. Zarea, *Tetrahedron Lett.*, **46**, 6879 (2005); (e) A. Khalafi-Nezhad, M. N. Soltani Rad, and A. Khoshnood, *Synthesis*, 583 (2004); (f) A. Khalafi-Nezhad, B. Mokhtari, and M. N. Soltani Rad, *Tetrahedron Lett.*, **44**, 7325 (2003); (g) A. Khalafi-Nezhad, M. N. Soltani Rad, and G. H. Hakimelahi, *Helv. Chim. Acta*, **86**, 2396 (2003); (h) A. Khalafi-Nezhad and A. Hashemi, *Iran. J. Chem. Chem. Eng.*, **20**, 9 (2001); (i) A. Khalafi-Nezhad, F. Alamdari, and N. Zekri, *Tetrahedron*, **56**, 7503 (2000); (j) A. Khalafi-Nezhad and A. Hashemi, *J. Chem. Res. (S)*, 720 (1999).
- [30] In the case of solid aldehydes, they were thoroughly ground with the other reactants to give a homogeneous mixture.