Rhodium-Catalyzed Ketone Methylation Using Methanol Under Mild Conditions: Formation of α-Branched Products**

Louis K. M. Chan, Darren L. Poole, Di Shen, Mark P. Healy, and Timothy J. Donohoe*

Abstract: The rhodium-catalyzed methylation of ketones has been accomplished using methanol as the methylating agent and the hydrogen-borrowing method. The sequence is notable for the relatively low temperatures that are required and for the ability of the reaction system to form α -branched products with ease. Doubly alkylated ketones can be prepared from methyl ketones and two different alcohols by using a sequential onepot iridium- and rhodium-catalyzed process.

ydrogen borrowing is a powerful method for functionalgroup interconversion, which involves reversible changes in the oxidation state of the reacting partners. In essence, a catalyst alters the reactivity of a compound by removing two hydrogen atoms in a formal oxidation of the substrate. This temporarily generates a highly reactive intermediate and permits bond formation to take place. Finally, the intermediate is reduced with the redelivery of two hydrogen atoms, giving a product without a net change in the overall oxidation state.^[1]

Since a discovery by Grigg et al., who used alcohols as alkylating reagents,^[2] the transition-metal-catalyzed α -alkylation of carbonyl compounds using hydrogen borrowing has emerged as a productive area of research. Contributions from the groups of Cho/Shim,^[3] Ishii,^[4] Nishibayashi,^[5] Ramón/ Yus,^[6] Park,^[7] Uozumi,^[8] Williams,^[9] and others^[10] have demonstrated variations of such alkylation reactions using different metal catalyst systems (Scheme 1 A).^[11] These methods have allowed ketone alkylation with benzyl and long-chain *n*-alkyl alcohols (but with surprisingly few, if any, examples using short-chain alcohols such as methanol). Moreover, the scope of the ketone substrates is predominantly limited to methyl ketones with linear products being obtained after monoalkylation, whereafter the reaction stops.



- [⁺] These authors contributed equally to this work.
- [**] We thank the EPSRC/Pharma Organic Synthetic Chemistry Studentships program and EPSRC, Pfizer, and Novartis for support.
- Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201307950.
- © 2014 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

A) Existing ketone alkylation methods via hydrogen borrowing



B) Precedent for the formation of branched products adjacent to a



C) This work: Application in methylation using methanol; form branched products?



Scheme 1. Different alkylation methods.

The formation of α -branched products from a more-substituted carbonyl starting material is much more challenging and restricted to a handful of examples (Scheme 1B).^[2g,3,4a,c,6b,7,9c]

Our research program seeks to overcome these problems, and we chose to investigate the possibility of performing a methylation at the α -position to a ketone using methanol as the alkylating reagent. We also wanted to address the formation of branched products (i.e., cause the formation of a stereogenic center) because a general procedure would represent an important and useful advance.

The use of methanol in catalytic alcohol dehydrogenation reactions is a worthwhile and challenging goal, with fundamental developments made by the groups of Beller^[12] and Milstein^[13] and demonstrated further with methanol by the groups of Tincado/Grützmacher^[14] and Glorius.^[15] However, methods that utilize methanol as an alkylating reagent with C–C bond formation (reported on PhCH₂CN^[2e,16]) or as a substrate for formal C–H functionalization (such as the work by Krische et al.^[17]) are extremely rare and represent an important area in need of development.

Herein, we report our findings on the synthesis of α branched ketone products (Scheme 1 C). They are significant because 1) they provide a general solution to the problem of ketone methylation using methanol; 2) the conditions are mild and the catalyst commercially available; 3) the formation of α -branched products is now possible.

Angew. Chem. Int. Ed. 2014, 53, 761-765

© 2014 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Our initial reactions employed valerophenone in methanol, with various transition metal catalysts and Cs_2CO_3 as a base; early on we discovered the beneficial effect of performing reactions under an atmosphere of oxygen, which is unusual in the context of hydrogen-borrowing chemistry. Interestingly, we also found that the alkylation of valerophenone by methanol was observed at temperatures as low as $65 \,^{\circ}C$,^[18] implying that methanol dehydrogenation was taking place under these relatively mild conditions.

Early experiments using catalytic amounts of $[{Cp*IrCl_2}_2]$ or $[{Cp*RuCl_2}_n]$ gave a mixture of the desired α -methylated ketone **2a**, dimer **3a**, enone **4a**, and β -methoxyketone **5a** (Table 1, entries 1,2). The product distribution shows that,

Table 1: Optimization of the α -methylation of ketone **1 a** (0.2 M).^[a]



Entry	Conditions	Yield [%]			
		(±)-2a	3 a -	4a	(±)-5 a
1	5 mol% [{Cp*IrCl ₂ } ₂]	42	12	14	27
2	10 mol% [Cp*RuCl ₂]	< 2	19	23	39
3	5 mol % [{Cp*RhCl ₂ } ₂]	98	n/o	n/o	n/o
4 ^[b]	5 mol% [{Cp*RhCl ₂ } ₂]	53	33	n/o	n/o
5	10 mol % RhCl ₃ ·H ₂ O	14	10	18	30
6	10 mol% RhCl₃•H₂O, 10 mol% Cp*H	74	8	n/o	n/o
7	10 mol% RhCl₃•H₂O, 10 mol% PPh₃	15	12	27	44
8	10 mol% [RhCl(PPh ₃) ₃]	23	n/o	n/o	n/o
9	5 mol% [Rh₂(OAc)₄]	10	21	5	23
10	no catalyst	n/o (98) ^[c]	< 5	n/o	n/o
11	5 mol% [{Cp*RhCl ₂ } ₂], argon	57	9	n/o	n/o
12	10 mol% Cp*H	n/o (86) ^[c]	trace	n/o	n/o

[a] n/o=not observed. All yields given are of isolated material.	Compound 3a was
isolated as a 1:1 mixture of diastereomers. [b] 5 equiv of KOH i	nstead of Cs_2CO_3 .
[c] Yield of recovered starting material.	

along with consumption of the starting material 1a, the conjugate reduction process was interrupted by the addition of nucleophiles to the enone 4a. Thus, in an attempt to improve the yield of 2a, we turned to using Rh, a well-known catalyst for 1,4-reduction.^[19] When using 5 mol% of [{Cp*RhCl₂]₂], α -methylation could be accomplished in 98% yield (entry 3); the use of a stronger base such as KOH resulted in a higher amount of dimer 3a formed (entry 4); RhCl₃ gave a mixture of unreduced intermediates (entry 5), whereas a respectable 74% yield of 2a was achieved by using a mixture of RhCl₃ and added Cp* ligand (entry 6).^[20] We were pleased to find that the reaction does not require the use of anhydrous methanol or extra precautions such as the exclusion of moisture, making this a highly practical method when compared to the traditional lithiumenolate alkylation methods.

When testing other Rh complexes, we found that Rh^I, Rh^{II}, and Rh^{III} complexes such as RhCl₃/PPh₃, [RhCl(PPh₃)₃],

and $[Rh_2(OAc)_4]$ gave small amounts of the desired product (entries 7–9). As a control experiment, the reaction was performed in the absence of the Rh catalyst and no product (**2a**) or intermediates were observed (entry 10). In reactions that were run under argon (entry 11), the yield of **2a** was acceptable but inferior to that shown earlier (entry 3). A further reaction with Cp*H as an additive, in the absence of RhCl₃, did not promote a reaction (entry 12).

We turned our attention to explore the substrate scope of the methylation reaction (Table 2) and found that a ketone with a shorter ethyl sidechain gave the product (2b) in 73% yield. When varying the electronic properties of the aromatic ring, good to high yields were obtained with electron-rich (2d,

> 2h, 2k) and electron-poor (2c, 2e, 2g, 2i, 2j) rings and with heteroaromatic alkyl ketones (21). We found that electron-poor substrates required a lower catalyst loading as well as shorter reaction times to achieve optimal yields. For example, 2i was obtained in 73% yield with 2.5 mol% of [{Cp*RhCl₂}] after $2\,h$ versus in only $41\,\%$ yield with $5\,mol\,\%$ of [{Cp*RhCl₂]₂] after 4 h.^[21] We think that this variation derives from 1,2-reduction of the carbonyl group in both starting ketone and product and leads to an undesired consumption of material. On the other hand, an electron-rich aniline substrate (2h) required portionwise catalyst addition and a longer reaction time to reach full conversion. Notably, halogen-substituted substrates (2c, 2e, 2g, 2j) are tolerated under the reaction conditions.

> Many products **2** were formed in good to excellent yields, significantly extending the scope of the reaction and showing that other primary alkyl chains were compatible with the methylation. Furthermore, the reaction was found to work well with β -substitution on the alkyl chains: high to excellent yields of the α -methylation products **2m**,**n** were obtained with KOH as a base. Unfortunately, a stereogenic center at the β -position did not induce diastereoselectivity under these reaction conditions.

We also investigated the methylation of alkyl/ alkyl ketone substrates. As alkylation does not occur on secondary centers, site-selective alkylation was accomplished on α -n-alkyl- α '-dialkyl ketones in 81 % and 49 % yields for **2s** and **2u**, respectively. The lower yield obtained for **2u** may be due to the higher p K_a value of the dialkylketone versus the aryl-alkyl ketones. In this case, the slower reaction led to decomposition of both substrate and product under the reaction conditions. With double (α,α') dimethylation reactions we obtained **2t** in 55 % yield (74 % yield per methylation), although no 1,3-diastereoselectivity was observed.

During our investigation, we noted that several of the starting ketones required multi-step synthesis, usually via the corresponding Weinreb amides. The wide commercial accessibility of many methyl ketones encouraged us to examine the double alkylation of these substrates (Scheme 2A). Ketone **2k** was synthesized directly from the corresponding methyl ketone in 56% yield (i.e., 75% yield per methylation). Direct access to isopropyl ketones **2v** and **2w** from commercially

Table 2: Scope of the α -methylation of ketones: structures of the products formed. All yields are of isolated material and d.r. was determined from ¹H NMR spectroscopy of the crude reaction mixture.



[a] [{Cp*RhCl₂}₂] (7.5 mol%) was added over 72 h in 3 portions. [b] 2 equiv of Cs_2CO_3 was used. [c] S_NAr product (2 k) was also isolated in 21 % yield. [d] 5 equiv of KOH was used as a base instead of Cs₂CO₃.

available material was achieved in 53% and 44% yields, respectively.

To extend this concept further, we examined the conditions for ketone alkylation reported by Ishii et al.^[4a] and combined them with our methylation reaction in a one-pot procedure (Scheme 2B). This sequence further demonstrates the orthogonal reactivity between the new system and the existing alkylation methods which use hydrogen-borrowing chemistry (and which do not form branched products). We were able to perform the one-pot consecutive alkylation of methyl ketones using tandem alcohol alkylation reactions, firstly with a benzyl (2r, 2s, 2o) or primary alkyl alcohol (2q) and secondly with methanol. Once the first alkylation was complete (Ir catalysis; no over-alkylation observed), methanol, oxygen, Cs₂CO₃, and Rh catalyst were added directly to the reaction mixture without any solvent removal or purification.^[5] Moderate to good yields of doubly alkylated products 20,q,r,s were obtained in this one-pot process (note that the overall yields are comparable to the multiples of the individual steps). We suggest that this double alkylation method holds considerable promise for the generation of

Angew. Chem. Int. Ed. 2014, 53, 761-765

© 2014 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

complex carbonyl compounds from simple methyl ketone precursors.

It is important to expand the synthetic utility of the methylation reaction so that other alkylated carbonyl derivatives can be prepared. At this point we have only investigated the methylation of ketones, however, we were able to transform the products formed through this reaction into ester-derived products in a single, high-yielding step. Regioselective Baeyer-Villiger oxidation of 2k and 20 with mCPBA was accomplished in 98% and 92% yield, respectively (Scheme 2C), thus providing access to α -branched alkylated (and activated) esters (7k, 7o).^[4d]



Scheme 2. A) Rh-catalyzed double methylation. B) One-pot sequential Ir/ Rh-catalyzed dialkylation; typical conditions: 1 mol% of [{IrCl(cod)}₂], KOH, Ph₃P, 1.2–5 equiv of alcohol, 100°C, 4 h; then 5 mol% of [{Cp*RhCl₂}₂], 5 equiv of Cs₂CO₃, MeOH, 65 °C, O₂. In addition to the overall yields, those of the individual steps are given. C) Baeyer-Villiger oxidation.-See the Supporting Information for experimental details; structures of the products shown.

To provide insight into the reaction mechanism, we carried out a series of reactions whereby the proposed (and in some cases observed) intermediates were resubjected to the reaction conditions (Scheme 3). In every case but one, we found that these compounds were converted into the methylation product 2a, providing evidence for their participation in a reaction sequence involving an aldol reaction with formaldehyde, elimination, and alkene reduction. However, when we used the reaction conditions on 3a (which might derive from the conjugate addition of an enolate to enone

www.angewandte.org

763



Scheme 3. Mechanistic experiments.

4a), we recovered only starting material, which suggests that3a is a dead-end that does not contribute to the formation of methylated product 2a.

At this point, we interpret our results as shown below (Scheme 4). The key steps are methanol oxidation to formaldehyde, followed by an aldol reaction and elimination to



Scheme 4. A preliminary mechanistic interpretation of the results. The exact nature of the ligands on the active Rh catalyst is not currently known.^[24]

form enone 4. Reduction of the enone by hydride appears to be competitive with the addition of methanol, although we cannot rule out the formation of 5 directly from 1 and methanol through oxidation.

Further investigation is required to elucidate the exact role of oxygen in this process, although we would draw attention to a report from Gabrielsson et al. who showed that methanol oxidation can be achieved under aerobic and mild conditions with an Ir catalyst.^[22] One working hypothesis at this point is that the oxygen may reoxidize any Rh^I species formed by deprotonation (and elimination of a leaving group) of Rh^{III}–H, thus regenerating the active catalyst in situ, although other possibilities exist.^[23]

Finally, it should be noted that the reactive and unhindered nature of formaldehyde as an electrophile may explain why this system is uniquely effective for making branched alkyl products, as only this particular electrophile is able to couple effectively with a substituted enolate under these conditions.

In conclusion, we have reported a method for the use of methanol as an alkylating reagent in a rhodium-catalyzed reaction. We found that an atmosphere of oxygen is beneficial to the yields of methylation and have shown that methanol oxidation could be achieved under remarkably mild conditions. The scope of this reaction includes aromatic and aliphatic ketones and consecutive one-pot double alkylation reactions provide a convenient route to branched ketones from simple methyl ketones. A brief study into the mechanism of the reaction has given evidence for an aldol-based reaction pathway.

Received: September 10, 2013 Published online: November 29, 2013

Keywords: homogeneous catalysis \cdot hydrogen borrowing \cdot ketones \cdot methanol

- a) M. H. S. A. Hamid, P. A. Slatford, J. M. J. Williams, Adv. Synth. Catal. 2007, 349, 1555; b) T. D. Nixon, M. K. Whittlesey, J. M. J. Williams, Dalton Trans. 2009, 753; c) G. Guillena, D. J. Ramón, M. Yus, Angew. Chem. 2007, 119, 2410; Angew. Chem. Int. Ed. 2007, 46, 2358; d) F. Shibahara, M. J. Krische, Chem. Lett. 2008, 37, 1102; e) J. F. Bower, I. S. Kim, R. L. Patman, M. J. Krische, Angew. Chem. 2009, 121, 36; Angew. Chem. Int. Ed. 2009, 48, 34; for a review of the different strategies involved, see f) C. Gunanathan, D. Milstein, Science 2013, 341, 1229712.
- [2] a) C. E. Bibby, R. Grigg, R. Price, J. Chem. Soc. Dalton Trans.
 1977, 872; b) R. Grigg, T. R. B. Mitchell, S. Sutthivaiyakit, Tetrahedron Lett. 1979, 20, 1067; c) R. Grigg, T. R. B. Mitchell, S. Sutthivaiyakit, N. Tongpenyai, J. Chem. Soc. Chem. Commun.
 1981, 611; d) R. Grigg, T. R. B. Mitchell, S. Sutthivaiyakit, Tetrahedron 1981, 37, 4313; e) R. Grigg, T. R. B. Mitchell, S. Sutthivaiyakit, N. Tongpenyai, Tetrahedron Lett. 1981, 22, 4107; f) C. Löfberg, R. Grigg, M. Whittaker, A. Keep, A. Derrick, J. Org. Chem. 2006, 71, 8023; g) C. Löfberg, R. Grigg, A. Keep, A. Derrick, V. Sridharan, C. Kilner, Chem. Commun. 2006, 5000.
- [3] a) C. S. Cho, B. T. Kim, T.-J. Kim, S. C. Shim, J. Org. Chem. 2001, 66, 9020; b) C. S. Cho, B. T. Kim, T.-J. Kim, S. C. Shim, Tetrahedron Lett. 2002, 43, 7987; c) C. S. Cho, J. Mol. Catal. A 2005, 240, 55.
- [4] a) K. Taguchi, H. Nakagawa, T. Hirabayashi, S. Sakaguchi, Y. Ishii, *J. Am. Chem. Soc.* 2004, *126*, 72; b) K. Maeda, Y. Obora, S. Sakaguchi, Y. Ishii, *Bull. Chem. Soc. Jpn.* 2008, *81*, 689; c) M. Morita, Y. Obora, Y. Ishii, *Chem. Commun.* 2007, 2850; d) Y. Iuchi, Y. Obora, Y. Ishii, *J. Am. Chem. Soc.* 2010, *132*, 2536.
- [5] G. Onodera, Y. Nishibayashi, S. Uemura, Angew. Chem. 2006, 118, 3903; Angew. Chem. Int. Ed. 2006, 45, 3819.
- [6] a) R. Martínez, G. L. Brand, D. J. Ramón, M. Yus, *Tetrahedron Lett.* 2005, 46, 3683; b) R. Martínez, D. J. Ramón, M. Yus, *Tetrahedron* 2006, 62, 8988.
- [7] M. S. Kwon, N. Kim, S. H. Seo, I. S. Park, R. K. Cheedrala, J. Park, Angew. Chem. 2005, 117, 7073; Angew. Chem. Int. Ed. 2005, 44, 6913.

- [8] a) Y. M. A. Yamada, Y. Uozumi, Org. Lett. 2006, 8, 1375;
 b) Y. M. A. Yamada, Y. Uozumi, Tetrahedron 2007, 63, 8492.
- [9] a) M. G. Edwards, J. M. J. Williams, Angew. Chem. 2002, 114, 4934; Angew. Chem. Int. Ed. 2002, 41, 4740; b) P. J. Black, G. Cami-Kobeci, M. G. Edwards, P. A. Slatford, M. K. Whittlesey, J. M. J. Williams, Org. Biomol. Chem. 2006, 4, 116; c) P. A. Slatford, M. K. Whittelsey, J. M. J. Williams, Tetrahedron Lett. 2006, 47, 6787; d) S. J. Pridmore, J. M. J. Williams, Tetrahedron Lett. 2008, 49, 7413.
- [10] a) X. Cui, Y. Zhang, F. Shi, Y. Deng, *Chem. Eur. J.* 2011, *17*, 1021;
 b) T. Kuwahara, T. Fukuyama, I. Ryu, *Org. Lett.* 2012, *14*, 4703.
- [11] The transformation of alcohols into more-substituted derivatives using vigorous basic conditions and without a transition metal catalyst (the Guerbet reaction) has been known and studied for some time, see: a) S. Veibel, J. I. Nielson, *Tetrahedron* **1967**, *23*, 1723; b) L. J. Allen, R. H. Crabtree, *Green Chem.* **2010**, *12*, 1362.
- [12] M. Nielsen, E. Alberico, W. Baumann, H.-J. Drexler, H. Junge, S. Gladiali, M. Beller, *Nature* 2013, 495, 85, and references therein.
- [13] a) R. Langer, I. Fuchs, M. Vogt, E. Balaraman, Y. Diskin-Posner, L. J. W. Shimon, Y. Ben-David, D. Milstein, *Chem. Eur. J.* 2013, 19, 3407; b) C. Gunanathan, D. Milstein, *Acc. Chem. Res.* 2011, 44, 588.
- [14] R. E. Rodríguez-Lugo, M. Trincado, M. Vogt, F. Tewes, G. Santiso-Quinones, H. Grützmacher, *Nat. Chem.* 2013, 5, 342.
- [15] N. Ortega, C. Richter, F. Glorius, Org. Lett. 2013, 15, 1776.
- [16] a) K. Motokura, N. Fujita, K. Mori, T. Mizugaki, K. Ebitani, K. Jitsukawa, K. Kaneda, *Chem. Eur. J.* 2006, *12*, 8228; for related

reactions that form branched products from α -aryl nitriles, but do not use methanol, see Ref. [2f].

- [17] a) J. Moran, A. Preetz, R. A. Mesch, M. J. Krische, *Nat. Chem.* **2011**, *3*, 287; b) F. Li, J. Xie, H. Shan, C. Sun, L. Chen, *RSC Adv.* **2012**, *2*, 8645.
- [18] Slow conversion of **1a** was also observed at 50 °C, indicating that reaction is possible at that temperature.
- [19] a) S. J. Taylor, M. O. Duffey, J. P. Morken, J. Am. Chem. Soc. 2000, 122, 4528; b) J. S. M. Samec, J.-E. Bäckvall, P. G. Andersson, P. Brandt, Chem. Soc. Rev. 2006, 35, 237.
- [20] X. Li, L. Li, Y. Tang, L. Zhong, L. Cun, J. Zhu, J. Liao, J. Deng, J. Org. Chem. 2010, 75, 2981.
- [21] The methylation of 1i was carried out on a 2.5 mmol scale with 2.5 mol% of [{Cp*RhCl₂}] in 70% yield. See the Supporting Information for details.
- [22] a) A. Gabrielsson, P. van Leeuwen, W. Kaim, *Chem. Commun.* **2006**, 4926; another report details further mechanistic possibilities for oxygen participation: b) B. Jiang, Y. Feng, E. A. Ison, *J. Am. Chem. Soc.* **2008**, *130*, 14462.
- [23] Y. Iwashita, A. Hayata, J. Am. Chem. Soc. 1969, 91, 2525.
- [24] For a study of the role of carbonate in related oxidation reactions, see: D. Balcells, A. Nova, E. Clot, D. Gnanamgari, R. H. Crabtree, O. Eisenstein, *Organometallics* 2008, 27, 2529. Although we have shown the fate of the hydrogen generated from methanol as being the formation of Rh-H and a proton, at this point we do not rule out a MH₂ species.