

# CHEMISTRY

## A European Journal

A Journal of



### Accepted Article

**Title:** Perfluoro Alkyl Hypofluorites and Peroxides Revisited

**Authors:** Sebastian Hasenstab-Riedel, Jan Hendrik Nissen, Thomas Drews, Benjamin Schröder, Helmut Beckers, and Simon Steinhauer

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

**To be cited as:** *Chem. Eur. J.* 10.1002/chem.201903620

**Link to VoR:** <http://dx.doi.org/10.1002/chem.201903620>

Supported by  
**ACES**

WILEY-VCH

## FULL PAPER


## Perfluoro Alkyl Hypofluorites and Peroxides Revisited

Jan H. Nissen, Thomas Drews, Benjamin Schröder, Helmut Beckers, Simon Steinhauer, and Sebastian Riedel<sup>[a]</sup>

**Abstract:** A more convenient synthesis of the perfluoro alkyl hypofluorite (F<sub>3</sub>C)<sub>3</sub>COF as well as the hitherto unknown (C<sub>2</sub>F<sub>5</sub>)(F<sub>3</sub>C)<sub>2</sub>COF compound is reported. Both hypofluorites can be prepared by use of the corresponding tertiary alcohols R<sup>F</sup>OH and elemental fluorine in the presence of CsF. An appropriate access to these highly reactive hypofluorites is crucial. The hypofluorites are then transferred into their corresponding perfluoro bisalkyl peroxides R<sup>F</sup>OOR<sup>F</sup> [R<sup>F</sup> = (F<sub>3</sub>C)<sub>3</sub>C, (C<sub>2</sub>F<sub>5</sub>)(F<sub>3</sub>C)<sub>2</sub>C] by treatment with partially fluorinated silver wool. NMR, gas-phase infrared and solid-state Raman spectra of the perfluoro bisalkyl peroxides are presented and their chemical properties are discussed.

## Introduction


Cady has described the synthesis of the first perfluoro alkyl hypofluorite, trifluoromethyl hypofluorite, F<sub>3</sub>COF, by the AgF<sub>2</sub> catalyzed direct fluorination of methanol in 1948.<sup>[1]</sup> Since then a variety of perfluoro<sup>[2]</sup> as well as partially chlorinated (e. g. Cl<sub>3</sub>CCF<sub>2</sub>OF)<sup>[2,3]</sup> and nitrogen (e. g. NF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>OF)<sup>[4]</sup> or sulfur (e. g. FSO<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>OF)<sup>[5]</sup> containing and later even hydrogen (e. g. H<sub>3</sub>COF)<sup>[6]</sup> substituted alkyl hypofluorites ROF (**1**) have been isolated and characterized. An alternative route to perfluoro alkyl hypofluorites is the fluorination of perfluoro ketones in the presence of metal fluorides, MF (M = K, Rb, Cs) [Eq. (1)], as described in 1966 by Ruff and Lustig.<sup>[7]</sup> Moreover, the direct fluorination (with 10% F<sub>2</sub> in N<sub>2</sub>) of sodium trifluoroacetate, F<sub>3</sub>CCO<sub>2</sub>Na,<sup>[8]</sup> or trifluoroacetic acid, F<sub>3</sub>CCO<sub>2</sub>H,<sup>[9]</sup> leads to formation of hypofluorite compounds in a temperature dependent ratio. These hypofluorites were used in the pioneering work of Hesse<sup>[10]</sup> and Rozen<sup>[11]</sup> as electrophilic fluorination agents or as etching gas<sup>[12]</sup> for semiconductors. The relatively low dissociation energy of the O–F bond in F<sub>3</sub>COF (184.2 kJ mol<sup>−1</sup>)<sup>[13]</sup> facilitates insertion of carbon monoxide<sup>[14]</sup> into the O–F bond to produce fluoroformates, F<sub>3</sub>COC(O)F, and the addition of F<sub>3</sub>COF to (per)fluorinated olefins which yield (per)fluorinated bisalkyl ethers, F<sub>3</sub>COR.<sup>[15]</sup> Further reactions of functionalized ethers like F<sub>3</sub>COCFCI–CF<sub>2</sub>Cl may then be reduced to the vinyl ether,



F<sub>3</sub>COCF=CF<sub>2</sub>, which has become a valuable fluorinated monomer for the industrial synthesis of perfluoroxy alkanes (PFA).<sup>[16]</sup>

Hypofluorites are highly hazardous compounds, which requires sophisticated handling and safety precautions. A high reactivity and oxidation power is often paired with an intrinsic instability which may lead to a rapid exothermic decomposition by contact to organic impurities or metal surfaces.<sup>[17]</sup> Decomposition of R<sup>F</sup>OF to yield the corresponding carbonyl compounds [Eq. (1)] is highly exothermic with reaction enthalpies ΔH<sub>R</sub> of up to −400 kJ mol<sup>−1</sup> (see Supporting Information, Table S2.1). The susceptibility to hydrolysis of perfluorinated hypofluorites, R<sup>F</sup>OF, to alcohols increases with increasing numbers of fluorinated groups [Scheme (1)]. Trifluoromethyl hypofluorite, F<sub>3</sub>COF, is thermally rather stable up to temperatures above 450 °C and hydrolysis in aqueous solution is very slow.<sup>[1]</sup> The chemistry of perfluoro alkyl hypofluorites has been recently reviewed.<sup>[17]</sup>

**Scheme 1:** Increasing tendency of hydrolysis of selected perfluoro alkyl hypofluorites, R<sup>F</sup>OF.



We found that perfluoroalkyl hypofluorites R<sup>F</sup>OF are excellent starting compounds for the synthesis of otherwise difficult accessible perfluoro bisalkyl peroxides R<sup>F</sup>OOR<sup>F</sup> (**2**),<sup>[18]</sup> which are synthetically valuable sources of fluoroalkoxy radicals, R<sup>F</sup>O•.<sup>[19]</sup> Until very recently<sup>[18]</sup> our knowledge about perfluoro bisalkyl peroxides was very scarce and limited to the two homologues bis(trifluoromethyl) peroxide, (F<sub>3</sub>CO)<sub>2</sub> (**2a**)<sup>[20]</sup> and bis(nonafluoro-*tert*-butyl) peroxide, [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> (**2b**).<sup>[21]</sup> The volatile and rather resistant peroxide **2a** (bp. −37 °C) was reported in 1933 by Swarts<sup>[22]</sup> and prepared in 1957 by Cady *et al.*<sup>[23]</sup> from trifluoromethyl hypofluorite and carbonyl difluoride, F<sub>2</sub>CO. One decade later Anderson *et al.* detected peroxide **2b** among other products obtained by the reaction of chlorine trifluoride and perfluoro *tert*-butyl alcohol, (F<sub>3</sub>C)<sub>3</sub>COH (**3b**), in a high-pressure stainless-steel reactor.<sup>[21]</sup> Later also the low-temperature photolysis (−40 °C) of perfluoro *tert*-butyl hypofluorite, (F<sub>3</sub>C)<sub>3</sub>COF (**1b**), in the presence of tetrafluorohydrazine, N<sub>2</sub>F<sub>4</sub>, was described to yield peroxide **2b**.<sup>[19]</sup> The gas-phase molecular structure of **2a** was investigated by electron diffraction<sup>[20]</sup> while the solid-state structure of **2b** has only very recently been published.<sup>[18]</sup> The chemistry of these perfluoroalkyl peroxides have only been scarcely investigated in the past. Among them are reactions with carbon-carbon double bonds of fluorinated olefins<sup>[24]</sup> and thiophenes.<sup>[25]</sup> Furthermore, the closely related compound class of fluoroformyl peroxides, R<sup>F</sup>–OO–C(O)F,<sup>[26]</sup> as well as perfluorodiacyl peroxides, R<sup>F</sup>C(O)–OO–C(O)R<sup>F</sup> have been

[a] J. H. Nissen, T. Drews, B. Schröder, H. Beckers, S. Steinhauer, Prof. S. Riedel  
 Fachbereich für Biologie, Chemie, Pharmazie, Institut für Chemie und Biochemie – Anorganische Chemie, Freie Universität Berlin, Fabeckstraße 34/36, 14195 Berlin (Germany)  
 E-mail: [s.riedel@fu-berlin.de](mailto:s.riedel@fu-berlin.de)  
 Supporting information for this article is given via a link at the end of the document.

## FULL PAPER


investigated and represent important intermediates in the industrial synthesis of fluorous polyether chains<sup>[27]</sup>.

Here we report a modified synthesis of tertiary perfluoroalkyl hypofluorites  $R^F OF$  [ $R^F = (F_3C)_3C$  **1b**,  $R^F = (C_2F_5)(F_3C)_2C$  **1c**] from the corresponding alcohols  $R^F OH$  [ $R^F = (F_3C)_3C$  **3b**,  $R^F = (C_2F_5)(F_3C)_2C$  **3c**] with elemental fluorine in the presence of excess CsF. These hypofluorites **1** were then treated with fluorinated silver wool to form the perfluoro bisalkyl peroxides  $R^F OOR^F$  [ $R^F = (F_3C)_3C$  (**2b**),  $R^F = (C_2F_5)(F_3C)_2C$  (**2c**)].

## Results and Discussion


Perfluoroalkyl hypofluorites,  $R^F OF$  (**1**), were obtained according to Eq. (1) by treatment of the corresponding carbonyl compound with elemental fluorine in a stainless steel reactor under the presence of any alkali metal fluoride, MF. The mixture was initially cooled to  $-196^\circ C$ , and then slowly heated to  $-78^\circ C$ .<sup>[17,28,29]</sup> Because these mixtures are highly hazardous and may explode spontaneously,<sup>[28]</sup> we first improved this synthesis to avoid any possible local heat formation and pressure increase during the reaction.

The number of perfluoroalkyl hypofluorites **1** obtainable by this procedure can be further increased by using perfluoro alkylalcohols **3** instead of the perfluoro carbonyl compounds according to Eq. (2). First, the perfluoro alkylalcohols **3b,c** are added to an excess of caesium fluoride in a stainless steel reactor and the mixture is thoroughly shaken at room temperature in order to dissipate the reaction heat of the subsequent fluorination reaction. It is assumed that the alcohol reacts with CsF to the corresponding caesium alcoholate and  $CsHF_2$ .<sup>[29]</sup> The reactor is cooled to  $-78^\circ C$  again and elemental fluorine is added in small portions via a stainless steel line until no fluorine is consumed anymore and the pressure remains constant. Excess fluorine is then removed from the reaction mixture at  $-196^\circ C$  and the hypofluorites **1** thus formed are distilled out of the reactor and purified by trap-to-trap distillation. For the hypofluorites  $(F_3C)_3COF$  (**1b**) and  $(C_2F_5)(F_3C)_2COF$  (**1c**) the conversion is quantitative.




This fluorination procedure was finally also used for the synthesis of the known hypofluorites  $F_3COF$  (**1a**),  $CF_3CF_2OF$  (**1d**) and  $(F_3C)_2CFOF$  (**1e**) starting from the perfluoro carbonyl compounds  $F_2CO$ ,  $F_3CC(O)F$  and  $(F_3C)_2CO$ , respectively, according to Eq. (1). Their IR and NMR spectra match those previously reported.<sup>[2]</sup> Quantum-chemical calculations indicate that apart from **1a** all these perfluoro alkyl hypofluorites are thermodynamically unstable by up to  $-400 \text{ kJ mol}^{-1}$  in terms of elimination of  $CF_4$  and formation of the corresponding carbonyl compounds (see Supporting Information, Table S2.1). As expected, elimination of elemental fluorine is endothermic by 120

to  $150 \text{ kJ mol}^{-1}$ . Nevertheless, these perfluoro alkyl hypofluorites are kinetically stable in the gas-phase up to  $110^\circ C$ .<sup>[2]</sup> Therefore, we were able to provide APCI mass spectra of *tert*-butyl hypofluorite **1b** (see Supporting Information, Figure S1.2). The predominant species in the negative mode is the alkoxide ion  $[(F_3C)_3CO]^-$  ( $m/z = 235$ ), while the molecular ion peak at  $m/z = 254$  is not observed. Interestingly, the peak at  $m/z = 285$  can be assigned to an adduct or insertion of a  $CF_2$ -fragment to the alkoxide ion  $[(F_3C)_3CO + CF_2]^-$ , which was previously observed for an perfluorinated ether bearing the perfluoro-*tert*-butyl group.<sup>[30]</sup> Only perfluoro *tert*-pentyl hypofluorite (**1c**), which is stable in solution for several minutes, is found to decompose readily at room temperature within seconds to yield selectively  $(F_3C)_2CO$  and  $C_2F_6$ , as proved by their gas-phase IR spectra [Eq. (3)].<sup>[31],[32]</sup>




Probably due to the lack of stability, hypofluorite **1c** has not yet been characterized.<sup>[10,33]</sup> We therefore present low-temperature NMR as well as the gas-phase IR spectrum of this poorly known compound. These spectra clearly confirm the presence of **1c**. The fully coupled  $^{13}C$  NMR spectrum (see Supporting Information, Figure S1.3) shows two quartets at  $\delta = 118.5 \text{ ppm}$  ( $C-CF_3$ ) and  $\delta = 116.8 \text{ ppm}$  ( $F_3C-CF_2$ ), where the latter one partly overlaps with the  $CF_2$ -triplet at  $\delta = 109.8 \text{ ppm}$ . The chemical shift of the quaternary carbon atom appears at  $\delta = 87.7 \text{ ppm}$ . The first order  $^{19}F$  NMR spectrum of hypofluorite **1c** (Figure 1) shows a doublet of septets for the  $CF_2$  nuclei at  $\delta = -117.8 \text{ ppm}$ , indicating a  $^4J(F,F)$  coupling to  $OF$  and to the two  $CF_3$  nuclei of 11.6 Hz. The  $F_3CCF_2$  signal at  $\delta = -81.9 \text{ ppm}$  also shows a doublet of septets with similar coupling constants of  $^5J(F,F) = 5.9 \text{ Hz}$  to both, the  $OF$  and the  $C-CF_3$  nuclei. The signal for the two  $CF_3$  groups occurs as a doublet of triplet of quartets (d t q) at  $\delta = -68.3 \text{ ppm}$  with a large  $^4J$  coupling constant of 17.7 Hz to the  $OF$  nucleus, which resonates at  $\delta = 150.8 \text{ ppm}$  (t sept q). The  $^3J$  coupling constant between the fluorine atoms of the pentafluoroethyl group is smaller than 0.5 Hz.




**Figure 1.**  $^{19}F$  NMR spectrum of  $(C_2F_5)(F_3C)_2COF$  (**3c**) (376.88 MHz, external  $[D_6]Acetone$ ,  $-60^\circ C$ ).

## FULL PAPER



**Figure 2.** Gas-phase IR spectrum of  $(\text{C}_2\text{F}_5)(\text{F}_3\text{C})_2\text{COF}$  (**3c**) (bottom) and the calculated spectrum for the most stable *t*-1 conformer (see Figure 3) at the B3LYP/aug-cc-pVTZ level of theory (top trace).

The IR spectrum of hypofluorite **1c** in the gas-phase is compared to a computed IR spectrum at the DFT-B3LYP/aug-cc-pVTZ level of theory in Figure 2. The experimental bands in the mid-IR from 1107 to 878  $\text{cm}^{-1}$  are split into two components, probably due to the presence of at least two rotational conformers in the gas-phase. Indeed, the DFT calculations revealed slightly different IR spectra for the different *trans* and *gauche* rotational conformers of **1c** (Figure 3, Table 1 and the SI) and a global minimum for the *t*-1 structure. This result agrees well with the experimental IR spectrum, which shows strong absorption for the C–F stretching modes in the region from 1291 to 1232  $\text{cm}^{-1}$  and the characteristic deformation bands of the  $\text{CF}_3$ -groups at 766, 743, and 730  $\text{cm}^{-1}$ . Furthermore, the bands at 1107 and 1078  $\text{cm}^{-1}$  are tentatively assigned to C–O stretching modes of different rotamers of **1c**. According to the calculations the O–F stretching mode has a relatively low intensity and its position varies up to 43  $\text{cm}^{-1}$  depending on the conformer of **1c** (see Table 1 and SI). It can tentatively be assigned in the experimental gas-phase IR spectrum to weak absorptions at 925 and 913  $\text{cm}^{-1}$ , but we cannot exclude that this band is superimposed by the asymmetric stretching mode of the  $\text{C}-(\text{CF}_3)_2$  fragment located at 1003  $\text{cm}^{-1}$ . The bands at 895 and 878  $\text{cm}^{-1}$  represent C–C stretching modes of the pentafluoroethyl group of **1c**. The calculated position of this band varies for the different conformers by 36  $\text{cm}^{-1}$ . The weak band at 613  $\text{cm}^{-1}$  can be associated with the  $\text{CC}_3$  deformation, while weaker deformation modes of the  $\text{CF}_3$ -group are found at 539 and 513  $\text{cm}^{-1}$ . A very weak absorption at 484  $\text{cm}^{-1}$  fits well to a computed deformation mode of the  $\text{C}_2\text{F}_5$ -group.



**Figure 3.** Relative energies of *trans* and *gauche* conformers of  $(\text{C}_2\text{F}_5)(\text{F}_3\text{C})_2\text{COF}$  (**1c**) obtained at the B3LYP/aug-cc-pVTZ level of theory (fluorine atoms bound to carbon are not shown).

**Table 1.** Gas-phase vibrational frequencies  $\tilde{\nu}$  [ $\text{cm}^{-1}$ ] and relative IR band intensities<sup>[a]</sup> compared to computed values for different rotational conformers of  $(\text{C}_2\text{F}_5)(\text{F}_3\text{C})_2\text{COF}$  (**1c**) at the B3LYP/aug-cc-pVTZ level of theory.<sup>[b]</sup>


experiment	DFT		assignment
	<i>t</i> -1	<i>g</i> -1	
1342 (m)	1290 (75)	1308 (22)	$\nu(\text{F}_2\text{C}-\text{CF}_3)$
1291 (s,sh)	1273 (272)	1260 (421)	$\nu(\text{CF}_3)$
1276 (vs)	1248 (606)	1246 (541)	$\nu(\text{CF}_3)$
1262 (vs)	1239 (403)	1237 (338)	$\nu(\text{CF}_3)$
1255 (vs)	1233 (58)	1221 (157)	$\nu(\text{CF}_3)$
1251 (vs)	1218 (110)	1216 (240)	$\nu(\text{CF}_3)$
1232 (s)	1197 (297)	1201 (256)	$\nu(\text{CF}_3)$
1181 (m)	1161 (25)	1155 (5)	$\nu(\text{CF}_3)$
1161 (w, sh)	1154 (63)	1153 (11)	$\nu(\text{CF}_2)$
1107 (m), 1078 (w)	1064 (121)	1113 (14)	$\nu(\text{CO})$
1038 (vw)	1055 (12)	1047 (49)	$\nu(\text{C}-\text{CF}_2)$
1003 (m), 977 (w)	972 (100)	963 (93)	$\nu_{\text{as}}(\text{C}-(\text{CF}_3)_2)$
925 (w), 913 (w)	995 (25)	967 (34)	$\nu(\text{OF})$
895 (m), 878 (m)	888 (117)	898 (150)	$\nu(\text{F}_2\text{C}-\text{CF}_3)$
766 (vw)	766 (5)	765 (1)	$\delta(\text{CF}_3)$
743 (m)	739 (59)	740 (59)	$\delta(\text{CF}_3)$
730 (m)	725 (39)	725 (40)	$\delta(\text{C}(\text{CF}_3)_2)$
613 (vw)	619 (6)	621 (12)	$\delta(\text{CC}_3)$
558 (sh)	537 (4)	537 (3)	$\delta(\text{CF}_3)$
539 (vw)	525 (9)	527 (10)	$\delta(\text{CF}_3)$
513 (vw)	501 (7)	499 (8)	$\delta(\text{CF}_3)$
484 (vw)	444 (2)	445 (3)	$\delta(\text{C}_2\text{F}_5)$

[a] Relative intensities in parentheses: vw = very weak, w = weak,

m = medium, s = strong, vs = very strong, sh = shoulder.

[b] For the different *trans* and *gauche* rotational conformers see Figure 3 and Figures S2.

The hypofluorites **1b** and **1c** are converted in rather good yield up to <70% into the corresponding symmetric perfluoro bis-alkyl peroxides **2b** and **2c**, respectively, by using partially fluorinated silver wool [see Exp. Details and Eq. (4)]. This reaction does not proceed at low temperatures of  $-78^\circ\text{C}$ , while at  $0^\circ\text{C}$  mainly decomposition products of the hypofluorites are formed [**1b**:  $\text{CF}_4$ ,  $(\text{F}_3\text{C})_2\text{CO}$ ; **1c**:  $\text{C}_2\text{F}_6$ ,  $(\text{F}_3\text{C})_2\text{CO}$ ].<sup>[31,32,34]</sup> When the reaction vessel is held at temperatures of  $-50$  to  $-45^\circ\text{C}$  for 48 to 72 h, peroxide **2b** is obtained from **1b** and separated by trap-to-trap distillation in a  $-78^\circ\text{C}$  trap from the more volatile side products  $\text{CF}_4$  and  $(\text{F}_3\text{C})_2\text{CO}$  in an overall yield of >70%. Pure peroxide **2b** is rather stable at ambient temperature and decomposes at temperatures above  $100^\circ\text{C}$  to yield  $(\text{F}_3\text{C})_2\text{CO}$  and  $\text{C}_2\text{F}_6$  with an activation energy of  $148.7 \pm 4.4 \text{ kJ mol}^{-1}$ .<sup>[35]</sup> Similarly, hypofluorite **1c** reacts to the peroxide **2c** and it was purified from the volatile side products  $\text{C}_2\text{F}_6$  and  $(\text{F}_3\text{C})_2\text{CO}$  by trap-to-trap distillation, where it remains in a  $-78^\circ\text{C}$  trap in a yield of up to 66%.




However, attempts to convert the hypofluorites  $\text{CF}_3\text{CF}_2\text{OF}$  (**1d**) and  $(\text{F}_3\text{C})_2\text{CFOF}$  (**1e**) with fluorinated silver wool under similar conditions to the corresponding perfluoro bisalkyl peroxides failed and led solely to decomposition products (**1d**:  $\text{CF}_4$ ,<sup>[34]</sup>  $\text{F}_2\text{CO}$ ,<sup>[36]</sup> **1e**:  $\text{CF}_4$ ,<sup>[34]</sup>  $\text{F}_3\text{CC}(\text{O})\text{F}$ <sup>[37]</sup>) which were identified by IR spectroscopy. The composition of the fluorinated silver wool used in the synthesis of the peroxides **2b** and **2c** according to Eq. (4) prepared as described in the experimental details was



## FULL PAPER

investigated by a powder X-ray diffraction analysis. It consists mainly of silver(I) fluoride, AgF, but contains also some silver subfluoride, Ag<sub>2</sub>F, silver(II) fluoride, AgF<sub>2</sub>, and elemental silver, Ag (see Figure S1.1). However, attempts to reproduce the above described conversion of hypofluorites **1** to peroxides **2** by using either elemental Ag or commercial AgF instead of the fluorinated silver wool failed and only decomposition products of the hypofluorites were obtained.


The mechanism of this solid-gas reaction for the formation of peroxides from hypofluorites is still unknown and further studies are necessary to explore this reaction. Powder diffraction measurements of the fluorinated silver wool prior and after several batches (Figure S1.1) indicates an increase of silver(I) fluoride at the expense of silver(0) or silver subfluorides within several reactions. From this result it can be assumed that the active site of the partially fluorinated silver wool consists of an incompletely coordinated silver subfluoride or silver(0) species which acts as a fluorine-atom acceptor. We noticed that the fluorine-atom acceptor ability of this species depletes, and thus, the yield of peroxide formation decreases after several successful batches, very likely due to fluorination of the active silver species and formation of inactive silver(I) fluoride. Based on this assumption, the following reaction mechanism can thus be postulated for the solid-gas reaction: First, the active silver site may undergo an oxidative addition of R<sup>F</sup>OF to form an alkoxide silver(II) intermediate, which then decomposes into alkoxyl radicals, R<sup>F</sup>O•, and silver(I) fluoride [Eq. (5)]. The free or loosely bound R<sup>F</sup>O• radicals may then combine to form the peroxides (**2**).




There are precedents for the formation of silver (II) alkoxides and their decomposition into alkoxyl radicals. Wechsberg previously described the reaction of AgF<sub>2</sub> with F<sub>2</sub>CO and fluorine and assumed the formation of an Ag(II)(OR<sup>F</sup>)<sub>2</sub> intermediate.<sup>[38]</sup> Owing to the high oxidation potential of Ag(II) (electron affinity: 21.45 eV),<sup>[39]</sup> the proposed Ag(II) alkoxide intermediates are prone to a ligand-to-metal electron-transfer (LMCT) and even to the formation of alkoxyl radical intermediates like Ag(I)(O•R<sup>F</sup>)(OR<sup>F</sup>).<sup>[39]</sup> A similar radical mechanism has been proposed for the AgF<sub>2</sub> catalyzed low temperature reaction of F<sub>2</sub> with SO<sub>3</sub> to yield peroxy disulfuryl difluoride, (FSO<sub>2</sub>O)<sub>2</sub>.<sup>[40]</sup>

The <sup>13</sup>C {<sup>19</sup>F} DEPTQ NMR spectrum<sup>[41]</sup> of neat [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> (**2b**) shows two signals at δ = 118.8 and 84.3 ppm associated with the CF<sub>3</sub> and the quaternary carbon nuclei. The <sup>19</sup>F NMR spectrum shows a singlet at δ = -69.6 ppm (lit.:<sup>[42]</sup> -70.0 ppm) while the resonance in the <sup>17</sup>O NMR spectrum occurred in the characteristic region of a peroxide compound<sup>[43]</sup> at δ = 246 ppm, see Figure S1.4. The oxygen atoms of (F<sub>3</sub>CO)<sub>2</sub> (**2a**) resonate at 262 ppm in the <sup>17</sup>O NMR spectrum (Figure S1.5). The base peak in the APCI<sup>-</sup> mass spectrum of **2b** (Figure S1.6) at 235 m/z represents the [(F<sub>3</sub>C)<sub>3</sub>CO]<sup>-</sup> fragment. Also the [(F<sub>3</sub>C)<sub>3</sub>C]<sup>-</sup> fragment can be assigned at 219 m/z, while the molecular ion peak at 470 m/z is not present. The <sup>13</sup>C {<sup>19</sup>F} DEPTQ NMR spectra of [(C<sub>2</sub>F<sub>5</sub>)(F<sub>3</sub>C)<sub>2</sub>CO]<sub>2</sub> (**2c**) (Figure S1.7) with optimized <sup>1</sup>J coupling

constants of 290 Hz and 35 Hz, respectively, show the expected signals for the fluorine substituted carbon atoms at 118.5, 116.7 and 115.2 ppm and the resonance of the quaternary carbon atom at 85.6 ppm slightly shifted to lower field compared to the corresponding hypofluorite **1c**. Three signals observed in the <sup>19</sup>F NMR spectrum (Figure S1.8) are also shifted by Δδ = 2 ppm to lower field compared to the spectrum of the reactant **1c**. This consistent low field shift of the NMR signals underlines the strong electron withdrawing effect of the perfluorinated *tert*-pentyl group of peroxide **2c**.



**Figure 4.** Gas-phase IR spectrum of [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> (**2b**) (bottom) in comparison to a computed spectrum at the B3LYP/aug-cc-pVTZ level (top trace).



**Figure 5.** Gas-phase IR spectrum of [(C<sub>2</sub>F<sub>5</sub>)(F<sub>3</sub>C)<sub>2</sub>CO]<sub>2</sub> (**2c**) (bottom) in comparison to a computed spectrum at the B3LYP/aug-cc-pVTZ level (top trace).

The IR spectrum of [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> (**2b**) in the gas phase is shown in Figure 4 together with the computed spectrum at the DFT-B3LYP/aug-cc-pVTZ level of theory. The strongest absorptions are associated with the CF<sub>3</sub> stretching bands in the region around 1300 cm<sup>-1</sup> (Table S1.1). The sharp IR band at 1110 cm<sup>-1</sup> is assigned to a C–O stretching mode while the second C–O stretch appears at 1129 cm<sup>-1</sup> in the low temperature Raman spectrum (Figure S1.9). The characteristic C–C<sub>3</sub> stretching bands

## FULL PAPER

of the *tert*-butyl group are found at 1002 and 982 cm<sup>-1</sup> in the IR. That agrees well with their computed band positions at the DFT-B3LYP/aug-cc-pVTZ level of theory at 994 and 970 cm<sup>-1</sup>, respectively. The corresponding Raman band shows a strong absorption at 1027 cm<sup>-1</sup>. The calculated O–O stretching mode at 902 cm<sup>-1</sup> can clearly be assigned to a Raman band at 865 cm<sup>-1</sup>. Very strong Raman bands are also found for the symmetric CF<sub>3</sub> deformation modes at 783 and 749 cm<sup>-1</sup>. Their counterparts in the gas-phase IR spectrum of peroxide **2b** appear at 739 and 731 cm<sup>-1</sup>. The asymmetric CF<sub>3</sub> deformation modes are located at 541 and 496 cm<sup>-1</sup> in the IR spectrum and at 569, 541 and 523 cm<sup>-1</sup> in the Raman spectrum. A characteristic deformation of the C–O–O–C peroxide moiety is assigned to a Raman band at 356 cm<sup>-1</sup>, close to the CF<sub>3</sub> rocking modes in the region from 339 to 296 cm<sup>-1</sup>. Two strong Raman bands at 241 and 123 cm<sup>-1</sup> represent CC<sub>3</sub> deformation modes of [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> (**2b**).

Figure 5 shows the IR spectrum of [(C<sub>2</sub>F<sub>5</sub>)(F<sub>3</sub>C)<sub>2</sub>CO]<sub>2</sub> (**2c**) together with the computed spectrum at the DFT-B3LYP/aug-cc-pVTZ level of theory. As expected, it is very similar to the spectrum of **2b** and to that of its precursor **1c**. The IR spectrum shows in addition to the strong CF<sub>3</sub> stretching modes in the region from 1277 to 1229 cm<sup>-1</sup> the characteristic C–C stretching mode of the C<sub>2</sub>F<sub>5</sub> group at 1340 cm<sup>-1</sup>. Weak and broad bands at 1187 and 1177 cm<sup>-1</sup> in the IR and the Raman spectrum (Figure S1.10), respectively, are assigned to stretching modes of the CF<sub>2</sub>-group, and a strong IR absorption at 1105 cm<sup>-1</sup> to the out-of-phase C–O stretching mode. The in-phase C–O and C–CF<sub>2</sub> stretching modes of **2c** are found in the Raman spectrum at 1132 and 1082 cm<sup>-1</sup>, respectively. The corresponding out-of-phase C–CF<sub>2</sub> stretching mode appeared in the IR spectrum at 1086 cm<sup>-1</sup>. Weak to medium intensity bands around 1000 cm<sup>-1</sup> in both, the IR and Raman spectra, are due to CC<sub>3</sub> stretching modes and a strong antisymmetric CF<sub>2</sub> stretching mode is found in the IR Spectrum at 898 cm<sup>-1</sup> (calc.: 929 cm<sup>-1</sup>). The Raman active O–O stretching mode appears at 853 cm<sup>-1</sup>, in excellent agreement with the calculation at 852 cm<sup>-1</sup>, and also the C–O–O–C deformation of the peroxide, located at 352 cm<sup>-1</sup> in the Raman spectrum is very close to that of **2b** (356 cm<sup>-1</sup>).

A full list of all experimental and computed wavenumbers together with a tentative assignment is given in the Supporting Information, Table S1.2.

In previous studies, the synthetically valuable fluorinated radicals F<sub>3</sub>CO<sup>•</sup> or (F<sub>3</sub>C)<sub>3</sub>CO<sup>•</sup> were generated by photolysis of the corresponding peroxide (F<sub>3</sub>CO)<sub>2</sub> (**2a**) or [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> (**2b**), respectively.<sup>[19]</sup> The recently recorded gas-phase UV/Vis spectra of the perfluorinated bisalkyl peroxides (R<sup>F</sup>O)<sub>2</sub> (R<sup>F</sup> = F<sub>3</sub>C (**2a**), (F<sub>3</sub>C)<sub>3</sub>C (**2b**) and (C<sub>2</sub>F<sub>5</sub>)(CF<sub>3</sub>)<sub>2</sub>C (**2c**)) show that for **2a** the lowest UV transition is below 200 nm, while the bulkier substituted peroxides **2b,c** exhibit weaker red-shifted transitions at 253 and 250 nm, respectively.<sup>[18]</sup>

As described earlier, ferrocene, Fe<sup>II</sup>Cp<sub>2</sub>, is oxidized to ferrocenium, [Fe<sup>III</sup>Cp<sub>2</sub>]<sup>+</sup>, by addition of peroxide **2b** [Eq. (6)].<sup>[18]</sup> An immediate color change of the solid from orange to dark green is observed during the reaction, which is typical for the formation of

a ferrocenium cation. Indeed, the IR spectrum of the solid (Figure S1.11) shows the characteristic vibration modes of the ferrocenium cation.<sup>[44]</sup> For example, the weak ν(CH) mode is blueshifted by 36 cm<sup>-1</sup> to 3121 cm<sup>-1</sup> with respect to ferrocene and the ν(CC) mode assigned at 1421 cm<sup>-1</sup> is also hypsochromically shifted by 14 cm<sup>-1</sup> in comparison to the reactant ferrocene. The bands at 965, 724 and 536 cm<sup>-1</sup> as well as strong absorption bands in the region from 1300 to 1100 cm<sup>-1</sup> are characteristic for the anion, [OC(CF<sub>3</sub>)<sub>3</sub>]<sup>-</sup>.<sup>[45]</sup> Additionally, the APCI mass spectra (Figure S1.12) show a peak at 235 m/z in the negative mode, characteristic for the [OC(CF<sub>3</sub>)<sub>3</sub>]<sup>-</sup> alkoxide anion. In the positive mode an analogous decomposition pathway, compared to that of ferrocene, is observed. These spectra confirm the formation of [FeCp<sub>2</sub>][OC(CF<sub>3</sub>)<sub>3</sub>].

To a sample of [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> (**2b**) elemental fluorine was added to a total pressure of about 1 bar at r.t. in a PFA tube. The tube was then flame-sealed at liquid nitrogen temperatures and after reaching r.t. NMR spectra were recorded (see the Supporting Information, Figure S1.13). The elemental fluorine is detected at a chemical shift of δ = 425 ppm (liquid: 422±1 ppm, gaseous 419±1 ppm)<sup>[46]</sup>. Peroxide **2b** resists elemental fluorine, and its solubility in **2b** is consistent with the low dipole moment as indicated by the dihedral angle Θ of the peroxide unit of 180° in the solid state and the perfluorinated nature of this compound.<sup>[18]</sup> The longitudinal relaxation time T<sub>1</sub> of the dissolved F<sub>2</sub> was determined by an inversion recovery experiment to be T<sub>1</sub> = 13.5 ms (Figure S1.13). This is approximately 300 times larger than T<sub>1</sub> for gaseous fluorine of 0.045 ms in the pressure range from 1 to 2 bar.<sup>[47]</sup> This indicates that the fast relaxation due to the spin-rotation mechanism observed for gaseous fluorine is hindered and proves that the fluorine is indeed dissolved in peroxide **2b**.

## Conclusions

We present a convenient synthesis to highly reactive perfluoro alkyl hypofluorite compounds R<sup>F</sup>OF from the corresponding alcohol and fluorine with excess CsF. Spectroscopic analysis of the hitherto undescribed (C<sub>2</sub>F<sub>5</sub>)(F<sub>3</sub>C)<sub>2</sub>COF with support of quantum-chemical calculations are reported. We provide a new synthetic approach for perfluoro bisalkyl peroxides R<sup>F</sup>OOR<sup>F</sup> by the reaction of hypofluorites with fluorinated silver wool. Furthermore, we also show the inertness of [(F<sub>3</sub>C)<sub>3</sub>CO]<sub>2</sub> towards strong oxidizers such as elemental fluorine. The liquid temperature range from 16 to 99 °C for the unpolar peroxide **2b** together with its inertness towards strong oxidizing halogens demonstrates its potential as a solvent for oxidation and halogenation reactions. By irradiation with UV light perfluoro alkyl peroxides **2** can be activated to generate valuable R<sup>F</sup>O<sup>•</sup> radicals for synthetic applications such as perfluoro alkoxy group transfer reagents.



## FULL PAPER

## Experimental Section

Experiments were carried out at strictly dry and oxygen free conditions in glass tubes using Teflon® valves or in stainless steel vessels. Purchased starting material was used without further purification. NMR spectra of neat liquid substances were recorded on a JEOL 400 MHz ECS or ECZ spectrometer using a capillary filled with [D6]Acetone (<sup>1</sup>H NMR: 2.05 ppm, 400.53 MHz; <sup>13</sup>C NMR: 29.8 ppm, 100.51 MHz) and CFCl<sub>3</sub> (<sup>19</sup>F NMR: 0 ppm, 376.13 MHz) as external standards. The chemical shift and scalar coupling constants were obtained by the program Mestrenova 10.0.<sup>[48]</sup> Raman spectra were measured at liquid nitrogen temperature on a Bruker MultiRAM II spectrometer equipped with a 1064 nm CW DPSS laser and a LN<sub>2</sub> cooled germanium detector at a resolution of 4 cm<sup>-1</sup>. Gas-phase infrared spectra were recorded using a Bruker Vector 22 spectrometer at a resolution of 2 cm<sup>-1</sup>. UV/Vis-spectra of gaseous samples were obtained using a Perkin-Elmer Lambda-900 spectrophotometer. Mass spectra were measured with an Advion expression<sup>+</sup> compact mass spectrometer. The m/z values of the monoisotopic peaks are given. The NMR relaxation time T<sub>1</sub> was determined by the 180°-τ-90° pulse sequence technique. Powder diffraction data were collected on a STOE IPDS II/T instrument at 290 K with Mo K<sub>α</sub> radiation (λ = 0.71073 Å) using a graphite monochromator. Integration was performed with STOE X-Area V1.56, data analysis and Rietveld refinement was performed with X'Pert HighScore Plus V2.2c.

**Safety note:** Extreme caution should be exercised when working with elemental fluorine and hypofluorites. Explosions have been reported<sup>[2,49]</sup> handling these extremely hazardous compounds. Although the described perfluoroalkyl peroxides were found to be insensitive to shock and friction<sup>[18]</sup> according to the U. N. Recommendations on the Transport of Dangerous goods,<sup>[50]</sup> we cannot exclude explosive reactions in mixtures with other substances.

## Acknowledgements

We gratefully acknowledge the Zentraleinrichtung für Datenverarbeitung (ZEDAT) of the Freie Universität Berlin for the allocation of computer time. The authors thank the Graduate School 1582 "Fluorine as a Key Element" (RTN 1582) as well as the CRC 1349 "Fluorine-Specific Interactions: Fundamentals and Function" (project number 387284271) for financial support. We are grateful to the Solvay Fluor GmbH for providing starting materials and for fruitful discussions and the expertise of Holger Pernice.

## Conflict of Interest

The authors do not declare any conflict of interest.

**Keywords:** gas-phase fluorine chemistry • hypofluorites • perfluoro bisalkyl peroxides • vibrational spectroscopy

- [1] K. B. Kellogg, G. H. Cady, *J. Am. Chem. Soc.* **1948**, *70*, 3986-3990.
- [2] J. H. Prager, P. G. Thompson, *J. Am. Chem. Soc.* **1965**, *87*, 230-238.
- [3] P. G. Thompson, J. H. Prager, *Fluoroxy Compounds*, US3442927A.
- [4] M. Lustig, A. R. Pitochelli, J. K. Ruff, *J. Am. Chem. Soc.* **1967**, *89*, 2841-2843.
- [5] W. Storzer, D. D. Desmarteau, *Inorg. Chem.* **1991**, *30*, 4122-4125.
- [6] a) M. Kol, S. Rozen, E. Appelman, *J. Am. Chem. Soc.* **1991**, *113*, 2648-2651; b) E. H. Appelman, D. French, E. Mishani, S. Rozen, *J. Am. Chem. Soc.* **1993**, *115*, 1379-1382.
- [7] J. K. Ruff, A. R. Pitochelli, M. Lustig, *J. Am. Chem. Soc.* **1966**, *88*, 4531-4532.
- [8] G. K. Mulholland, R. E. Ehrenkauf, *J. Org. Chem.* **1986**, *51*, 1482-1489.
- [9] A. Sekiya, D. D. Desmarteau, *Inorg. Chem.* **1980**, *19*, 1328-1330.
- [10] D. H. R. Barton, R. H. Hesse, M. M. Pechet, G. Tarzia, H. T. Toh, N. D. Westcott, *J. Chem. Soc., Chem. Commun.* **1972**, *0*, 122-123.
- [11] S. Rozen **1996**, 1717-1736.
- [12] T. Kashiwaba, H. Oomori, A. Yao, *Dry Etching Agent, Dry Etching Method and Method for Producing Semiconductor Device*, WO 2018/159368.
- [13] R. C. Kennedy, J. B. Levy, *J. Phys. Chem.* **1972**, *76*, 3480-3488.
- [14] a) P. J. Aymonino, *Chem. Commun.* **1965**, 241; b) L. Du, D. D. Desmarteau, V. Tortelli, M. Galimberti, *J. Fluorine Chem.* **2009**, *130*, 830-835.
- [15] K. K. Johri, D. D. DesMarteau, *J. Org. Chem.* **1983**, *48*, 242-250.
- [16] W. Navarrini, V. Tortelli, A. Russo, S. Corti, *J. Fluorine Chem.* **1999**, *95*, 27-39.
- [17] V. Francesco, M. Sansotera, W. Navarrini, *J. Fluorine Chem.* **2013**, *155*, 2-20.
- [18] J. H. Nissen, T. Stüker, T. Drews, S. Steinhauer, H. Beckers, S. Riedel, *Angew. Chem. Int. Ed.* **2019**, *58*, 3584-3588; *Angew. Chem.* **2019**, *131*, 3622-3626.
- [19] M. S. Toy, R. S. Stringham, *J. Fluorine Chem.* **1976**, *7*, 375-383.
- [20] C. J. Marsden, L. S. Bartell, F. P. Diodati, *J. Mol. Struct.* **1977**, *39*, 253-262.
- [21] D. E. Gould, C. T. Ratcliffe, L. R. Anderson, W. B. Fox, *J. Chem. Soc. D* **1970**, 216.
- [22] F. Swarts, *Bull. Soc. Chim. Belg.* **1933**, 102-113.
- [23] R. S. Porter, G. H. Cady, *J. Am. Chem. Soc.* **1957**, *79*, 5628-5631.
- [24] H. L. Roberts, *J. Chem. Soc.* **1964**, 4538-4540.
- [25] W. J. Peláez, G. A. Argüello, *Tetrahedron Letters* **2010**, *51*, 5242-5245.
- [26] S.-L. Yu, D. D. Desmarteau, *Inorg. Chem.* **1978**, *17*, 304-306.
- [27] W. Navarrini, M. Sansotera, P. Metrangolo, P. Cavallotti, G. Resnati, *Modification of carbonaceous materials*, WO 2009/019243 A1.
- [28] D. D. DesMarteau, *Inorg. Chem.* **1970**, *9*, 2179-2181.
- [29] M. E. Redwood, C. J. Willis, *Can. J. Chem.* **1965**, *43*, 1893-1898.
- [30] M. S. Toy, R. S. Stringham, *J. Fluorine Chem.* **1975**, *5*, 25-30.
- [31] I. M. Mills, W. B. Person, J. R. Scherer, B. Crawford, *J. Chem. Phys.* **1958**, *28*, 851-853.
- [32] C. V. Berney, *Spectrochim. Acta* **1965**, *21*, 1809-1823.
- [33] a) I. Mori, T. Tamura, M. Ohashi, *Hypofluorite gas for removal of deposits by solid-gas reaction in cleaning or etching*, EP 924282; b) I. Mori, T. Kawashima, M. Ohashi, T. Tamura, *Method for stripping hyposulfites from flue gases in cleaning or etching of semiconductor devices*, JP 2001321638.
- [34] Woltz, P. J. H., A. H. Nielsen, *J. Chem. Phys.* **1952**, *20*, 307-312.
- [35] R. Ireton, A. S. Gordon, D. C. Tardy, *Int. J. Chem. Kinet.* **1977**, *9*, 769-775.
- [36] M. J. Hopper, J. W. Russell, J. Overend, *J. Chem. Phys.* **1968**, *48*, 3765-3772.
- [37] M. D. Hurley, T. J. Wallington, M. S. Javadi, O. J. Nielsen, *Chem. Phys. Lett.* **2008**, *450*, 263-267.
- [38] M. Wechsberg, G. H. Cady, *J. Am. Chem. Soc.* **1969**, *91*, 4432-4436.

## FULL PAPER

- [39] W. Grochala, Z. Mazej, *Phil. Trans. R. Soc. A* **2015**, 373.
- [40] F. B. Dudley, G. H. Cady, *J. Am. Chem. Soc.* **1957**, 79, 513-514.
- [41] R. Burger, P. Bigler, *J. Magn. Reson.* **1998**, 135, 529-534.
- [42] N. S. Walker, D. D. Desmarteau, *J. Fluorine Chem.* **1975**, 5, 135-139.
- [43] J.-P. Kintzinger, H. Marsmann, *Oxygen-17 and Silicon-29*, Springer Berlin Heidelberg, Berlin, Heidelberg, **1981**.
- [44] I. Pavlík, J. Klikorka, *Coll. Czech. Chem. Comm.* **1965**, 30, 664-674.
- [45] A. Reisinger, N. Trapp, I. Krossing, *Organometallics* **2007**, 26, 2096-2105.
- [46] J. W. Nebgen, W. B. Rose, F. I. Metz, *J. Mol. Spectrosc.* **1966**, 20, 72-74.
- [47] V. R. Celinski, M. Ditter, F. Kraus, F. Fujara, J. Schmedt auf der Gönne, *Chemistry* **2016**, 22, 18388-18393.
- [48] C. Cobas, S. Domínguez, N. Larin, I. Iglesias, C. Geda, F. Seoane, M. Sordo, P. Monje, S. Fraga, R. Cobas et al., *MestReNova*, Mestrelab Research S.L., **2015**.
- [49] C. Lu, J.-H. Kim, D. D. Desmarteau, *J. Fluorine Chem.* **2010**, 131, 17-20.
- [50] *Recommendations on the transport of dangerous goods. Model regulations*, UNITED NATIONS, New York, **2007**.



## FULL PAPER

## FULL PAPER

## Once Upon a Time... Hypofluorites

The convenient synthesis of alkyl hypofluorites RFOF [RF = (F<sub>3</sub>C)<sub>3</sub>C, (C<sub>2</sub>F<sub>5</sub>)(F<sub>3</sub>C)<sub>2</sub>C] is the germ cell for perfluoro bisalkyl peroxides. Hypofluorites are prepared from the corresponding alcohols RFOH and elemental fluorine. The perfluoro bis-tert-butyl peroxide is highly inert towards oxidizers and even dissolves elemental fluorine without decomposition.



J. H. Nissen, T. Drews, B. Schröder, H. Beckers, S. Steinhauer, Prof. S. Riedel\*

**Page No. – Page No.**

Perfluoro Alkyl Hypofluorites and Peroxides Revisited