


Copper-Catalyzed Intramolecular Alkene Carboetherification:
Synthesis of Fused-Ring and Bridged-Ring Tetrahydrofurans

Yan Miller, Lei Miao, Azade S. Hosseini, and Sherry R. Chemler*

Department of Chemistry, The State University of New York at Buffalo, Buffalo, New York 14260, United States

Supporting Information

ABSTRACT: Fused-ring and bridged-ring tetrahydrofuran scaffolds are found in a number of natural products and biologically active compounds. A new copper-catalyzed intramolecular carboetherification of alkenes for the synthesis of bicyclic tetrahydrofurans is reported herein. The reaction involves Cu-catalyzed intramolecular addition of alcohols to unactivated alkenes and subsequent aryl C–H functionalization provides the C–C bond. Mechanistic studies indicate a primary carbon radical intermediate is involved and radical addition to the aryl ring is the likely C–C bond-forming mechanism. Preliminary catalytic enantioselective reactions are promising (up to 75% ee) and provide evidence that copper is involved in the alkene addition step, likely through a *cis*-oxycupration mechanism. Catalytic enantioselective alkene carboetherification reactions are rare and future development of this new method into a highly enantioselective process is promising. During the course of the mechanistic studies a protocol for alkene hydroetherification was also developed.



1. INTRODUCTION

Tetrahydrofurans are frequently found in bioactive compounds.¹ These structurally complex, oftentimes chiral compounds are isolated from natural products, usually in minute quantities, so *de novo* chemical synthesis of such tetrahydrofurans has been actively pursued.² Methods for tetrahydrofuran synthesis include intramolecular S_N2 reactions involving alcohols and leaving groups, [3+2] dipolar cycloadditions, addition of alcohols to electrophile-activated alkenes, addition of nucleophiles to Lewis acid-activated lactol derivatives and hydrogenation of furans.^{2,3} This article reports a new method for the formation of fused-ring and bridged-ring bicyclic tetrahydrofurans via a copper-catalyzed doubly intramolecular alkene carboetherification/C–H functionalization of aryl-substituted 4-pentenols (Scheme 1). Fused-ring and





Figure 1. Bioactive fused-ring and bridged-ring tetrahydrofurans.

Scheme 1. Copper-Facilitated Alkene Carboetherifications



bridged-ring tetrahydrofurans analogous to the ones described in this report can be found in a number of terpenoid natural products including those that possess antibiotic and anti-HIV activity (Figure 1).⁴

Previous syntheses of fused and bridged bicyclic tetrahydrofurans include tetrahydrofuran formation followed by intramolecular electrophilic aromatic substitution,⁵ [4+3]

annulations,⁶ [3+2] annulations,⁷ [5+2] annulations,⁸ ring-expanding reactions⁹ and palladium-catalyzed doubly intramolecular alkene carboetherification/C–H functionalization.¹⁰ While these methods are synthetic advances, most are limited in scope to either fused-ring or bridged-ring tetrahydrofuran synthesis and few are enantioselective. The method reported herein is general for both fused- and bridged-ring synthesis and the potential for development of an efficient catalytic enantioselective protocol is promising (*vide infra*).

The synthesis of tetrahydrofurans via transition metal-catalyzed carboetherification, addition of alcohols and carbon units to alkenes, represents an efficient transformation of two very common functional groups.^{3a,d} Transition metal-catalyzed

Received: April 16, 2012

Published: June 21, 2012

alkene carboetherification has been developed most extensively with palladium catalysts^{10,11} and to a lesser extent with gold catalysts.¹² Similar transformations have also been reported to occur under cobalt catalysis where carbon radical intermediates are thought to be involved.¹³

Few catalytic asymmetric examples of carboetherification reactions of unactivated alkenes have been reported and all of them occur under palladium catalysis.^{3a,14} A common limitation of these methods is the potential for β -hydride elimination which is either built into the method^{14c,d} or avoided by use of 1,1-disubstituted alkene substrates.^{14a} The racemic Pd-catalyzed reactions most analogous to the reactions reported herein also appear to require use of 1,1-disubstituted alkene substrates.¹⁰ Herein is reported the first copper-catalyzed intramolecular alkene carboetherification. Notably, β -hydride elimination does not occur in this reaction. This reaction likely occurs via a novel *cis*-oxycupration and prognosis for its development into a catalytic enantioselective protocol is promising (*vide infra*).

There are few reports of Cu-catalyzed additions of alcohols to electronically unactivated alkenes. To date, such reactions are limited to alkene hydration and the role of Cu in these reactions is primarily as a controlled source of triflic acid [e.g., from Cu(OTf)₂].¹⁵ In the presence of PhI(OAc)₂, Cu(OTf)₂ has also been shown to catalyze the diacetoxylation of alkenes.¹⁶ A Cu-catalyzed intermolecular addition of alkenes and anhydrides that forms γ -lactones (a net carboetherification) has recently been reported, but is quite different with respect to substrates and likely mechanism than the one reported herein.¹⁷ Likewise, Cu-promoted cyclization–acetoxylation of alkenes¹⁸ and Cu-catalyzed oxyalkylation of vinyl arenes¹⁹ have been reported but these reaction involve initial addition of carbon radicals to alkenes.

Our laboratory has reported the copper-catalyzed intramolecular addition of a variety of amines and carbon moieties across alkenes (carboamination) for the synthesis of pyrrolidines and to a lesser extent piperidines.²⁰ Based on the success of these substrates, we turned our attention to 4-pentenols to ascertain if Cu(II) could promote intramolecular addition of an alcohol to an unactivated alkene. Specifically, we designed a doubly intramolecular carboetherification wherein a projected carbon radical intermediate could add to a nearby aryl ring, thereby resulting in a fused-ring or bridged-ring tetrahydrofuran (*vide supra*, Scheme 1).

2. RESULTS AND DISCUSSION

2.1. General Reaction Conditions and Optimization.

The intramolecular carboetherification of 3,3-dibenzyl-4-penten-1-ol (**1a**) was evaluated using a number of Cu(II) salts, ligands and reaction conditions (Table 1). The *cis*-fused tetrahydronaphthofuran **2a** was obtained with high diastereoselectivity and variable yield depending upon the reaction conditions. We first established the feasibility of the reaction by treatment of **1a** with excess of Cu(2-ethylhexanoate)₂ [Cu(eh)₂, 300 mol %] as the copper promoter (Table 1, entry 1). Lowering the amount of Cu(eh)₂ to 150 and 100 mol % using either K₂CO₃ or *n*-BuLi as base led to roughly 50% conversion (Table 1, entries 2 and 3). While this reaction (Table 1, entry 1) was efficient in providing the cyclized adduct **2a**, we sought a catalytic method. Use of MnO₂ (3 equiv) as stoichiometric oxidant and Cu(eh)₂ (20 mol %) as catalyst provided only 33% conversion (Table 1, entry 4). Changing to 2,2'-bipyridine-complexed Cu(OTf)₂ (20 mol %) provided only a trace amount of **2a** (Table 1, entry 6). [Use of Cu(OTf)₂ without

Table 1. Optimization of Reaction Conditions^a

entry	CuX ₂ (mol %)	ligand (mol %)	oxidant ^b	temp (°C)	yield (%) ^b
1	Cu(eh) ₂ (300)			120	>95 ^c
2	Cu(eh) ₂ (150)			120	55 ^c
3 ^d	Cu(eh) ₂ (100)			120	46
4 ^e	Cu(eh) ₂ (20)		MnO ₂	120	33 ^c
5 ^f	Cu(eh) ₂ (300)			120	82 ^c
6 ^e	Cu(OTf) ₂ (20)	Bipy (25)	MnO ₂	120	trace
7 ^e	Cu(OTf) ₂ (20)	3 (25)	MnO ₂	120	98
8 ^e	Cu(OTf) ₂ (20)	3 (25)	MnO ₂	100	98
9 ^e	Cu(OTf) ₂ (20)	3 (25)	MnO ₂	90	66 ^c
10 ^e	Cu(OTf) ₂ (15)	3 (19)	MnO ₂	120	78
11	Cu(OTf) ₂ (20)	3 (25)	O ₂ (1 atm)	100	20 ^c
12 ^{e,f}	Cu(OTf) ₂ (20)	3 (25)	MnO ₂	100	95
13 ^f	Cu(OTf) ₂ (20)	3 (25)	O ₂ (1 atm)	100	20 ^c
14	Cu(eh) ₂ (20)	3 (25)	MnO ₂	100	0 ^c
15 ^g	Cu(OTf) ₂ (20)	3 (25)	MnO ₂	100	80 ^c

^aReactions were run on 0.188 mmol scale of **1a** in PhCF₃ (0.1 M with respect to **1a**) with K₂CO₃ (1 equiv), CuX₂, and the designated ligand in a sealed pressure tube. For reactions using Cu(OTf)₂, the Cu(OTf)₂ was complexed with the ligand in CF₃Ph for 1 h at 60 °C prior to substrate addition. ^bYield of isolated product (>20:1 dr). ^cConversion (%) based on crude ¹H NMR. ^dAlcohol **1a** was treated with *n*-BuLi (1.1 equiv) at 0 °C, the resulting solution was stirred at rt for 15 min, then Cu(eh)₂ was added and the mixture was stirred at rt for 1 h prior to heating at 120 °C for 24 h. ^e300 mol % MnO₂ was used. ^fNo K₂CO₃ was added in this reaction. ^gToluene was used instead of PhCF₃. The Cu catalyst appeared less soluble in this solvent. Cu(eh)₂ = Cu(2-ethylhexanoate)₂, Bipy = 2,2'-bipyridine.

additional ligand also gave trace **2a** (not shown).] Fortunately bis(oxazoline) **3** complexed to Cu(OTf)₂ (20 mol %) provided 98% yield of **2a** using MnO₂ (3 equiv) as oxidant at 120 °C in CF₃Ph (Table 1, entry 7).

The reaction temperature could be lowered to 100 °C without affecting the yield (Table 1, entry 8), but lower temperature and lower catalyst loading gave lower yields (Table 1, entries 9 and 10). Oxygen (1 atm, balloon) was not an effective oxidant (Table 1, entry 11). It is also noteworthy that K₂CO₃ as base is not necessary for reaction to occur but provides a slightly higher yield in the reaction with Cu(eh)₂ (Table 1, entries 1 and 5). K₂CO₃ in the Cu(OTf)₂·bis(oxazoline)-catalyzed reactions does not appear necessary (Table 1, compare entries 8 and 12) but was generally used in subsequent reactions for simplicity. Use of oxygen as oxidant was also attempted under neutral (no K₂CO₃) conditions but again, no turnover occurred (Table 1, entry 13). When Cu(eh)₂ was used in conjunction with ligand **3** under the otherwise optimal reaction conditions (Table 1, entry 8) no conversion to **2a** was observed, indicating the Cu(OTf)₂·bis(oxazoline) catalyst is more active (Table 1, entry 14). Finally, use of the

more common solvent, toluene, was investigated. The desired product was indeed obtained (Table 1, entry 15) but the conversion was lower (80%), possible due to lower solubility of the copper catalyst in this solvent.

2.2. Scope of Fused-Ring Synthesis. Various substituted 3,3-dibenzyl-4-penten-1-ols were submitted to the optimized reaction conditions (Table 1, entry 8). While the electron-rich *p*-Me, *p*-OMe and *p*-SMe substrates provided excellent yield under these conditions (Table 2, entries 1–3), the more electron-deficient *p*-Br, *p*-Cl, *p*-F and *p*-CF₃-substituted substrates required heating to 110 or 120 °C for complete conversion (Table 2, entries 4–7). All of the reactions of the 3,3-disubstituted-4-pentenols were completely selective for the *cis*-fused diastereomer. The cyclization reaction forms two adjacent stereocenters where one is a quaternary carbon.

Regioisomers were formed in cases where substrates had ortho and meta-aryl substituents. The *m*-MeO substrate 4 produced a 1.9:1 mixture of ortho:para substituted regioisomers 5 and 6, respectively (Table 2, entry 8). This regioselectivity is consistent with addition of a carbon radical to an aryl ring (*vide infra*).^{20a,b,f,21}

The *o*-Me and *o*-MeO-substituted substrates 7a and 7b provided regioisomeric mixtures as well, where in products 9a and 9b, the substituent had apparently moved over to the meta position (Table 2, entries 9 and 10). This is consistent with ipso addition of a carbon radical to the aryl ring followed by alkyl migration/rearrangement to provide either product 8 or 9, depending on which group migrates (Scheme 2).^{20f}

Reaction of the mono- γ -substituted pentenol 10a required heating to 120 °C and favored formation of the *trans*-fused diastereomer 11a with modest selectivity under the Cu(OTf)₂·3-catalyzed conditions (Table 2, entry 11). The diastereoselectivity of the reaction could be improved with use of Cu(eh)₂ (20 mol %) as catalyst even though higher temperature (140 °C) was required for good conversion (Table 2, entry 12). Two other monobenzyl substrates provided analogous results (Table 2, entries 13 and 14). 3,3-Diphenyl-4-penten-1-ol 13 provided the indanofuran 14 in 66% yield upon Cu-catalyzed carboetherification (Table 2, entry 15). The monophenyl analog of 13, 3-phenyl-4-penten-1-ol, however, gave modest conversion (64%) of a 2:1 mixture of carboetherification and hydroetherification products even with higher Cu(OTf)₂·3 loading (30 mol %) and longer reaction time (48 h) (see Supporting Information for details). (±)-1-Phenyl and 1-allyl-3,3-dibenzyl-4-penten-1-ols 15 underwent the Cu(OTf)₂·bis(oxazoline) 3-catalyzed reactions, providing moderate diastereoselectivity for the 2,5-*trans*-tetrahydrofuran diastereomers 16 (Table 3, entries 1 and 5). The diastereoselectivity was improved when Cu(2-ethylhexanoate)₂ was used to promote the reactions (Table 3, entries 2 and 6). The diastereoselectivity further improved when the catalytic conditions with Cu(eh)₂ as catalyst were used (Table 3, entries 3, 4, 7–9). Although we have previously observed that choice of Cu(II) ligands can affect diastereoselectivity of related alkene aminooxygenation reactions,²² the oxidant contribution to diastereoselectivity was unexpected. Addition of ligand 3 to the Cu(eh)₂-catalyzed reaction had little effect on the reaction (compare entries 3 and 4, Table 3).


2.3. Bridged-Ring Tetrahydrofuran Synthesis. A transannular carboetherification for the formation of oxabicyclo[3.2.1]octanes was next explored (Table 4). These are rare examples of transannular additions of carbon radicals to

Table 2. Scope of Fused-Ring Tetrahydrofuran Synthesis via Alkene Carboetherification^a

entry	substrate	product(s) (ratio) ^b	yield (%) ^c
1			94
	1b , Ar = 4-MeC ₆ H ₄	2b , X = Me	
2	1c , Ar = 4-MeOC ₆ H ₄	2c , X = OMe	94
3	1d , Ar = 4-MeSC ₆ H ₄	2d , X = SMe	88
4 ^d	1e , Ar = 4-BrC ₆ H ₄	2e , X = Br	88
5 ^e	1f , Ar = 4-ClC ₆ H ₄	2f , X = Cl	85
6 ^e	1g , Ar = 4-FC ₆ H ₄	2g , X = F	72
7 ^d	1h , Ar = 4-CF ₃ C ₆ H ₄	2h , X = CF ₃	90
8	4 , Ar = 3-MeOC ₆ H ₄	 5 6 (ratio = 1.9 : 1)	98
9	7a , Ar = 2-MeOC ₆ H ₄	 8a (X = OMe) 9a (X = OMe) (ratio = 1.3 : 1)	96
10	7b , Ar = 2-MeC ₆ H ₄	8b, 9b , X = Me (ratio = 4.5 : 1)	94
11 ^e	 10a , X = OMe	 11a 12a 11a : 12a = 1.8-3.2 : 1	78
12 ^f	10a	11a : 12a = 5-6.3 : 1	80
13 ^f	10b , X = SMe	11b : 12b = 4-5 : 1	67
14 ^f	10c , X = CH ₃	11c : 12c = 4 : 1	56
15 ^e	 13	 14	66

^aReactions were run at 100 °C in CF₃Ph with Cu(OTf)₂ (20 mol %) complexed with ligand 3 (25 mol %) in the presence of K₂CO₃ (1 equiv), and MnO₂ (300 mol %) for 24 h. ^bRatio determined by analysis of the ¹H NMR spectra of the crude and flash chromatographed product mixtures. ^cIsolated yield after flash chromatography on SiO₂. ^dReaction was run at 110 °C. ^eReaction was run at 120 °C. ^fReaction was catalyzed by Cu(eh)₂ instead of Cu(OTf)₂·3 and was run at 140 °C. All reactions were performed in duplicate.

Scheme 2. Rationale for Regioisomer Formation

Table 3. Reactions of 1-Substituted 3,3-Dibenzyl-4-pentenols^a

entry	substrate	CuX ₂ -ligand	oxidant	yield (%) ^b	dr ^c
1	15a, R = Ph	Cu(OTf) ₂ ·3	MnO ₂	98	2.6:1
2 ^d	15a	Cu(eh) ₂		86	12:1
3	15a	Cu(eh) ₂ ·3	MnO ₂	81	>20:1
4	15a	Cu(eh) ₂	MnO ₂	79	>20:1
5	15b, R = Allyl	Cu(OTf) ₂ ·3	MnO ₂	98	2.5:1
6 ^d	15b	Cu(eh) ₂		87	5.5:1
7	15b	Cu(eh) ₂ ·3	MnO ₂	44	>20:1
8	15b	Cu(eh) ₂	MnO ₂	50	>20:1
9 ^e	15b	Cu(eh) ₂	MnO ₂	95	>20:1

^aSame general reaction conditions as Table 2 unless otherwise noted.^bIsolated yield after flash chromatography on SiO₂. ^cRatio determined by analysis of the ¹H NMR spectra of the crude reaction mixture.^dReaction run with Cu(eh)₂ (300 mol %). ^eReaction run at 120 °C for 48 h.

arenes (*vide infra*). These reactions occurred best at 120 °C using the Cu(OTf)₂-bis(oxazoline) 3-catalyzed reaction conditions. The oxabicyclo[3.2.1]octane **21** formed in entry 6 resembles the natural product bruguirol (Figure 1). We also examined the reactivity of substrates with higher alkene substitution in this carboetherification reaction (Table 4, entries 7–9). 2,2-Diphenyl-4-methyl-4-penten-1-ol **22** underwent the oxidative cyclization to afford the 2-methyloxabicyclo[3.2.1]octane **23** in 87% yield. (*E*)-2,2-diphenyl-4-hexen-1-ol **24a** also underwent the cyclization efficiently, producing adducts **25a** as a 1:1 diastereomeric mixture. This result was surprising as we have previously observed poor reactivity with internal alkene substrates in copper-promoted carboamination reactions.^{20b} Substrate **24b**, with phenyl substitution on the terminal carbon of the alkene, reacted to give a complex mixture that included aldehyde peaks in the crude ¹H NMR (Table 4, entry 9).

2.4. Catalytic Enantioselective Carboetherification.

Development of an enantioselective carboetherification protocol was next explored. The [Cu(*R,R*)-Ph-box](OTf)₂ catalyst was explored due to our previous success in rendering the copper-catalyzed alkene carboamination reaction enantioselective (eq 1).^{20f} Unfortunately 4-pentenols **1a** and **18b** provided racemic carboetherification products using this catalyst (eqs 2 and 3).

We believe the enantioselectivity in the carboamination reaction (eq 1) is due in large part to the steric interaction between the substrate's *N*-tosyl substituent and the phenyl group on the ligand's oxazoline ring (e.g., **TS-A**). Assuming an


Table 4. Synthesis of Bridged Bicyclic Tetrahydrofurans^a

entry	substrate	product(s)	yield (%) ^b
1			93
2	18b , X = OMe	19b , X = OMe	91
3	18c , X = SMe	19c , X = SMe	87
4	18d , X = Cl	19d , X = Cl	93
5	18e , X = F	19e , X = F	92
6 ^c			67
7			87
8			81
9	24b , R = Ph	complex mixture	--

^aReactions were run at 120 °C in CF₃Ph (0.1 M with respect to substrate) with Cu(OTf)₂ (20 mol %) complexed with ligand **3** (25 mol %) in the presence of K₂CO₃ (1 equiv) and MnO₂ (300 mol %) for 12 h unless otherwise noted. ^bIsolated yield after flash chromatography on SiO₂. ^c18% of starting alcohol **20** was recovered.

analogous cyclic *cis*-oxycupration transition state (e.g., **TS-B**) in the carboetherification reaction, this steric interaction is not present, which could contribute to loss of enantioselectivity (the enantiomeric **TS-C** may not be energetically very different from **TS-B**).

We reasoned that increasing the steric demands of the substrate near the alcohol region could render a cyclic oxycupration step more enantioselective by increasing steric interactions between the substrate and ligand. Thus, substrates with mono and disubstitution at the hydroxyl-bearing carbon were examined (Table 5). While 1-phenyl-3,3-dibenzyl-4-pentenol **15a** gave high diastereoselectivity for **16** with Cu(eh)₂·(*R,R*)-Ph-box as catalyst, the enantioselectivity was <5%. Changing to the [Cu(*R,R*)-Ph-box](OTf)₂ catalyst, we obtained a 2:1 diastereomeric ratio of **16a** to **17a** where **16a** was formed in 30% ee and **17a** was formed in 67% ee. This reaction is likely a kinetic resolution of the racemic alcohol substrate (absolute configurations tentatively assigned as shown, *vide infra*). The analogous 1-phenyl-3,3-di(*p*-chlorobenzyl)-4-pentenol **15c** reacted with similar selectivity (Table 5, entry 3). The 1,1-diphenyl-3,3-dibenzyl-4-pentenol **15d** provided one carboetherification diastereomer in ca. 75% ee (conservative estimate based on incompletely resolved enantiomers) and its 1,1-diphenyl-3,3-di(*p*-chlorobenzyl)-4-

Table 5. Catalytic Asymmetric Carboetherifications^a


entry	substrate	CuX ₂	yield (%)	dr (16:17) ^b	ee (%) ^c
1	15a, R ¹ = Ph, R ² = H, Ar = Ph	Cu(eh) ₂	62	>20:1	<5
2	15a	Cu(OTf) ₂	89	2:1	30, 67
3	15c, R ¹ = Ph, R ² = H, Ar = 4-Cl-C ₆ H ₄	Cu(OTf) ₂	77	2:1	32, 60
4	15d, R ¹ = Ph, R ² = Ph, Ar = Ph	Cu(OTf) ₂	89	~75	
5	15e, R ¹ = Ph, R ² = Ph, Ar = 4-Cl-C ₆ H ₄	Cu(OTf) ₂	85	nd	

^aReaction conditions: CuX₂ (0.376 mmol) and (R,R)-Ph-box (0.0470 mmol) were combined in PhCF₃ (0.6 mL) and heated in a sealed tube for 2 h. Alcohol **15** (0.188 mmol) in 0.9 mL of PhCF₃, K₂CO₃ (0.188 mmol), and MnO₂ (0.564 mmol) were added, and the reaction was stirred at 100 °C in a sealed tube for 24 h. ^bSubstrates are racemic or achiral. ^cIsolated yield after flash chromatography on SiO₂. ^dRatio determined by analysis of the ¹H NMR spectra of the crude reaction mixture. ^eEnantiomeric excess determined by chiral HPLC. nd = not determined. Enantiomers could not be separated on chiral HPLC or GC.

pentenol derivative (**15e**) gave a product with a comparable optical rotation (see Supporting Information), but the enantiomers could not be separated for %ee determination. The data in Table 5 clearly indicate that development of a catalytic enantioselective copper-catalyzed alkene carboetherification is feasible. Mechanistically, the enantioselective reactions provide strong support for the presence of Cu in the alkene addition step.

2.5. Absolute Stereochemistry Assignment. X-ray quality crystals of enantiomerically enriched heavy atom (Cl)-containing carboetherification products could not be obtained,


unfortunately. We have tentatively assigned the absolute stereochemistry of the products as shown in Table 5 and eqs 4 and 5 (*vide infra*) by analogy to the enantioselective



carboamination products (e.g., eq 1 and see optical rotation correlation discussion in the Supporting Information for **21**, *vide infra*).^{20f}

2.6. Mechanistic Probes. We further explored the enantioselective variant of the transannular carboetherification of 4-pentenol **20**. Although the reaction did not go to completion at 100 °C, the bicyclic furan **21** was obtained in 50% ee (Scheme 3). We next introduced 1,4-cyclohexadiene to

Scheme 3. Enantioselective Transannular Carboetherification and Hydroetherification




the reaction and obtained tetrahydrofuran **26** in 85% yield (along with 10% unreacted alcohol **20**, Scheme 3). Tetrahydrofuran **26** was obtained in 76% enantiomeric excess. Taken together these reactions indicate (1) a carbon radical, one that can perform H-atom abstraction, is a likely intermediate in the reaction, (2) the H-atom abstraction is faster than transannular C–C bond formation and (3) the carboetherification reaction may suffer some ring-opening/unselective ring-closing prior to C–C bond formation which could account for its diminished enantioselectivity relative to the apparently less reversible alkene hydroetherification involving H-atom abstraction. One potential reversibility scenario is depicted in Scheme 3. This

reversibility may be less prominent when addition to a more proximal aryl ring, e.g. as in Table 5, occurs. Formation of the favored *cis*-fused products, e.g. **16** and **17**, may be faster than furan ring-opening. In the case of α -substituted substrates **15a** and **15c–e**, selectivity can be as high as 75% ee, which is comparable to the hydroetherification reaction (Scheme 3).

The intermediacy of an sp^2 -hybridized carbon intermediate (e.g., a carbon radical, see Schemes 1–3, was further probed by the reaction of the isotopically labeled 3-*p*-methoxybenzyl-4-penten-1-ol (**10a-D**). Carboetherification provided a 1.8:1 ratio of *trans*-fused **11a-D** and *cis*-fused **12a-D**, both of which were composed of a 1:1 diastereomeric mixture at the deuterated carbon (eq 4). The presence of the carbon radical was further confirmed by running the reaction in the presence of 1,4-cyclohexadiene (eq 5). This hydroetherification reaction was highly diastereoselective and produced the 2,3-*trans*-tetrahydrofuran **27** in 82% yield. No carboetherification products were formed, indicating the H-atom transfer process is faster. This relative rate data, in conjunction with the much higher *trans*-diastereoselectivity for the hydroetherification, seems to indicate again that addition to the alkene is less reversible in the hydroetherification reaction. The 2,3-*trans* kinetic selectivity can be rationalized by **TS-D**, which places the allylic substituent in a pseudo-equatorial position in the cyclization transition state. The 2,3-*trans* diastereomer should experience less sterics in the H-atom abstraction step than its corresponding 2,3-*cis* diastereomer, which can contribute to a highly *trans*-selective hydroetherification (eq 5). Conversely, the C–C bond-forming step of the corresponding carboetherification reaction (radical addition to arene after O–C bond formation) in eq 4 should be more facile for the minor 2,3-*cis* carbon radical intermediate that leads to formation of the *cis*-fused diastereomer **12a-D**, than for the major 2,3-*trans* carbon radical to form the *trans*-fused diastereomer **11a-D**. The preference for *cis*-fused cyclization is clear in the highly selective conversion of **1** to **2** (*vide supra*). Thus, the 2,3-*trans* carbon radical or organocopper intermediate could suffer ring-opening and ring-closing prior to C–C bond formation, resulting in a lower overall reaction diastereoselectivity.

Based on the labeling and H-atom abstraction reactions, the intermediacy of a primary carbon radical is reasonable. The C–C bond-forming process in the aryl addition step of the carboetherification could, however, still feasibly take place either by a radical addition or an electrophilic aromatic substitution mechanism. Aryl addition to an organocopper(III) intermediate, either directly at carbon or via addition to copper(III) followed by reductive elimination, could feasibly lead to the carboetherification product. Analogous mechanisms have been proposed for Pd- and Au-catalyzed reactions.²³ (Carbon–carbon bond formation through a Friedel–Crafts-type addition to a primary carbocation intermediate is less likely as such an intermediate is expected to be very high energy and unlikely to be formed under the reaction conditions.)

The stereochemistry of the deuterio-alkene cannot be used in our analysis since stereochemical information is lost due to the presence of the carbon radical intermediate. Regiochemical trends in electrophilic and radical aryl substitution reactions are known, however, and can be used to infer the likely reaction mechanism. It is known that radicals favor addition ortho to substituents on aromatic rings, possibly due to electronic stabilization of aryl radical intermediates (e.g., eq 6).²¹ Conversely, electrophilic addition of metals, electrophilic organometallics or activated carbonyls (Friedel–Crafts acyla-




tion) to aromatic rings generally occurs at the less sterically hindered para-position (e.g., eq 7).²³

Our reaction provides a regioisomeric mixture where the ortho-substituted product is formed as the major product (Table 2, entry 8, ortho:para = 1.9:1). This regioselectivity is most consistent with a carbon radical addition to the aromatic ring.

2.7. Proposed Mechanism. A proposed catalytic cycle is illustrated in Scheme 4. A *cis*-oxycupration transition state (**TS-E**)

Scheme 4. Proposed Carboetherification Catalytic Cycle



E) is proposed in analogy to related aminocuprations.²⁰ For α -substituted pentenols **15**, the R group is placed in an equatorial position in the cyclic transition state (**TS-E**) resulting in formation of the major 2,5-*trans*-tetrahydrofuran diastereomer **16**. The electronic effect observed in the substrate reactivity in Table 1 seems to indicate that more electron-rich alkenes and/or alcohols are more reactive. Homolysis of the unstable C–Cu(II) bond of intermediate **28** results in the primary carbon radical intermediate **29**. Addition of the resulting carbon radical to the nearest (*cis*) aryl ring produces the new C–C bond in aryl radical **30**, and aryl oxidation provides the product **16**. Oxidation of Cu(I) with MnO_2 regenerates the Cu(II) catalyst.

We have observed that the reaction diastereoselectivity can vary depending upon the nature of the copper(II) ligands, the presence or absence of MnO_2 and the presence or absence of 1,4-cyclohexadiene (Tables 2 and 3 and eqs 4 and 5). Reaction reversibility due to a ring-opening and unselective ring-closing process should be less prevalent if the subsequent step, reaction

termination by H atom abstraction or addition of the carbon radical to the aromatic ring, is rapid. The Cu(eh)₂ catalyst and the presence of MnO₂ increase reaction diastereoselectivity which may indicate these reactions are also less reversible. This is possibly due to an increase in stability of the primary carbon radical intermediate by its reversible conversion to an organocopper(III) species. This species can be formed by direct oxidation of the organocopper(II) species (e.g., **28**) or addition of the primary carbon radical (e.g., **29**) to Cu(II). Both processes should be increased by greater concentration of oxidizing species or greater propensity of the Cu(I) or Cu(II) species to undergo oxidation. A ligand affect on copper oxidation potential is feasible.²⁴

The extent of the reversibility of the tetrahydrofuran ring formation is also highly dependent upon the rate of the subsequent C–C or C–H bond-forming step. As we observed in our enantioselective reactions, 1-substituted-4-pentenol substrates that are able to form *cis*-fused tetrahydronaphthofuran products, e.g. **16**, and reactions involving C–H bond formation via H-atom abstraction (e.g., formation of **26** in Scheme 3) give the highest enantioselectivities (up to 76%). It can be inferred with these substrates and reactions that the oxycupration step is enantioselective and suffers little if any ring-opening/unselective ring-closing.

Carbon–oxygen bond formation via *trans*-oxycupration is a mechanistic alternative to the proposed *cis*-oxycupration shown in Scheme 4. The product stereochemistry cannot be used to differentiate the two mechanisms since stereochemical information from the C–Cu bond formation is lost (*vide supra*, eq 4). The *cis*-oxycupration mechanism requires coordination of the alcohol to the copper center prior to alkene addition. This seems likely due to the high affinity the hard alcohol nucleophile should have for the hard Cu(II) catalyst. The olefin, in comparison, is a softer nucleophile.

An alternative, oxygen radical alkene addition mechanism (from homolysis of an O–Cu bond) can be ruled out since we have observed that the ligand on Cu(II) can influence the enantioselectivity of the reaction. A mechanism involving internal charge transfer of the RO–Cu(II) intermediate to give an RO–Cu(I) complex where both the metal and alkoxyradical participate in the olefin addition step, in analogy to that proposed for related Co-catalyzed reactions,¹³ cannot be ruled out, but seems unlikely since such N to Cu charge transfer does not appear to occur in the analogous aminocupration reaction.²⁵

3. CONCLUSIONS

In conclusion, the first copper-catalyzed intramolecular carboetherification of 4-pentenols has been described. This reaction efficiently provides fused and bridged-ring tetrahydrofurans from common and unactivated alkenols. Evidence for oxycupration as the alkene addition mechanism is presented, notably in the form of ligand-based asymmetric catalysis. The C–C bond-forming step in the carboetherification reaction is thought to occur via carbon radical arene addition and this step constitutes an efficient C–H functionalization. A novel, efficient, diastereoselective and enantioselective hydroetherification reaction, the result of trapping the carbon radical intermediate with 1,4-cyclohexadiene in an intermolecular H-atom abstraction step, was also presented. This hydroetherification reaction could prove useful in generating 5-methyl-2,2-disubstituted tetrahydrofurans currently of interest to the perfume industry.²⁶ The novel reactivity reported herein

will serve as the basis for the development of other alkene oxidative difunctionalization reactions. We anticipate we can broaden the scope of tetrahydrofurans formed through carboetherification by introducing external alkene radical acceptors^{20g} to enable intermolecular C–C bond formation. Application of this method in target-oriented synthesis, optimization of the enantioselective protocol, and additional product analysis for definitive absolute configuration determination are underway and will be reported in due course.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental procedures, spectroscopic data, and crystallographic data for **11a** and **16e** (CIFs). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

schemler@buffalo.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The National Institute of General Medical Science (GM078383) is gratefully acknowledged for support of this research. We thank William W. Brennessel and the X-ray Crystallographic Facility at the University of Rochester for the X-ray structures of **11a** and **16e**. We thank Barbara Casavant, Shulendu Kariakarte, and Timothy Liwosz for kind assistance in the synthesis of five substrates.

■ REFERENCES

- (1) Reviews: (a) Boivin, T. L. B. *Tetrahedron* **1987**, *43*, 3309–3362. (b) Cardillo, G.; Orena, M. *Tetrahedron* **1990**, *46*, 3321–3408. (c) Harmange, J.-C.; Figadere, B. *Tetrahedron: Asymmetry* **1993**, *4*, 1711–1754. (e) Miura, K.; Hosomi, A. *Synlett* **2003**, 143–155.
- (2) (a) Koert, U. *Synthesis* **1995**, 115–132. (b) Bates, R. H.; Chen, M.; Roush, W. R. *Curr. Opin. Drug Disc. Dev.* **2008**, *11*, 778–792. (c) Kuwano, R. *Heterocycles* **2008**, *76*, 909–922. (d) Wolfe, J. P.; Hay, M. B. *Tetrahedron* **2007**, *63*, 261–290. (e) Li, N.; Shi, Z.; Tang, Y.; Chen, J.; Li, X. *Beilstein J. Org. Chem.* **2008**, *4* (48), doi: 10.3762/bjoc.4.48.
- (3) (a) McDonald, R. I.; Liu, G.; Stahl, S. S. *Chem. Rev.* **2011**, *111*, 2981–3019. (b) Piccialli, V. *Synthesis* **2007**, 2585–2607. (c) Majumdar, K. C.; Debnath, P.; Roy, B. *Heterocycles* **2009**, *78*, 2661–2728. (d) Wolfe, J. P. *Synlett* **2008**, 19, 2913–2937. Selected electrophile-activated additions/cyclization of ROH and alkenes: (e) Kang, S. H.; Lee, S. B.; Park, C. M. *J. Am. Chem. Soc.* **2003**, *125*, 15748–15749. (f) Zhou, L.; Tan, C. K.; Jiang, X.; Chen, F.; Yeung, Y.-Y. *J. Am. Chem. Soc.* **2010**, *132*, 15474–15476. (g) Chen, G.; Ma, S. *Angew. Chem., Int. Ed.* **2010**, *49*, 8306–8308. (h) Veitch, G. E.; Jacobsen, E. N. *Angew. Chem., Int. Ed.* **2010**, *49*, 7332–7335. (i) Murai, K.; Matsushita, K. T.; Nakamura, A.; Fukushima, S.; Shimura, M.; Fujioka, H. *Angew. Chem., Int. Ed.* **2010**, *49*, 9174–9177. (j) Whitehead, D. C.; Yousefi, R.; Jaganathan, A.; Borhan, B. *J. Am. Chem. Soc.* **2010**, *132*, 3298–3299. (k) Yousefi, R.; Whitehead, D. C.; Mueller, J. M.; Staples, R. J.; Borhan, B. *Org. Lett.* **2011**, *13*, 608–611. (l) Hennecke, U.; Muller, C. H.; Frohlich, R. *Org. Lett.* **2011**, *13*, 860–863. (m) Denmark, S. E.; Burk, M. T. *Org. Lett.* **2012**, *14*, 256–259.
- (4) (a) Yoshikawa, K.; Kokudo, N.; Tanaka, M.; Nakano, T.; Shibata, H.; Aragaki, N.; Higuchi, T.; Hashimoto, T. *Chem. Pharm. Bull.* **2008**, *56*, 89–92. (b) Yao, S.; Tang, C. P.; Ke, C. Q.; Ye, Y. *J. Nat. Prod.* **2008**, *71*, 1242–1246. (c) Peranza-Sanchez, S. R.; Chavez, D.; Chai, H.-B.; Shin, Y. G.; Garcia, R.; Mejia, M.; Fairchild, C. R.; Lane, K. E.; Menendez, A. T.; Farnsworth, N. R.; Cordell, G. A.; Pezzuto, J. M.

- Kinghorn, A. D. *J. Nat. Prod.* **2000**, 63, 492–495. (d) Xiao, W.-L.; Yang, L.-M.; Gong, N.-B.; Wu, L.; Wang, R.-R.; Pu, J.-X.; Li, X.-L.; Huang, S.-X.; Zheng, Y.-T.; Li, R.-T.; Lu, Y.; Zheng, Q.-T.; Sun, H.-D. *Org. Lett.* **2006**, 8, 991–994. (e) Han, L.; Huang, X.; Sattler, I.; Moellmann, U.; Fu, H.; Lin, W.; Grabley, S. *Planta Med.* **2005**, 71, 160. (f) Wang, J.; Soisson, S. M.; Young, K.; Shoop, W.; Kodali, S.; Galgocsi, A.; Painter, R.; Parthasarathy, G.; Tang, Y. S.; Cummings, R.; Ha, S.; Dorso, K.; Motyl, M.; Jayasuriya, H.; Ondeyka, J.; Herath, K.; Zhang, C.; Hernandez, L.; Allocco, J.; Basilio, A.; Tormo, J. R.; Genilloud, O.; Vincente, F.; Pelaez, F.; Colwell, L.; Lee, S. H.; Michael, B.; Felcetto, T.; Gill, C.; Silver, L. L.; Hermes, J. D.; Bartizal, K.; Barrett, J.; Schmatz, D.; Becker, J. W.; Cully, D.; Singh, S. B. *Nature* **2006**, 441, 358–361.
- (5) (a) Solorio, D. M.; Jennings, M. P. *J. Org. Chem.* **2007**, 72, 6621–6623. (b) Fananas, F. J.; Fernandez, A.; Cevic, D.; Rodriguez, F. J. *Org. Chem.* **2009**, 74, 932–934.
- (6) Molander, G. A.; Cameron, K. O. *J. Am. Chem. Soc.* **1993**, 115, 830–846.
- (7) (a) Padwa, A.; Hornbuckle, S. F.; Fryxell, G. E.; Stull, P. D. *J. Org. Chem.* **1989**, 54, 817–824. (b) Hodgson, D. M.; Labande, A. H.; Pierard, F. Y. T. M.; Castro, M. A. E. *J. Org. Chem.* **2003**, 68, 6153–6159. (c) Hodgson, D. M.; Stupple, P. A.; Pierard, F. Y. T. M.; Labande, A. H.; Johnstone, C. *Chem.—Eur. J.* **2001**, 7, 4465–4476. (d) Ishida, K.; Kusama, H.; Iwasawa, N. *J. Am. Chem. Soc.* **2010**, 132, 8842–8843.
- (8) Rodriguez, J. R.; Rumbo, A.; Castedo, L.; Mascarenas, J. L. *J. Org. Chem.* **1999**, 64, 4560–4563.
- (9) (a) McGrath, N. A.; Bartlett, E. S.; Sittihan, S.; Njardarson, J. T. *Angew. Chem., Int. Ed.* **2009**, 48, 8543–8546. (b) Loy, R. N.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2009**, 131, 2786–2787.
- (10) (a) Matuura, B. S.; Condie, A. G.; Buff, R. C.; Karahalas, G. J.; Stephenson, C. R. J. *Org. Lett.* **2011**, 13, 6320–6323. (b) Zhu, R.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2012**, 51, 1926–1929.
- (11) Representative Pd-catalyzed alkene carboetherifications: (a) Semmelhack, M. F.; Bodurow, C. *J. Am. Chem. Soc.* **1984**, 106, 1496–1498. (b) Wolfe, J. P.; Rossi, M. A. *J. Am. Chem. Soc.* **2004**, 126, 1620–1621. (c) Hay, M. B.; Wolfe, J. P. *J. Am. Chem. Soc.* **2005**, 127, 16468–16476. (d) Ward, A. F.; Wolfe, J. P. *Org. Lett.* **2009**, 11, 2209–2212. (e) Nakhla, J. S.; Kampf, J. W.; Wolfe, J. P. *J. Am. Chem. Soc.* **2006**, 128, 2893. (f) Nicolai, S.; Erard, S.; Fernandez-Gonzales, D.; Waser, J. *Org. Lett.* **2010**, 12, 384–387. (g) Jiang, D.; Peng, J.; Chen, Y. *Org. Lett.* **2008**, 10, 1689–1698. (h) Hayashi, S.; Yorimitsu, H.; Oshima, K. *J. Am. Chem. Soc.* **2009**, 131, 2052–2053. (i) Han, X.; Lu, X. *Org. Lett.* **2009**, 11, 2381–2384. (j) Coy, B., E. D.; Jovanovic, L.; Sefkow, M. *Org. Lett.* **2010**, 12, 1976–1979. (k) Satterfield, A. D.; Kubota, A.; Sanford, M. S. *Org. Lett.* **2011**, 13, 1076–1079.
- (12) Au-catalyzed alkene carboetherifications: (a) Zhang, L.; Kozmin, S. A. *J. Am. Chem. Soc.* **2005**, 127, 6962–6963. (b) Sherry, B. D.; Maus, L.; Laforteza, B. N.; Toste, F. D. *J. Am. Chem. Soc.* **2006**, 128, 8132–8133. (c) Zhang, G.; Cui, L.; Wang, Y.; Zhang, L. *J. Am. Chem. Soc.* **2010**, 132, 1474–1475. (d) Melhado, A. D.; Brenzovich, W. E.; Lackner, A. D.; Toste, F. D. *J. Am. Chem. Soc.* **2010**, 132, 8885–8887. (e) Ball, L. T.; Green, M.; Lloyd-Jones, G. C.; Russell, C. A. *Org. Lett.* **2010**, 12, 4724–4727. (f) Brenzovich, W. E., Jr.; Brazeau, J.-F.; Toste, F. D. *Org. Lett.* **2010**, 12, 4728–4731.
- (13) Cobalt-catalyzed 4-pentenol cyclizations: (a) Schuch, D.; Fries, P.; Donges, M.; Menendez Perez, B.; Hartung, J. *J. Am. Chem. Soc.* **2009**, 131, 12918–12920. (b) Inoki, S.; Mukaiyama, T. *Chem. Lett.* **1990**, 67–70. (c) Fries, P.; Halter, D.; Kleinschek, A.; Hartung, J. *J. Am. Chem. Soc.* **2011**, 133, 3906–3912. (d) Palmer, C.; Morra, N. A.; Stevens, A. C.; Bajtos, B.; Machin, B. P.; Pagenkopf, B. L. *Org. Lett.* **2009**, 11, 5614–5617.
- (14) Catalytic enantioselective alkene carboetherification-type reactions: (a) Tietze, L. F.; Sommer, K. M.; Zinngrebe, J.; Stecker, F. *Angew. Chem., Int. Ed.* **2005**, 44, 257–259. (b) Pathak, T. P.; Gligorich, K. M.; Welm, B. E.; Sigman, M. S. *J. Am. Chem. Soc.* **2010**, 132, 7870–7871. (c) Arai, M. A.; Kuraishi, M.; Arai, T.; Sasai, H. *J. Am. Chem. Soc.* **2001**, 123, 2907. (d) Kawamura, Y.; Kawano, Y.; Matsuda, T.; Ishitobi, Y.; Hosokawa, T. *J. Org. Chem.* **2009**, 74, 3048.
- Selected other catalytic enantioselective additions of alcohols to alkenes to form tetrahydrofurans: (e) Burke, S. D.; Jiang, L. *Org. Lett.* **2001**, 3, 1953–1955.
- (15) (a) Taylor, J. G.; Whittall, N.; Hii, K. K. M. *Chem. Commun.* **2005**, 5103–5105. (b) Tschan, M. J.-L.; Thomas, C. M.; Strub, H.; Carpentier, J.-F. *Adv. Synth. Catal.* **2009**, 351, 2496–2504. (c) Adrio, L. A.; Quek, L. S.; Taylor, J. G.; Hii, K. K. M. *Tetrahedron* **2009**, 65, 10334–10338.
- (16) (a) Seayad, J.; Seayad, A. M.; Chai, C. L. L. *Org. Lett.* **2010**, 12, 1412–1415. (b) Kang, Y.-B.; Gade, L. H. *J. Am. Chem. Soc.* **2011**, 133, 3658–3667.
- (17) Huang, L.; Jiang, H.; Qi, C.; Liu, X. *J. Am. Chem. Soc.* **2010**, 132, 17652–17654.
- (18) Wang, Y.; Du, H. *J. Org. Chem.* **2010**, 75, 3503–3506.
- (19) (a) Cheng, K.; Huang, L.; Zhang, Y. *Org. Lett.* **2009**, 11, 2908–2911. An Fe-catalyzed radical carboetherification: Liu, W.; Li, Y.; Liu, K.; Li, Z. *J. Am. Chem. Soc.* **2011**, 133, 10756–10759.
- (20) Cu-promoted carboamination: (a) Sherman, E. S.; Chemler, S. R.; Tan, T. B.; Gerlits, O. *Org. Lett.* **2004**, 6, 1573–1575. (b) Sherman, E. S.; Fuller, P. H.; Kasi, D.; Chemler, S. R. *J. Org. Chem.* **2007**, 72, 3896–3905. (c) Fuller, P. H.; Chemler, S. R. *Org. Lett.* **2007**, 9, 5477–5480. Cu-catalyzed carboamination: (d) Zeng, W.; Chemler, S. R. *J. Am. Chem. Soc.* **2007**, 129, 12948–12949. (e) Sherman, E. S.; Chemler, S. R. *Adv. Synth. Catal.* **2009**, 351, 467–471. (f) Miao, L.; Haque, I.; Manzoni, M. R.; Tham, W. S.; Chemler, S. R. *Org. Lett.* **2010**, 12, 4739–4741. (g) Liwosz, T. W.; Chemler, S. R. *J. Am. Chem. Soc.* **2012**, 134, 2020–2023.
- (21) Addition of radicals to arenes: (a) Ito, R.; Migita, T.; Morikawa, N.; Simamura, O. *Tetrahedron* **1965**, 21, 955. (b) Pryor, W. A.; Davis, W. H., Jr.; Gleaton, J. H. *J. Org. Chem.* **1975**, 40, 2099–2102. (c) Conrad, J. C.; Kong, J.; Laforteza, B. N.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2009**, 131, 11640–11641.
- (22) Paderes, M. C.; Chemler, S. R. *Eur. J. Org. Chem.* **2011**, 3679–3684.
- (23) (a) Yip, K.-T.; Yang, D. *Org. Lett.* **2011**, 13, 2134–2137. (b) Sibbald, P. A.; Rosewall, C. F.; Swartz, R. D.; Michael, F. E. *J. Am. Chem. Soc.* **2009**, 131, 15945–15951. (c) de Haro, T.; Nevado, C. *Angew. Chem., Int. Ed.* **2011**, 50, 906–910. (d) Chiche, B.; Finiels, A.; Gauthier, C.; Geneste, P. *J. Org. Chem.* **1986**, 51, 2128–2130.
- (24) Jones, G. O.; Liu, P.; Houk, K. N.; Buchwald, S. L. *J. Am. Chem. Soc.* **2010**, 132, 6205–6213.
- (25) Paderes, M. C.; Belding, L.; Fanovic, B.; Dudding, T.; Keister, J. B.; Chemler, S. R. *Chem.—Eur. J.* **2012**, 18, 1711–1726.
- (26) Kraft, P.; Givaudan, S. A. Trisubstituted furans suitable for the preparation of fragrance compositions. U.S. patent 7,632,790, Dec 15, 2009.
- (27) Fontiadou, A. D.; Zografos, A. L. *Org. Lett.* **2011**, 13, 4592–4595.

NOTE ADDED AFTER ASAP PUBLICATION

It has come to the authors' attention that a copper-promoted carboetherification reaction involving endo cyclization of a phenol onto a pendant alkene was reported recently. This work is now cited as ref 27, and this paper reposted July 11, 2012.