

Efficient and Selective Multicomponent Oxidative Coupling of Two Different Aliphatic Primary Amines into Thioamides by Elemental Sulfur


Thanh Binh Nguyen,* Ludmila Ermolenko, and Ali Al-Mourabit

Centre de Recherche de Gif-sur-Yvette, Institut de Chimie des Substances Naturelles
CNRS, 91198 Gif-sur-Yvette Cedex, France

nguyen@icsn.cnrs-gif.fr

Received July 23, 2012

ABSTRACT



An efficient and selective multicomponent oxidative coupling of two different aliphatic primary amines into thioamides by elemental sulfur under solvent-free conditions has been developed.

Simple and effective reactions including multicomponent coupling processes using readily available reagents are seen as a key solution for the 21st century pollution problems generated by large scale chemistry. With this vision, the use of elemental sulfur in organic syntheses appears to be highly desirable to maximize atom economy and to avoid expensive complex catalysts. For this purpose, we have selected the study of the synthesis of thioamides by a coupling reaction between two amines. Molecules containing a thioamide¹ moiety play an important role in chemistry. Thioamide functions are also present in a variety of biologically active molecules.² They are also known as building blocks in the synthesis of

heterocycles and other compounds containing both nitrogen and sulfur within their backbones.³

Different synthetic methods have been discovered for the synthesis of thioamides. Among these strategies, thioanation of amide analogues with Lawesson's reagent is the most common,⁴ but this reaction cannot be classified as an atom economical approach because of crucial limitations: only one oxygen atom is replaced by a sulfur atom, and no other new bond was created. Thus, it is worthwhile to provide a practical and environmentally benign method to synthesize thioamides. Because sulfur is nontoxic, stable under ambient conditions, easy to handle, and readily available, its use in the preparation of thioamides is highly desirable as exemplified by the Willgerodt–Kindler reaction, starting from aryl alkyl ketones, elemental sulfur, and secondary amines such as morpholine.⁵

(1) (a) Hurd, R. N.; Delamater, G. *Chem. Rev.* **1961**, *61*, 45. (b) Bauer, W.; Kühlein, K. *Houben-Weyl's Methoden der Organischen Chemie*; Georg Thieme Verlag: Stuttgart, New York, 1985; Vol. E5, pp 1218–1279. (c) Jagodziński, T. S. *Chem. Rev.* **2003**, *103*, 197. (d) Leung, P. H.; Qin, Y.; He, G.; Mok, K. F.; Vittal, J. J. *J. Chem. Soc., Dalton Trans.* **2001**, 309. (e) Murai, T.; Sano, H.; Kawai, H.; Aso, H.; Shibahara, F. *J. Org. Chem.* **2005**, *70*, 8148.

(2) (a) Mehanna, A. S.; Belani, J. D.; Kelley, C. J.; Pallans, L. A. *J. Med. Chem.* **2007**, *50*, 513. (b) Yu, K. L.; Torri, A. F.; Luo, G.; Cianci, C.; Grant-Young, K.; Danetz, S.; Tiley, L.; Krystalb, M.; Meanwella, N. A. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3379.

(3) For selected recent examples, see: (a) Lo, W. S.; Hu, W. P.; Lo, H. P.; Chen, C. T.; Kao, C. L.; Vandavasi, J. K.; Wang, J. J. *Org. Lett.* **2010**, *12*, 5570. (b) Iwata, M.; Yazaki, R.; Chen, I. H.; Sureshkumar, D.; Kumagai, N.; Shibasaki, M. *J. Am. Chem. Soc.* **2011**, *133*, 5554. (c) Goossen, L. J.; Blanchot, M.; Salih, K. S. M.; Karch, R.; Rivas-Nass, A. *Org. Lett.* **2008**, *10*, 4497.


(4) (a) Cava, M. P.; Levinson, M. I. *Tetrahedron* **1985**, *41*, 5061. (b) Brillon, D. *Sulfur Rep.* **1992**, *12*, 297.

(5) (a) Willgerodt, C. *Ber. Dtsch. Chem. Ges.* **1888**, *21*, 534. (b) Kindler, K. *Liebigs Ann. Chem.* **1923**, *431*, 187. (c) Wegler, R.; Kuhle, E.; Schafer, W. *Angew. Chem.* **1958**, *70*, 351. For recent examples of thioamide synthesis by reaction of sulfur with imines or aldehydes and amines, see: (d) Zbruyev, O. I.; Stiasni, N.; Kappe, C. O. *J. Comb. Chem.* **2003**, *5*, 145. (e) Okamoto, K.; Yamamoto, T.; Kanbara, T. *Synlett* **2007**, 2687.

(6) (a) Davis, R. E.; Nakshbendi, H. F. *J. Am. Chem. Soc.* **1962**, *84*, 2085. (b) Hodgson, W. G.; Buckler, S. A.; Peters, G. *J. Am. Chem. Soc.* **1963**, *85*, 543. (c) Thomson, J. W.; Nagashima, K.; Macdonald, P. M.; Ozin, G. A. *J. Am. Chem. Soc.* **2011**, *133*, 5036.

The interaction of sulfur with amines has been the subject of several studies.^{6–8} McMillan observed the formation of *N*-benzylthiobenzamide when benzylamine and sulfur were heated together which is, to our knowledge, the only example of such a transformation.⁷ In the same report, the reaction has also been applied for an equimolar mixture of benzylamine and morpholine and resulted in a low yield of the corresponding *N*-(thiobenzoyl) morpholine (40.6%). No information concerning the composition of the reaction mixture has been collected. To our knowledge, such a reaction with **two different aliphatic primary amines** has never been disclosed. In this paper, we wish to report a new synthetic approach to thioamide via a sulfur-mediated selective thionation–coupling reaction starting from two different amines (Scheme 1).

Scheme 1. Thionation–Coupling Reaction of Two Amines



The reaction of sulfur with two different aliphatic primary amines (benzylamine **1a** and 2-phenethylamine **2a**, for instance), in principle, can afford four possible thioamides (Table 1).

Such a reaction that is capable of providing a single cross-coupled thioamide selectively from two amines is highly desirable. To evaluate the potential of elemental sulfur as a thionation–coupling reagent, the homocoupling reaction conditions of McMillan were reproduced by heating benzylamine **1a** with an equimolar amount of sulfur at reflux (190 °C, 1 h) (Table 1, entry 1). Since full conversion of **1a** into *N*-benzylthiobenzamide **3a** was achieved rapidly (less than 1 h), we decided to lower the reaction temperature. At 110 °C, high conversion in **3a** (>98%) was obtained after 16 h (entry 2; compare with entry 3). The same reaction was next investigated for 2-phenethylamine **2a** and required a higher temperature (150 °C) to afford a comparable high conversion (>98%) of homocoupled thioamide (entry 4). Lowering the temperature to 130 °C resulted in incomplete conversion of **3b** (40%, entry 5). We decided to use an equimolar mixture of **1a** and **2a** for reaction at 130 °C (entry 6). In this case, the ¹H NMR spectrum of the crude mixture displayed the presence of both homocoupled thioamides **3a** and **3b**, only one heterocoupled thioamide **3c**, a small quantity of **2a** (~5%), and only a trace of **1a**. Interestingly, the percentage of heterocoupled thioamide **3c** is very high (92% of thioamides). Because **3c** can be yielded by the reaction of

2a and **3a** (vide infra), an appropriate additional quantity of **2a** would help increase the conversion into **3c** and at the same time lower the amount of **3a**. To our delight, adding just 20% of **2a** to the reaction mixture substantially increased the conversion of **3a** (entry 7). However, adding more **2a** favored the formation of **3b** (entry 8).

Table 1. Reaction Conditions Screening^a

entry ^a	1a (mmol)	2a (mmol)	temp (°C)	conversion (%) ^b	3a:3b:3c
1 ^e	5	0	190	100 ^c	100:0:0
2	5	0	110	>98 ^c	100:0:0
3	5	0	100	80 ^c	100:0:0
4	0	5	150	>98 ^d	0:100:0
5	0	5	130	60 ^d	0:100:0
6	5	5	130	>98 ^c	13:3:86
7	5	6	130	>98^c (87)^f	5:3:92
8	5	7.5	130	>98 ^c	4:6:90

^a Conditions: sulfur (7.5 mmol for entries 1–5 or 15 mmol for entries 6–8, 32.1 g/mol), 16 h unless otherwise stated. ^b Based on ¹H NMR.

^c Based on the consumption of **1a**. ^d Based on the consumption of **2a**.

^e Reaction time 1 h. ^f 87% Isolated yield of **3c**.



































In subsequent studies, we applied the optimized conditions to investigate the generality of this reaction and to demonstrate its synthetic utility (Table 2). With *N*-methylbenzylamine **1b** (entry 1), **3c** was obtained in high yield. In this reaction, no product resulting from the homocoupling of **1b** could be observed. Moreover, *N*-methylthiobenzamide was the only observable intermediate of the reaction. With *N,N*-dimethylbenzylamine **1c** (entry 2), because the formation of homocoupling of **1c** is not possible, the only intermediate of the reaction *N,N*-dimethylthiobenzamide can react exclusively with **2a** to yield **3c**. When dibenzylamine **1d** (0.5 equiv, entry 3) or tribenzylamine **1e** (0.33 equiv, entry 4) was used with **2a**, all benzyl groups of **1d–e** are successively transformed into the thiobenzoyl moiety of the final thioamide **3c**. Other functional benzylamines (**1f–g**) and 3- and 4- picolinamines (**1h–i**) could also be used as the **amine 1** component in this reaction without any incidents. Different substrates could be used as the **amine 2** component, including functionalized amine (**2b**), simple aliphatic amines (*n*-octylamine and *n*-hexylamine **2c–d**), alicyclic amine (**2e**), and secondary amine (morpholine **2f**).

To our delight, 4-picolinamine is more prone to oxidation, allowing the reaction to occur at lower temperature (entries 8–12). Interestingly, 1-phenethylamine **1j** could also be successfully employed in place of the benzylamine, leading to phenylthioacetamides **3b** and **3m** in moderate to high yields. Obviously, in these two cases, the transformation

(7) (a) McMillan, F. H. *J. Am. Chem. Soc.* **1948**, *70*, 868. For related work, see: (b) Aghpoor, K.; Darabbi, H. R.; Tabar-Heydar, K. *Phosphorus, Sulfur and Silicon* **2002**, *177*, 1183.

(8) For a mechanistic study of the room temperature reaction of sulfur with benzylamine, see: Sasaki, Y.; Olsen, F. P. *Can. J. Chem.* **1971**, *49*, 283.

Table 2. Conversion of Two Different Amines into *N*-Substituted Thioamides^a

$ \begin{array}{c} \text{R}-\text{NH}_2 + \text{S} + \text{H}_2\text{N}-\text{R}' \xrightarrow{110-130\text{ }^\circ\text{C}} \text{R}-\text{C}(=\text{S})-\text{NH}-\text{R}' \\ \text{amine 1} \qquad \qquad \qquad \text{amine 2} \qquad \qquad \qquad \text{3c-m} \end{array} $						
entry	amine 1	amine 2	temp (°C)	time (h)	product	yield (%) ^b
1	 1b	 2a	130	24	 3c	90
2	 1c	2a	130	24	3c	95
3	 1d^c	2a	130	24	3c	87
4	 1e^d	2a	130	24	3c	80
5	 1a	 2b	130	16	 3d	88
6	 1f	2b	130	24	 3e	85
7	 1g	 2c	130	24	 3f	88
8	 1h	2a	130	24	 3g	90
9	 1i	 2b	110	20	 3h	91
10	1i	 2d	110	20	 3i	80
11	1i	 2e	110	24	 3j	85
12	1i	 2f	110	15	 3k	91
13	 1a	2f	130	40	 3l	70
14	 1j	 2a	130	16	 3b	93
15	1j	2f	120	40	 3m	65
16	 1a	 2g	120	60	 3n	62

^a Reaction conditions: **amine 1** (5 mmol), **amine 2** (6 mmol), sulfur (15 mmol). ^b Isolated yield. ^c 2.5 mmol. ^d 1.7 mmol.

involves a Willgerodt rearrangement. Finally, the coupled product of **1a** with 4-hydroxybutylamine **2g** provides an

attractive example in which the hydroxy group of **2g** remained intact during the reaction and may be useful for

further functionalization. In general, higher conversions of cross-coupled thioamide **3** were observed when the reaction mixture was heated for a longer time at appropriate temperatures which could not induce the oxidation of **amine 2** components or other side reactions.

Plausible mechanisms are described in Scheme 2. The upper part presents the mechanism when benzylamine **1a** was the only amine component.⁸ The first step (step **a**) of the reaction could be the oxidation of **1a** into benzaldimine **4a**. Imine **4a** is transformed into thiobenzamide **3a** via two possible pathways: (i) oxidation by sulfur (step **b**) to afford thiobenzamide **5** followed by a transthioamidation (step **c**) with benzylamine **1a**;⁹ (ii) transimination¹⁰ (step **d**) of imine **4a** with amine **1a** to provide **4b** and subsequent oxidation (step **e**) of **4b** by sulfur. Since the oxidation using the sulfur of imine **4a–b** (steps **b** and **e**) is faster than that of an amine **1a** (step **a**), the presence of imines **4a–b** could not be observed under our conditions. Because the transthioamidation (step **c**) is highly favorable when ammonia is driven off on heating,¹¹ once generated, thiobenzamide is consumed immediately. Consequently, only **1a** and **3a** could be observed in the reaction mixture by NMR spectroscopy. The rate-determining step is therefore step **a**. Contrary to one of the McMillan hypotheses⁷ which suggested that **3a** could be obtained by condensation of two benzylamine molecules followed by oxidation, we reasoned that this condensation under gentle heating conditions is likely impossible, although reaction of dibenzylamine and sulfur can give rise to **3a**.


(9) In contrast to benzamide which is relatively inert to benzylamine, thiobenzamide is more labile towards a transamidation reaction. For uncatalyzed transthioamidation, see: (a) Schlatter, M. J. *J. Am. Chem. Soc.* **1942**, *64*, 2722. For selected examples of catalyzed transamidation, see: (b) Zhang, M.; Imm, S.; Bahn, Neubert, S. L.; Neumann, H.; Beller, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 3905. (c) Nguyen, T. B.; Sorres, J.; Tran, M. Q.; Ermolenko, L.; Al-Mourabit, A. *Org. Lett.* **2012**, *14*, 3202. (d) Hoerter, J. M.; Otte, K. M.; Gellman, S. H.; Stahl, S. S. *J. Am. Chem. Soc.* **2006**, *128*, 5177. (e) Kissounko, D. A.; Guzei, L. A.; Gellman, S. H.; Stahl, S. S. *Organometallics* **2005**, *24*, 5208. (f) Eldred, S. E.; Stone, D. A.; Gellman, S. H.; Stahl, S. S. *J. Am. Chem. Soc.* **2003**, *125*, 3422. (g) Stephenson, N. A.; Zhu, J.; Gellman, S. H.; Stahl, S. S. *J. Am. Chem. Soc.* **2009**, *131*, 10003. (h) Hoerter, J. M.; Otte, K. M.; Gellman, S. H.; Cui, Q.; Stahl, S. S. *J. Am. Chem. Soc.* **2008**, *130*, 647. (i) Kissounko, D. A.; Hoerter, J. M.; Guzei, L. A.; Cui, Q.; Gellman, S. H.; Stahl, S. S. *J. Am. Chem. Soc.* **2007**, *129*, 1776. (j) Allen, C. L.; Atkinson, B. N.; Williams, J. M. J. *Angew. Chem., Int. Ed.* **2012**, *51*, 1383.

(10) Belowich, M. E.; Stoddart, J. F. *Chem. Soc. Rev.* **2012**, *41*, 2003.

(11) Wallach reported that the reaction in a sealed tube (without ammonia removal) at 180° C of benzylamine with sulfur gave thiobenzamide: Wallach, O. *Ann.* **1890**, *259*, 300. On the other hand, step **c** was not favored if *N*-methyl- was used due to steric hindrance or was not possible if *N,N*-dimethylbenzylamine was used.

When 2-phenethylamine **2a** was involved in the reaction, it could react either with both imines **4a–b** via transimination (steps **f** and **h**) or with thioamide **5** and **3a** via transthioamidation (steps **i** and **j**). Due to the similar nucleophilic nature and boiling points of amines **1a** and **2a**, steps **h** and **j** might be reversible. However, the formation of **3c** in step **j** and **4c** in step **h** can be driven by continuous oxidation of the benzylamine. Since **4b** and **4c** could be oxidized further to **3a** and **3c** (steps **e** and **g**), step **j** might be the rate-determining step. Obviously, the success of the reaction depends on the difference in reactivity of benzylamine **1a** and 2-phenethylamine **2a** in the first step of amine oxidation.

Scheme 2. Proposed Reaction Mechanism



In summary, a three-component reaction involving elemental sulfur and two different amines has been achieved for the first time and offers a straightforward and an efficient strategy for the synthesis of various thioamides. Further studies of the reaction mechanism and applications are in progress.

Acknowledgment. Financial support from CNRS and ICSN is gratefully acknowledged.

Supporting Information Available. Experimental procedures, product characterization, and copies of the ¹H and ¹³C NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.