

Syntheses and Reactivity of Aminobis(diorganylamino)phosphanes[☆]

Gerold Schick^a, Armin Loew^a, Martin Nieger^a, Karri Airola^b, and Edgar Niecke^{*a}

Anorganisch-Chemisches Institut der Universität Bonn^a,
Gerhard-Domagk-Straße 1, D-53121 Bonn, Germany

Department of Chemistry, University of Jyväskylä^b,
P.O. Box 35, FIN-40101, Jyväskylä, Finland

Received April 2, 1996

Key Words: Aminobis(diorganylamino)phosphanes / *P*-Hydrophosphane imine / Complexation reactions

Aminobis(diorganylamino)phosphanes (R₂N)₂PNH₂ (R = *i*Pr, *c*-Hex, Ph) (**2a–c**), were obtained by treatment of the corresponding chlorobis(diorganylamino)phosphanes **1a–c** with lithium amide in liquid ammonia. The transition metal phosphane complex (R₂N)₂(H₂N)PNi(CO)₃ (**3**) was prepared by reaction of **2a** with tetracarbonylnickel. *N*-Complexation of **2a** with trimethylalane affords the iminophosphorane–alane adduct (R₂N)₂HP=N(H)AlMe₃ (**4**) by 1,2-H shift from the nitrogen to the phosphorus atom. The corresponding reaction with aluminium trichloride leads to the formation of the salt [(R₂N)₂HP=N(H)]₂AlCl₂⁺[AlCl₄][–] (**5**), in which two imino-

phosphorane moieties are linked by an AlCl₂ bridge. Lithiation of **2a** and subsequent reaction with titanocene dichloride result in nucleophilic replacement of one chlorine atom to give the phosphanylamido-substituted titanocene compound (R₂N)₂P–N(H)Ti(Cl)Cp₂ (**6a**), which rearranges to its *P*-hydroiminophosphorane tautomer (R₂N)₂P(H)=NTi(Cl)Cp₂ (**6b**) on gentle heating. No isomerisation was observed in the case of *N*-borylated aminophosphane (R₂N)₂P–NH–(9-BBN) (**7**), obtained by reaction of lithiated **2a** with *B*-Cl-9-BBN. The X-ray structure analyses of compounds **2b**, **c**, **3**, **4**, **5**, and **6a** are discussed.

In the past triaminophosphanes have been the subject of extensive studies^[1]. The ready availability of substituted triaminophosphanes (R₂N)₃P by reaction of PCl₃ with secondary amines^[2], their versatile reactivity and their properties as complex ligands^[3] characterise them as valuable synthons in organophosphorus chemistry. Amongst other reactions, oxidation to phosphoranes^[4] and phosphonium salts^[5], transesterification and transamination reactions^[4] as well as displacement reactions with hydrogen halides and boron halides furnishing the corresponding aminohalophosphanes^[6] have been described. In contrast, analogous triaminophosphanes derived from primary amines (RHN)₃P lack thermal stability, due to facile amine elimination at ambient temperature, which leads to condensation reactions^[7]. Parent triaminophosphane (H₂N)₃P was postulated as an intermediate of the complete ammonolysis of phosphorus trichloride^[8]. However, there is no spectroscopic evidence known for this species. Elimination of ammonia results in the formation of not well-defined solids of varying composition. Mention should be made that thermally stable transition metal complexes (H₂N)₃PM(CO)₅ (M = Cr, Mo, W) are well-known^[9]. Mixed aminophosphanes of the type (R₂N)_{3–x}P(NH₂)_x (x = 1, 2) have not been described so far. Formation of polymeric not well-defined products was reported for the ammonolysis of (Ph₂N)PCl₂^[10], whilst oligomeric hydridophosphazenes were obtained by treatment of ammonia with (Me₂N)₂PCl, which themselves easily undergo condensation reactions at ambient temperature^[11].

Recognizing the potential synthetic utility of triaminophosphanes bearing NH₂ groups as substituents with respect to an additional reactive site, we have focussed our


attention on the stabilisation of such species by introduction of sterically protecting diorganylamino groups^[12] into the molecules.

In this paper we report on the synthesis, crystal structure, and reactivity of the first aminobis(diorganylamino)phosphanes (R₂N)₂P–NH₂.

Results and Discussion

The reactions of the bis(diorganylamino)chlorophosphanes **1a–c** with lithium amide in liquid ammonia at –35 °C afforded the corresponding aminobis(diorganylamino)phosphanes **2a–c** which were isolated as an air- and moisture-sensitive, colourless oil (**2a**) or colourless crystals (**2b**, **c**) (Scheme 1). The compounds can be stored at low temperatures for weeks without decomposition. However, in solution at ambient temperature they slowly undergo condensation reactions to give oligomeric and polymeric hydridophosphazenes^[13] by elimination of the corresponding diorganylamine. Elimination of ammonia and formation of diphosphanylamines R₂P–NH–PR₂ as reported for Me₂PNH₂^[2c] or Ph₂PNH₂^[14] were not detected in the case of compounds **2a–c**.


Scheme 1



Their constitution became evident from multinuclear NMR investigations. The replacement of the chlorine atom in the chlorophosphanes **1a–c** by the amido group causes a significant downfield shift of the phosphorus atom signal in the ^{31}P -NMR spectrum (**2a**: $\Delta\delta = 62.0$, $\delta = 77.0$; **2b**: $\Delta\delta = 64.0$, $\delta = 80.4$; **2c**: $\Delta\delta = 52.0$, $\delta = 71.5$), as observed in the case of tris(diorganylamino)-substituted derivatives^[15]. The presence of the NH_2 group in **2c** is evidenced by the coupling of the two amino protons with the phosphorus atom ($^2J_{\text{PH}} = 17.1$ Hz). This coupling was not observed in the case of compound **2a** or **2b**, which might be the result of a fast intermolecular proton exchange.

Suitable crystals of compounds **2b** and **c** for X-ray structure analyses were obtained from pentane/diethyl ether at -78°C . Their structures are given in Figures 1 and 2 and selected structural parameters are listed in Table 1. The geometry around the phosphorus atom in **2b** and **c** is distorted trigonal-pyramidal (**2b**: ΣP1 : 307° , **2c**: ΣP1 : 306°), with typical $\text{P(III)}-\text{N}^{[16]}$ bond lengths. Whilst in **2b** the P–N bond lengths between the phosphorus atom and the two R_2N groups [$\text{P1}-\text{N2}$: $169.1(2)$, $\text{P1}-\text{N3}$: $169.5(2)$] are slightly shorter than the bond length between the phosphorus and the nitrogen atom of the NH_2 group [$\text{P1}-\text{N1}$: $172.2(2)$], the situation is reversed in **2c** [$\text{P1}-\text{N1}$: $167.0(2)$, $\text{P1}-\text{N2}$: $173.4(2)$, $\text{P1}-\text{N3}$: $173.3(2)$]. In contrast to the nitrogen atoms in the diorganylamino groups, which exhibit a planar geometry (**2b**: ΣN2 : 360° , ΣN3 : 360° ; **2c**: ΣN2 : 359° , ΣN3 : 360°), the nitrogen atoms of the NH_2 groups show a distinct pyramidalisation (**2b**: ΣN1 : 326° ; **2c**: ΣN1 : 347°). Two molecules of **2b** are connected by van der Waals contacts between the phosphorus atom and one of the hydrogen atoms of the NH_2 group (288 pm). In **2c** the intramolecular distances between one NH proton and the carbon atoms of one phenyl group ($\text{H1}\cdots\text{C1}-\text{C6}$: 280 – 331

Figure 1. Molecular structure of aminophosphane **2b** in the solid state. C–H hydrogen atoms are omitted for clarity, second molecule generated by transformation $-x, 1-y, -z$



pm) lie in the range of the sum of the van der Waals radii.

Due to facile elimination of amine all attempts to crystallise **2a** have failed so far. However, the corresponding phosphane complex **3**, obtained by treatment of **2a** with tetra-

Figure 2. Molecular structure of aminophosphane **2c** in the solid state. C–H hydrogen atoms are omitted for clarity, second molecule generated by transformation $1-x, 1-y, 1-z$




Table 1. Selected bond lengths [pm] and bond angles $^\circ$ of compounds **2b**, **2c**, **3**, and **6a**

	2b	2c	3	3'	6a
P1–N1	172.2(2)	167.0(2)	168.3(2)	168.1(3)	172.3(3)
P1–N2	169.1(2)	173.4(2)	169.2(2)	169.5(2)	169.4(3)
P1–N3	169.5(2)	173.3(2)	168.4(3)	168.6(2)	169.7(3)
$\Sigma^\circ \text{P1}$	307	306	–	–	312
$\Sigma^\circ \text{N1}$	326	347	324	330	359
$\Sigma^\circ \text{N2}$	360	359	356	356	359
$\Sigma^\circ \text{N3}$	360	360	360	360	360
N1–P1–N2	98.7(1)	102.5(1)	101.7(1)	98.7(1)	102.7(2)
N1–P1–N3	97.4(1)	102.4(1)	106.3(1)	109.5(1)	101.6(1)
N2–P1–N3	110.6(1)	100.7(1)	103.0(1)	102.5(1)	107.2(2)
N–C ^[a]	148	143	148	148	148
Ni1–P1	–	–	224.4(1)	223.8(1)	–
Ti1–N1	–	–	–	–	193.7(3)
Ti–Cp (centroid ^[b])	–	–	–	–	209
Ti–Cl1	–	–	–	–	242.1(1)
P1–Ni1–Ti1	–	–	–	–	141.5(2)
N1–Ti1–Cl1	–	–	–	–	96.8(1)


^[a] Average N–C bond lengths are given. – ^[b] Average Ti–Cp(centroid) distance.

carbonylnickel, was isolated by crystallisation from toluene as colourless crystals suitable for an X-ray analysis (Figure 3).

Two independent, nearly identical molecules are found in the asymmetric unit of phosphane complex **3**. The phosphorus and the nickel atom exhibit a distorted tetrahedral environment, and the three P–N bond lengths vary only slightly around the average bond length of 169 pm (Table 1). The structural parameters of the $\text{P}-\text{Ni}(\text{CO})_3$ moiety are consistent with those reported previously for $(\text{CO})_3\text{Ni}$ -phosphane complexes^[17].


In contrast to the transition metal fragment, which attacks the phosphorus atom, a hard Lewis acid should prefer a complexation via the nitrogen atom of the amino group. This expectation was verified by reaction of **2a** with AlMe_3 in pentane at -78°C and resulted in the formation of the *P*-hydrophosphoranimine-*N*-alane adduct **4** (Scheme 2).

Figure 3. Molecular structure of one of the crystallographically independent molecules of **3**. C–H hydrogen atoms are omitted for clarity



Coordination of the Lewis acid at the less sterically protected nitrogen atom of the NH_2 group with subsequent hydrogen shift from the nitrogen to the phosphorus atom was also observed in the reaction of **2a** with AlCl_3 . However, here the salt $[(\text{R}_2\text{N})_2\text{HP}=\text{N}(\text{H})]_2\text{AlCl}_2^+ [\text{AlCl}_4]^-$ (**5**) was formed (Scheme 2).

Scheme 2




The isomerisation of N–H aminophosphanes $\text{R}_2\text{PN}(\text{H})\text{R}'$ to the corresponding P–H iminophosphoranes $\text{R}_2\text{P}(\text{H})\text{NR}'$ has been the subject of several theoretical and experimental investigations^[18]. The relative stability of both isomers is known to be dependent on the steric as well as the electronic influences exerted by the substituents R and R'. Electron-withdrawing groups R' such as $\text{SO}_2\text{C}_6\text{H}_4\text{CH}_3$ and $\text{PS}(\text{OC}_6\text{H}_5)_2$ as well as bulky substituents R and R' are known to stabilize the P–H isomer. To the best of our knowledge, compounds **4** and **5** are the first examples in which the rearrangement of an aminophosphane to its phosphorane tautomer is induced by complexation by Lewis acids at the nitrogen lone pair^[19].

Diagnostic of the 1,2-hydrogen shift from the nitrogen to the phosphorus atom in compound **4** are the high-field shift of the ^{31}P -NMR signal ($\delta = 13.7$) and the characteristic PH coupling constant^[20] ($^1J_{\text{PH}} = 563.3$ Hz). In the ^1H -NMR spectrum the proton bound directly to the phosphorus atom gives rise to a doublet of doublets ($\delta = 7.05$, $^1J_{\text{PH}} = 563.3$, $^3J_{\text{HH}} = 11.9$ Hz), which clearly indicates a species with the $(>\text{N})_2\text{P}(\text{H})\text{NH}$ - moiety. Very similar ^{31}P - and ^1H -NMR spectra were observed for compound **5** (^{31}P -NMR: $\delta = 13.8$; ^1H NMR: $\delta = 7.22$, $^1J_{\text{PH}} = 578.6$, $^3J_{\text{HH}} = 10.8$ Hz).

The formation of the salt **5** in solution becomes apparent from the ^{27}Al -NMR spectrum which shows the characteristic signal of AlCl_4^- ($\delta = 101.2$). The Al nuclei in the cation gives rise to a broadened resonance at $\delta = 100.2$. Single crystals of products **4** and **5** were obtained by recrystallisation from dichloromethane and dichloromethane/pentane. Their molecular structures were verified by X-ray structure analyses to be bis(diisopropylamino)phosphane imine-*N*-trimethylalane (**4**) and bis[bis(diisopropylamino)phosphane iminato]dichloroaluminium(III) tetrachloroaluminate(III) (**5**) (Figures 4 and 5).

Figure 4. Molecular structure of the alane adduct **4**. C–H hydrogen atoms are omitted for clarity



Whereas the structure of compound **4** proves the formation of a *P*-hydrophosphane imine- AlMe_3 adduct, two equivalents of both **2a** and AlCl_3 react to give the salt **5**.

Figure 5. Molecular structure of **5**. C–H hydrogen atoms are omitted for clarity




Table 2. Selected bond lengths [pm] and bond angles [°] of compounds **4** and **5**

Bond	4	5	Bond angle	4	5
P1–N1	154.9(5)	161.7(6)	P1–N1–Al1	136.4(3)	130.4(4)
P2–N4	–	161.0(6)	P2–N4–Al1	–	128.1(4)
P1–N2	162.3(4)	162.6(6)	N1–P1–N2	113.7(3)	109.9(3)
P1–N3	161.4(4)	1.626(6)	N1–P1–N3	121.8(3)	118.0(4)
P2–N5	–	162.4(6)	N2–P1–N3	109.2(2)	107.2(3)
P2–N6	–	165.1(6)	N4–P2–N5	–	117.8(3)
Al1–N1	192.8(6)	184.7(7)	N4–P2–N6	–	109.5(3)
Al1–N4	–	184.9(6)	N5–P2–N6	–	108.8(3)
			N1–Al1–N4	–	106.8(3)

The cation in **5** consists of two phosphane imine groups, which are linked by their imine nitrogen atoms by a bridging AlCl_2 group. The anion $[\text{AlCl}_4]^-$ shows only a slight deviation from the ideal tetrahedral geometry, whereby two chlorine atoms participate in the formation of weak $\text{Cl} \cdots \text{H} \cdots \text{N}$ hydrogen bonds ($\text{N1} \cdots \text{Cl3}$: 351, $\text{N4} \cdots \text{Cl4}$: 360 pm). The imine hydrogen atoms and the P–H hydrogen atoms adopt an almost antiperiplanar conformation in both compounds.

Whilst the Al–N bond length (Table 2) in **4** [192.8(6) pm] is of similar magnitude as reported for compounds containing an imine nitrogen–aluminium moiety [$(\text{Me}_2\text{C}=\text{NAlMe}_2)_2$: 195.5, 193.0 pm^[21]; $\text{N}(\text{PPh}_2\text{NSiMe}_3)_2\text{AlMe}_2$: 194.0 pm^[22]], the Al–N bond lengths in **5** are significantly shortened [184.7(7), 184.9(6) pm]. The situation is reversed in the case of the P–N(H) bond lengths for both compounds [**4**: 154.9(5); **5**: 161.7(6), 161.0(6) pm]. The bond lengths between the phosphorus and the two nitrogen atoms of the diorganylamino groups differ only slightly for both compounds [**4**, **5**: 163 pm (average)]. These results reflect a different extent of polarisation in the P–N–Al fragment in both compounds. The further bond lengths and angles show no particularity.


Replacement of the hydrogen at the NH_2 group in **2a** by a transition metal fragment was easily achieved by metalation with *n*-butyllithium (at -30°C in THF) and subsequent reaction with a transition metal halide. For **2a** and titanocene dichloride this afforded the titanocene complex

6a (Scheme 2), which was isolated by crystallisation from diethyl ether as dark red crystals. On gentle heating to 40°C , **6a** slowly rearranges in solution to the thermodynamically more stable *P*-hydrophosphane iminato isomer **6b**.

The rearrangement of **6a** to its phosphorane tautomer **6b** was unambiguously proven by a significant shielding of the phosphorus atom (**6a**: $\delta = 98.6$; **6b**: $\delta = -3.9$) as well as the characteristic PH coupling constant ($^1J_{\text{PH}} = 521.4$ Hz).

The results of an X-ray structure analysis of the titanocene complex **6a** are illustrated in Figure 6, and selected structural parameters are given in Table 1. The P–N bond lengths and angles in **6a** differ only slightly from those in **2b** except for the planarity at the titanium-bound nitrogen (ΣN1 : 359°). The Ti–N bond length of 193.7(3) pm is significantly shorter than those found in the related compound $(\eta^5\text{-Cp})_2\text{Ti}(\eta^1\text{-NC}_4\text{H}_4)_2$ ^[23] (208.5 pm). The planarity at N1, the short Ti–N bond, the nearly eclipsed conformation of the lone pair at N1 and the chlorine atom indicate $p_\pi\text{-}d_\pi$ interaction between titanium and nitrogen. The preference of this conformation was predicted by Lauher and Hoffmann^[24] for $\text{Cp}_2\text{M}(\text{NR}_2)\text{L}'$ complexes ($\text{L}' = \text{non } \pi\text{-donor ligand}$) and is explained in terms of a π -back donation from the N lone pair to an empty a_1 orbital of the bent titanocene unit.

Figure 6. Molecular structure of **6a**. C–H hydrogen atoms are omitted for clarity



A proton transfer from N to P was not observed in the case of the *N*-borylated phosphane **7**, which was obtained by treatment of lithiated **2a** with *B*-Cl-9-BBN at -78°C in THF (Scheme 2). The higher thermodynamic stability of aminophosphane **7** compared to its P–H isomer was already demonstrated by Bertrand et al., who observed the formation of analogous *N*-borylated aminophosphanes by hydrogen transfer from the corresponding *P*-hydrophosphane iminato isomer, obtained by reaction of bis(diorganylamino)phosphanyl azides with trialkylboranes^[25]. The structure of **7** was derived from its NMR spectroscopic data, which are in accordance with those found for compounds of the type $(i\text{Pr}_2\text{N})_2\text{P} \cdots \text{NH} \cdots \text{BR}_2$.

We thank the *Fonds der Chemischen Industrie* for financial support. K. A. thanks the *Finnish Academy* for financial support.

Experimental

All manipulations were carried out with the exclusion of air and moisture under argon. Solvents were dried by using standard procedures. The bis(diorganylamino)chlorophosphanes **1a**^[26], **1b**^[26], **1c**^[27], and *B*-Cl-9-BBN^[28] were prepared by literature methods. Tetracarbonylnickel, Ni(CO)₄, dichlorobis(cyclopentadienyl)titanium(IV), Cp₂TiCl₂, trimethylalane, AlMe₃ (2 M in toluene), and aluminium trichloride were used as supplied. — NMR: Bruker AMX 300; ³¹P: 121.5 MHz, external standard 85% H₃PO₄; ¹H: 300.1 MHz, external standard TMS; ¹³C: 75.5 MHz, external standard TMS; ¹¹B: 96.3 MHz, external standard BF₃ · O(C₂H₅)₂; ²⁷Al: 300.1 MHz, Al(H₂O)₆³⁺ external standard; positive signs denote shifts to lower frequencies. — MS: Kratos Instruments Concept 1H, Kratos Instruments, MS 50, VG Instruments VG 12-250 (EI, 70 eV). — Melting points were determined in sealed glass capillaries and are uncorrected.

Aminobis(diisopropylamino)phosphane (2a), *Aminobis(dicyclohexylamino)phosphane (2b)*, and *Aminobis(diphenylamino)phosphane (2c)*: To a suspension of 15 ml of LiNH₂ (1.1 g) in 75 ml of liquid ammonia a solution of 50 mmol of **1** (a: 13.3 g; b: 21.3 g; c: 20.1 g) in 25 ml of THF is slowly added at −40 °C. After stirring of the combined solutions at −35 °C for some time (a: 45 min; b: 6 h; c: 45 min) the ammonia is evaporated and the organic solvent removed under reduced pressure. To the remaining residue 30 ml of pentane (c: pentane/toluene, 1:1) is added, the precipitated LiCl is filtered off and the solvent removed under reduced pressure. Pure **2b** and **2c** are obtained as colourless solids by recrystallisation from diethyl ether at −78 °C, whereas the liquid **2a** is purified by distillation at 10^{−3} Torr at 45–50 °C.

2a: Yield: 16.8 g (68%), b.p. 40–43 °C/10^{−3} Torr. — ³¹P{¹H} NMR (C₆D₆): δ = 77.0 (s). — ¹³C{¹H} NMR (C₆D₆): δ = 44.2 (d, ²J_{CP} = 12.2 Hz, PNC), 25.1 (d, ³J_{CP} = 8.6 Hz, CH₃). — ¹H NMR (C₆D₆): δ = 3.6 (m, 4H, CH), 1.1 (d, ³J_{HH} = 6.7 Hz, 24H, CH₃), NH₂ not detected.

2b: Yield: 14.6 g (72%), m.p. 43–45 °C (dec.). — ³¹P{¹H} NMR (C₆D₆): δ = 80.4 (s). — ¹³C{¹H} NMR (C₆D₆): δ = 54.5 (d, ²J_{CP} = 9.9 Hz, PNC), 36.6 (d, ³J_{CP} = 8.4 Hz, NCC), 35.3 (d, ³J_{CP} = 6.8 Hz, NCC), 27.9 (s, NCCC), 27.7 (s, NCCC), 26.6 (s, NCCCC). — MS (180 °C/70 eV), *m/z* (%): 407 (6) [M⁺], 324 (1) [M⁺ − C₆H₁₁], 227 (52) [M⁺ − N(C₆H₁₁)₂], 181 (55) [HN(C₆H₁₁)₂]⁺, 138 (100) [NC₅H₇]⁺, and other fragments. — C₂₄H₄₆N₃P: calcd. 407.3429; found 407.3441 (MS).

2c: Yield: 13.6 g (67%), m.p. 72–76 °C (dec.). — ³¹P{¹H} NMR (C₆D₆): δ = 71.5 (s). — ¹³C{¹H} NMR (C₆D₆): δ = 146.3 (d, ²J_{CP} = 8.7 Hz, PNC), 129.8 (s, C-*p*-ph), 125.3 (d, ³J_{CP} = 8.4 Hz, C-*o*-ph), 123.8 (d, ⁴J_{CP} = 0.9 Hz, C-*m*-ph). — ¹H NMR (C₆D₆): δ = 7.25–6.95 (m, 20H, Ph), 1.70 (d, ²J_{HP} = 17.1 Hz, 2H, NH₂).

[*Aminobis(diisopropylamino)phosphane*]tricarboxynickel(0) (**3**): To a solution of 2.47 g (10 mmol) of **2a** in 15 ml of toluene 17 mg (10 mmol) of tetracarbonylnickel(0) is quickly added, and the mixture is stirred for 20 min. The resulting solution is stored for several days at −80 °C. The separated colourless crystals of **3** are collected by filtration. Yield: 2.3 g (59%), m.p. 20 °C. — ³¹P{¹H} NMR (C₆D₆): δ = 125.6 (s). — ¹³C{¹H} NMR (C₆D₆): δ = 197.7 (s, CO), 191.9 (s, CO), 46.7 (d, ²J_{CP} = 14.1 Hz, PNC), 23.5 (d, ³J_{CP} = 2.1 Hz, CCH₃). — ¹H NMR (C₆D₆): δ = 3.84 (m, 4H, CH), 1.97 [s (b), 2H, NH₂], 1.23 (d, ³J_{HH} = 6.9 Hz, 24H, CH₃). — MS (180 °C/70 eV), *m/z* (%): 389 (5) [M⁺], 361 (30) [M⁺ − CO], 247 (9) [M⁺ − Ni(CO)₃], 147 (100) [M⁺ − Ni(CO)₃ − N(C₃H₇)₂], and other fragments.

Bis(diisopropylamino)phosphane imine-N-Trimethylalane (4): To a solution of 1.8 g (7.3 mmol) of **2a** in 40 ml of pentane 3.7 ml (7.3 mmol, 2 M in hexane) of AlMe₃ is added dropwise at −78 °C. While stirring the reaction mixture is allowed to warm up to room temp. The resulting white precipitate of **4** is filtered off, dried in vacuo and recrystallised from a small amount of dichloromethane. Yield 1.74 g (75%), m.p. 123–125 °C (dec.). — ³¹P{¹H} NMR (C₆D₆): δ = 13.7 (s). — ¹³C{¹H} NMR (C₆D₆): δ = 46.2 (d, ²J_{CP} = 5.34 Hz, PNC), 23.8 (d, ³J_{CP} = 2.7 Hz, CCH₃), 22.2 (d, ³J_{CP} = 2.7 Hz, CCH₃), −5.5 [s (b), AlCH₃]. — ¹H NMR (C₆D₆): δ = 7.05 (dd, ¹J_{HP} = 563.3, ³J_{HH(imine)} = 11.9 Hz, 1H, PH), 3.18 (d sept, ³J_{HP} = 17.1, ³J_{HH} = 6.8 Hz, 4H, CH), 1.09 (d, ³J_{HH} = 6.8 Hz, 12H, CCH₃), 0.96 (d, ³J_{HH} = 6.8 Hz, 12H, CCH₃), −0.1 [s (b), 9H, AlCH₃].

Bis[bis(diisopropylamino)phosphane iminato]dichloroaluminum(III) tetrachloroaluminate(III) (5): To a solution of 0.56 g (2.3 mmol) of **2a** in 30 ml of dichloromethane 0.3 g (2.3 mmol) of AlCl₃ of AlCl₃ is added at −78 °C. While stirring the reaction mixture is allowed to warm up to room temp. After evaporation of the solvent, **5** is obtained as a white powder and recrystallised from dichloromethane/pentane/toluene (1:1:1). Yield: 0.31 g, m.p. 162–165 °C (dec.). — ³¹P{¹H} NMR (C₆D₆): δ = 13.8 (s). — ¹³C{¹H} NMR (C₆D₆): δ = 47.3 [s (b), PNC], 24.3 (s, CCH₃), 23.3 (s, CCH₃). — ¹H NMR (C₆D₆): δ = 7.22 (dd, ¹J_{HP} = 578.6, ³J_{HH(imine)} = 10.9 Hz, 2H, PH), 3.45 (d sept, ³J_{HP} = 17.1, ³J_{HH} = 6.8 Hz, 8H, CH), 1.19 (d, ³J_{HH} = 6.8 Hz, 24H, CCH₃), 1.13 (d, ³J_{HH} = 6.8 Hz, 24H, CCH₃). — ²⁷Al NMR: δ = 101.2 (AlCl₄), 100.2 (AlCl₂).

[*Bis(diisopropylamino)phosphanyl-amido*]chlorobis(cyclopentadienyl)titanium(IV) (**6a**) and [*Bis(diisopropylamino)phosphane imido*]chlorobis(cyclopentadienyl)titanium(IV) (**6b**): To a solution of 0.46 g (1.9 mmol) of **2a** in 20 ml of THF 1.8 ml (1.9 mmol, 1.6 M in *n*-hexane) of *n*-butyllithium is added at −40 °C. The reaction mixture is allowed to warm up to −15 °C, then a solution of 0.47 g (1.9 mmol) of Cp₂TiCl₂ in 30 ml of THF is added rapidly. The resulting dark red solution is stirred for 1 h, then the organic solvent is removed in vacuo. To the residue 30 ml of pentane/toluene (1:1) is added. The precipitated LiCl is filtered off and the solvent removed in vacuo. **6a** is recrystallised from diethyl ether at −30 °C. If the reaction solution is gently heated at 30 °C for 4 h, **6a** rearranges to compound **6b**. The rearrangement is accompanied by partial decomposition of **6a**.

6a: Yield: 0.46 g (53%), m.p. 139 °C (dec.). — ³¹P{¹H} NMR (CDCl₃): δ = 98.9 (s). — ¹³C{¹H} NMR (CDCl₃): δ = 114.5 (s, C-*cp*), 45.6 (d, ²J_{CP} = 11.4 Hz, PNC), 25.2 (d, ³J_{CP} = 8.8 Hz, PNCC), 24.7 (³J_{CP} = 5.7 Hz, PNCC). — ¹H NMR (CDCl₃): δ = 6.24 (s, 10H, H-*cp*), 3.3 (m, 4H, NCH), 1.27 (d, ³J_{HH} = 6.8 Hz, 12H, CCH₃), 1.17 (d, ³J_{HH} = 6.8 Hz, 12H, CCH₃), NH not detected. — MS (180 °C/70 eV), *m/z* (%): 459 (1) [M⁺], 394 (4) [M⁺ − *cp*], 359 (17) [M⁺ − N(C₃H₇)₂], 323 (11) [M⁺ − N(C₃H₇)₂ − Cl], 246 (42) [M⁺ − ClTi(Cp)₂], 213 (40) [ClTi(Cp)₂], and other fragments. — C₂₂H₃₉ClN₃PTi: calcd. 457.2049; found 457.2078 (MS).

6b: ³¹P{¹H} NMR (CDCl₃): δ = −3.9 (s). — ¹³C{¹H} NMR (CDCl₃): δ = 113.7 (s, C-*cp*), 45.1 (d, ²J_{CP} = 5.7 Hz, PNC), 24.0 (s, PNCC), 23.8 (s, PNCC). — ¹H NMR (CDCl₃): δ = 7.38 (d, ¹J_{HP} = 525.8 Hz, 1H, PH), 6.07 (s, 10H, H-*cp*), 3.5 (dsept, ³J_{HP} = 16.5, ³J_{HH} = 6.9 Hz, 4H, NCH), 1.29 (d, ³J_{HH} = 6.9 Hz, 12H, CCH₃), 1.20 (d, ³J_{HH} = 6.9 Hz, 12H, CCH₃).

[*(9-Borabicyclo[3.3.1]nonan-9-yl)amino*]bis(diisopropylamino)phosphane (**7**): To a solution of 0.49 g (2.0 mmol) of **2a** in 20 ml of THF 1.3 ml (2.0 mmol, 1.6 M in *n*-hexane) of *n*-butyllithium is added at −30 °C. The solution is then cooled to −78 °C, and a

Table 3. Crystallographic data and parameters of crystal structure determinations for **2b**, **c**, **3**, **4**, **5**, and **6a**

	2b	2c	3	4	5	6a
formula	C ₂₄ H ₄₆ N ₃ P	C ₂₄ H ₂₂ N ₃ P	C ₁₅ H ₃₀ N ₃ O ₃ PNI	C ₁₅ H ₃₉ AlN ₃ P	C ₂₄ H ₆₀ Al ₂ Cl ₆ N ₆ P ₂	C ₂₂ H ₃₉ ClN ₃ PTi
M _r	407.6	383.4	390.1	319.4	761.4	459.9
crystal system	triclinic	triclinic	triclinic	orthorhombic	monoclinic	monoclinic
space group	P1 (no. 2)	P1 (no.2)	P1 (no.2)	P2 ₁ 2 ₁ 2 ₁ (no. 19)	Cc (no. 9)	C2/c (no. 15)
a [Å]	9.846(1)	9.623(2)	9.208(2)	9.687(1)	9.624(2)	29.708(4)
b [Å]	10.858(2)	10.678(2)	14.358(2)	12.270(2)	27.826(2)	10.530(1)
c [Å]	12.796(2)	11.264(1)	16.671(2)	17.754(2)	15.018(3)	16.168(1)
α [°]	70.72(1)	77.65(1)	111.21(1)	90°	90°	90°
β [°]	74.07(1)	89.00(1)	92.12(1)	90°	91.19(1)	103.79(1)
γ [°]	80.61(1)	63.80(1)	91.55(1)	90°	90°	90°
V [Å ³]	1237.7(3)	1010.6(3)	2051(1)	2110.2(4)	4021(1)	4912.0(9)
Z	2	2	4	4	4	8
ρ (g cm ⁻³)	1.09	1.26	1.27	1.01	1.26	1.24
μ (mm ⁻¹)	1.06	1.30	2.23	1.52	5.26	4.65
F(000)	452	404	832	712	1616	1968
diffractometer	Enraf-Nonius CAD4	Enraf-Nonius CAD4	Enraf-Nonius CAD4	Enraf-Nonius CAD4	Enraf-Nonius CAD4	Enraf-Nonius CAD4
radiation	CuKα	CuKα	CuKα	CuKα	CuKα	CuKα
λ [Å]	1.54178	1.54178	1.54178	1.54178	1.54178	1.54178
crystal size [mm]	0.58 x 0.45 x 0.40	0.30 x 0.15 x 0.10	0.40 x 0.40 x 0.50	0.65 x 0.55 x 0.30	0.60 x 0.50 x 0.30	0.40 x 0.25 x 0.18
T [K]	200(2)	200(2)	193(2)	208(2)	200(2)	208(2)
2θ _{max} [°]	120	120	120	120	130	120
	-10 ≤ h ≤ 11	-9 ≤ h ≤ 10	-10 ≤ h ≤ 10	-10 ≤ h ≤ 3	-11 ≤ h ≤ 11	0 ≤ h ≤ 33
	0 ≤ k ≤ 12	0 ≤ k ≤ 11	-16 ≤ k ≤ 14	-13 ≤ k ≤ 5	-32 ≤ k ≤ 14	-11 ≤ k ≤ 0
	-13 ≤ l ≤ 14	-12 ≤ l ≤ 12	0 ≤ l ≤ 18	-18 ≤ l ≤ 19	-17 ≤ l ≤ 1	-18 ≤ l ≤ 17
no. of measured data	3956	3234	6313	3938	4530	3719
no. of unique data	3670	2995	6072 ^{a)}	2912	3622	3640
full matrix least-squares refinement on	F ²	F ²	F	F ²	F ²	F ²
no. of parameters/restraints	260/2	259/2	430	195/2	390/8	265/1
R [for I > 2σ(I)]	0.068	0.057	0.047	0.089	0.081	0.059
wR2	0.185	0.166	0.055 ^{b)}	0.234	0.210	0.170

^{a)} 5760 reflections with $F > 4\sigma(F)$ used for all calculations. — ^{b)} R_w .

solution of 0.31 g (2.0 mmol) of *B*-Cl-9-BBN in 10 ml of pentane is added dropwise. During stirring, the reaction mixture is allowed to warm up to room temp. The organic solvent is then removed under reduced pressure, pentane is added to the resulting residue, and the precipitated LiCl is filtered off. **7** is purified by crystallisation from a small amount of pentane at -80 °C. Yield: 0.58 g (80%), m.p. 87 °C (dec.). — ³¹P{¹H} NMR (CDCl₃): δ = 73.8 (s). — ¹³C{¹H} NMR (CDCl₃): δ = 45.4 (d, ²J_{CP} = 12.6 Hz, PNC), 33.7 (d, ⁴J_{CP} = 33 Hz, BCC), 29.2 (m, b, BC), 24.2 (s, BCCC), 25.2 (d, ³J_{CP} = 7.6 Hz, PNCC), 23.7 (d, ³J_{CP} = 5.7 Hz, PNCC). — ¹H NMR (CDCl₃): δ = 4.76 [s (b), NH], 3.48 (d sept, ³J_{HP} = 10.8, ³J_{HH} = 5.6 Hz, 4H, NCH), 1.9–1.3 (m, 14H, 9-BBN), 1.15 (d, ³J_{HH} = 5.6 Hz, 12H, NCCH₃), 1.13 (d, ³J_{HH} = 5.6 Hz, 12H, NCCH₃). — ¹¹B NMR (CDCl₃): δ = 59.4 [s (b), w/2 = 420 Hz]. — MS (180 °C/70 eV), *m/z* (%): 367 (12) [M⁺], 267 (100) [M⁺ - *i*Pr₂N], 231 (22) [M⁺ - C₈H₁₅BN], and other fragments. — C₂₀H₄₃BN₃P: calcd. 366.3288; found 366.3305 (MS).

X-Ray Structure Determination of 2b, c, 3, 4, 5, and 6a: The structures were solved by direct methods (SHELXTL-Plus^[29]). The non-hydrogen atoms were refined anisotropically, H atoms were refined by using a riding model [full-matrix least-squares refinement on F^2 (SHELXL-93^[30]), and F (SHELXTL-Plus^[29] for **3**), respectively]. An absorption correction was applied on the basis of Ψ scans (**2c**, min./max. transmission = 0.734/0.976; **5** = 0.147/0.953, **6a** = 0.475/0.972) and DIFABS^[31] (**2b**, **3**, **4**). In the case of **5** the absolute structure was determined by x refinement^[32] [Flack's x parameter -0.02(3)], in the case of **4** the absolute structure could not be determined reliably. With **2b** and **5** an extinction correction was applied.

Details on data collection and structure refinement are given in Table 3. — Further details of the crystal structure investigations may be obtained from the Fachinformationszentrum Karlsruhe, D-76344 Eggenstein-Leopoldshafen (FRG), on quoting the depositary number CSD-59318.

★ Dedicated to Prof. Dr. Drs. h.c. H. G. von Schnering on the occasion of his 65th birthday.

- [1] D. E. C. Corbridge, *Studies in Inorganic Chemistry, Phosphorus*, 2nd ed., Elsevier Scientific Publishing Company, Amsterdam, 1980, and references therein.
- [2] [2a] G. M. Kosolapoff, L. Maier, *Organophosphorus Compounds*, John Wiley & Sons, Inc., New York, 1973, vol. 5, p. 88. — [2b] R. B. King, N. D. Sadanani, *Synth. React. Org. Met.-Org. Chem.* **1985**, 15(2), 149–158. — [2c] A. B. Burg, P. J. Slota jr., *J. Am. Chem. Soc.* **1958**, 80, 1107–1109.
- [3] [3a] A. H. Cowley, R. E. Davis, K. Remadua, *Inorg. Chem.* **1981**, 20, 2146–2152. — [3b] H. Inoue, T. Nakagome, T. Kuroiwa, T. Shirai, E. Fluck, *Z. Naturforsch., Part B*, **1987**, 42, 573–578.
- [4] [4a] R. Appel, U. Baumeister, F. Knoch, *Chem. Ber.* **1983**, 116, 75–86. — [4b] H. Nöth, H. J. Vetter, *Chem. Ber.* **1963**, 96, 1308–1315.
- [5] H. Nöth, H. J. Vetter, *Chem. Ber.* **1965**, 98, 1981–1987.
- [6] H. Nöth, H. J. Vetter, *Chem. Ber.* **1963**, 94, 1109–1118.
- [7] [7a] R. A. Keat, *Top. Current. Chem.* **1982**, 102, 89–116. — [7b] A. Tarassoli, R. Haltiwanger, A. D. Norman, *Inorg. Chem.* **1982**, 21, 2684–2690. — [7c] R. R. Holmes, J. A. Forstner, *J. Am. Chem. Soc.* **1960**, 82, 5509–5510. — [7d] M. L. Thompson, A. Tarassoli, R. Haltiwanger, A. D. Norman, *Inorg. Chem.* **1987**, 26, 684–689.
- [8] E. Fluck, *Top. Phosphorus Chem.*, **1967**, 4, 295.
- [9] [9a] H. Nöth, H. Reith, V. Thorn, *J. Organomet. Chem.* **1978**,

- 159, 165–169. — [19b] H. Maisch, *Z. Naturforsch., Part B*, **1979**, *34*, 784–789.
- [10] M. Becke-Goehring, J. Schulze, *Chem. Ber.* **1958**, *91*, 1188–1195.
- [11] A. Schmidpeter, H. Roßknecht, *Chem. Ber.* **1974**, *107*, 3146–3148.
- [12] R. W. Reed, G. Bertrand in *Phosphorus-31 NMR Spectral Properties in Compound Characterisation and Structural Analysis* (Eds.: L. D. Quin, J. G. Verkade), VCH, Weinheim, **1994**, chapter IV, p. 15.
- [13] According to molecular mass determination the reaction product consists mainly of cyclotri- and cyclotetraphosphazenes which could be isolated in pure form: G. Schick, E. Niecke, unpublished results.
- [14] O. Schmitz du Mont, B. Ross, H. Klieber, *Angew. Chem.* **1967**, *79*, 869; *Angew. Chem. Int. Ed. Engl.* **1967**, *79*, 875.
- [15] J. P. Gouesnard, J. Dorie, G. J. Martin, *Can. J. Chem.* **1980**, *58*, 1295–1304.
- [16] G. Trinquier, M. T. Ashby, *Inorg. Chem.* **1994**, *33*, 1306–1313.
- [17] [17a] L. Horner, H. Kunz, *Chem. Ber.* **1971**, *104*, 717. — [17b] J. Pickardt, L. Rösch, H. Schumann, *Z. Anorg. Allg. Chem.* **1976**, *426*, 66.
- [18] [18a] D. Gonbeau, G. Pfister-Guillouzo, M. Mazieres, M. Sanchez, *Can. J. Chem.* **1985**, *63*, 3242–3248. — [18b] P. V. Sudhakar, K. Lammertsma, *J. Am. Chem. Soc.* **1991**, *113*, 1899–1906. — [18c] A. Schmidpeter, J. Ebeling, *Angew. Chem.* **1968**, *80*, 197–198; *Angew. Chem. Int. Ed. Engl.* **1968**, *7*, 209; A. Schmidpeter, H. Roßknecht, *Angew. Chem.* **1969**, *81*, 572–573; *Angew. Chem. Int. Ed. Engl.* **1969**, *8*, 614; *Z. Naturforsch., Part B*, **1971**, *26*, 81–82. — [18d] H. Binder, R. Fischer, *Chem. Ber.* **1974**, *107*, 205–209. — [18e] H. R. O'Neal, R. H. Neilson, *Inorg. Chem.* **1984**, *23*, 1372–1377. — [18f] N. Burford, J. A. C. Clyburne, S. Mason, J. F. Richardson, *Inorg. Chem.* **1993**, *32*, 4988–4989.
- [19] Isomerisation of a P-coordinated (diphosphanylamine)rhenium complex $[(OC)_5RhPPh_2NHPPPh_2]^+$ to its PH tautomer $[(OC)_5RhPPh_2N=P(H)Ph_2]^+$ has been reported: P. Steil, U. Nagel, W. Beck, *J. Organomet. Chem.* **1989**, *366*, 313–331.
- [20] A. Schmidpeter, J. Eberling, H. Stary, C. Weingand, *Z. Anorg. Allg. Chem.* **1972**, *394*, 171–186.
- [21] S. K. Seale, J. L. Atwood, *J. Organomet. Chem.* **1974**, *73*, 27–34.
- [22] R. Hasselbring, H. W. Roesky, A. Heine, D. Stalke, G. M. Sheldrick, *Z. Naturforsch., Part B*, **1993**, *48*, 43–49.
- [23] R. van Bynum, W. E. Hunter, R. D. Rogers, J. L. Atwood, *Inorg. Chem.* **1980**, *19*, 2368–2374.
- [24] J. W. Lauher, R. Hoffmann, *J. Am. Chem. Soc.* **1976**, *98*, 1729–1741.
- [25] G. Alcaraz, A. Bacereido, F. Dahan, G. Bertrand, *J. Am. Chem. Soc.* **1994**, *116*, 1225–1229.
- [26] H. M. Olmstead, P. P. Power, A. Sigel, *Inorg. Chem.* **1988**, *27*, 2045–2047.
- [27] H. Falius, M. Babin, *Z. Anorg. Allg. Chem.* **1976**, *420*, 65–71.
- [28] H. C. Brown, S. U. Kulkarni, *J. Organomet. Chem.* **1979**, *168*, 281–293.
- [29] G. M. Sheldrick, *SHELXTL-Plus*, Siemens Analytical X-Ray Instruments Inc., Madison, Wisconsin, USA, **1989**.
- [30] G. M. Sheldrick, *SHELXTL-93*, University of Göttingen, **1993**.
- [31] N. Walker, D. Stuart, *Acta Crystallogr., Sect. A*, **1983**, *39*, 158–166.
- [32] H. D. Flack, *Acta Crystallogr., Sect. A*, **1983**, *39*, 876–881. [96066]