

Article

Evaluation of Nitrosamide Formation in the Cytochrome P450 Mediated Metabolism of Tobacco-Specific Nitrosamines

Erik S. Carlson, Pramod Upadhyaya, and Stephen S. Hecht

Chem. Res. Toxicol., **Just Accepted Manuscript** • DOI: 10.1021/acs.chemrestox.6b00384 • Publication Date (Web): 16 Nov 2016Downloaded from <http://pubs.acs.org> on November 20, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

Chemical Research in Toxicology is published by the American Chemical Society.

1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society.

However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Evaluation of Nitrosamide Formation in the Cytochrome P450 Mediated Metabolism of
2 Tobacco-Specific Nitrosamines


3
4 Erik S. Carlson,^{†,‡} Pramod Upadhyaya,[†] and Stephen S. Hecht^{†,*}

5
6 [†] Masonic Cancer Center, University of Minnesota, 2231 6th Street SE, 2-210 CCRB, Minneapolis, MN
7 55455, United States

8 [‡] Department of Pharmacology, University of Minnesota Medical School, 321 Church Street SE, 6-120
9 Jackson Hall, Minneapolis, MN 55455

10
11
12
13
14
15
16
17
18
19 *To whom correspondence should be addressed:
20 Masonic Cancer Center, University of Minnesota, 2231 6th Street SE - 2-148 CCRB, Minneapolis, MN 55455, USA.
21 phone: (612) 624-7604 fax: (612) 624-3869 e-mail: hecht002@umn.edu

TOC Graphic:



Abstract:

N'-Nitrosonorcotine (NNN) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) are carcinogenic tobacco-specific nitrosamines believed to play a vital role in the initiation of tobacco-related cancers. To exhibit their carcinogenicity, both NNN and NNK must be metabolically activated by cytochrome P450s, specifically P450 2A6 and P450 2A13, respectively. Prior research has focused on α -hydroxylation, which leads to the formation of several DNA adducts that have been identified and quantified *in vivo*. However, some studies indicate that P450s can retain substrates within their active sites and perform processive oxidation. For nitrosamines, this would oxidize the highly unstable α -hydroxynitrosamines to potentially more stable nitrosamides, which could also alkylate DNA. Thus, we hypothesized that both NNN and NNK are processively oxidized *in vitro* to nitrosamides by P450 2A6 and P450 2A13, respectively. To test this hypothesis, we synthesized the NNN- and NNK-derived nitrosamides, determined their half-lives at pH 7.4 and 37 °C, and monitored for nitrosamide formation in an *in vitro* P450 system with product analysis by LC-ESI⁺-HRMS/MS. Half-lives of the nitrosamides were determined by HPLC-UV and ranged from 7–35 min, which is more than 40 times longer than the corresponding α -hydroxynitrosamines. Incubation of NNN in the P450 2A6 system resulted in the formation of the nitrosamide, *N*'-nitrosonorcotine (NNC) at low levels. Similarly, the nitrosamide 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanedione (CH₂-oxo-NNK), was detected in low amounts in the incubation of NNK with the P450 2A13 system. The other possible NNK-derived nitrosamide, 4-(nitrosoformamido)-1-(3-pyridyl)-1-butanone (CH₃-oxo-NNK), was not observed in the P450 2A13 reactions. CH₂-oxo-NNK readily formed O⁶meGua in reactions with dGuo and calf thymus DNA. These results demonstrate that NNC and CH₂-oxo-NNK are novel metabolites of NNN and NNK, respectively. Though low-forming,

1 their increased stability may allow for mutagenic DNA damage *in vivo*. More broadly, this study
2 provides the first account of a cytochrome-P450 mediated conversion of nitrosamines to
3 nitrosamides which warrants further studies to determine how general this phenomenon is in
4 nitrosamine metabolism.

1 Introduction:

Tobacco use is the leading preventable cause of cancer death in the United States resulting in an estimated 160,000 deaths annually,¹ or 30% of all cancer deaths nationwide.² 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK, **1**) and *N'*-nitrosornicotine (NNN, **2**) are tobacco-specific nitrosamines thought to play a critical role in tobacco carcinogenesis. Both compounds cause tumors in animal models and have been classified as carcinogenic to humans.^{1,3} To exhibit their carcinogenicity, NNK and NNN must first be metabolically activated by cytochrome P450s.⁴⁻⁶ In the case of NNK, P450 2A13-mediated hydroxylation at the methylene or methyl carbon adjacent to the nitrosamino group results in highly unstable α -hydroxy species **3** and **4**^{3,4} (Scheme 1A). Prior work indicates that α -hydroxynitrosamines have half-lives of ~ 5 sec^{7,8}, decomposing to diazohydroxides such as **7** and **8**,⁸ which lose H₂O producing diazonium ions that either hydrolyze to urinary products **10** and **11** or react with DNA and proteins to form adducts⁴ (Scheme 1A). If left unrepaired, DNA adducts can cause mutations in critical oncogenes or tumor suppressor genes, initiating tumor development. Methyl adducts such as O⁶-methylguanine (O⁶meG) are especially tumorigenic in animal models.⁹ For NNN, α -hydroxylation at the 2' or 5' position by P450 2A6 initiates a similar cascade leading to a variety of DNA adducts and urinary products (Scheme 1A). Both NNK- and NNN-derived DNA adducts have been identified and quantified in several animal models and show promise as biomarkers for carcinogen activation in humans.¹⁰⁻¹⁴

While the α -hydroxynitrosamine hypothesis outlined above accounts nicely for many of the metabolic products and DNA adducts produced in the metabolism of NNK, NNN, and other carcinogenic nitrosamines, there are also inconsistencies in this hypothesis.^{15,16} Important among these is the short lifetime of α -hydroxynitrosamines which raises questions about their ability to alkylate nuclear DNA after having been formed in the endoplasmic reticulum. A few studies have shown these intermediates exist long enough for glucuronidation, but measured levels were minor. In addition, it is unclear if this pathway mediates detoxification or intracellular transportation to the nucleus¹⁷⁻¹⁹. Elespuru et al and Guttenplan have explored the hypothesis that the α -hydroxynitrosamines are further oxidized to nitrosamides, which

are also direct DNA alkylating agents, but this alternate hypothesis lacks compelling supportive data^{15,16}. In the study reported here, we explore this hypothesis with respect to the metabolic activation of NNK and NNN. We hypothesize that the α -hydroxynitrosamines could be retained in the active site of P450 2A13 or P450 2A6 and further oxidized to the corresponding nitrosamides: 4-(methylnitrosamino)-1-(3-pyridyl)-1,4-butanedione (CH₂-oxo-NNK, **13**), 4-(nitrosoformamido)-1-(3-pyridyl)-1-butanone (CH₃-oxo-NNK, **14**) from NNK and *N'*-nitrosonorcotinine (NNC, **15**) from NNN.

There is precedent for retention of substrates in P450s leading to processive oxidation. Metabolism of nicotine by P450 2A6 proceeds through a retained iminium ion or hemiaminal intermediate before releasing cotinine as the major metabolite.²⁰ The Guengerich group has shown that P450 2E1 oxidizes ethanol directly to acetic acid with limited substrate dissociation.²¹ Likewise, they showed that formaldehyde and acetaldehyde formed in the metabolism of dimethylnitrosamine and diethylnitrosamine by P450 2A6 are directly oxidized to formic acid and acetic acid, respectively, without release of the intermediate aldehydes.²² They later proposed an alternate route for acid formation via a nitrosamide intermediate; however, it was undetectable.²³

Nitrosamides are recognized as direct-acting carcinogens with common half-lives being on the scale of minutes.²⁴ Extensive research shows that their stability and reactivity is dependent on temperature, steric and electronic factors, and solvent composition.^{25–29} In nucleophilic environments,³⁰ the major products are the corresponding carboxylic acid derivatives and the diazonium species discussed earlier (Scheme 1). Thus, if released into a cell, nitrosamides should not only have better stability for traversing the hydrolytic environment of the cytosol, but also the ability to alkylate DNA. If nitrosamines are oxidized to nitrosamides, this could lead to the identification of new classes of DNA adducts and a better understanding of the mechanisms of nitrosamine carcinogenesis.

In the present study, we examine the hypothesis that NNK and NNN are metabolized to their corresponding nitrosamides. We synthesized these three nitrosamides and evaluated their half-lives and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 major degradation products *in vitro*. We describe our finding that CH₂-oxo-NNK and NNC are minor
2 metabolites of NNK and NNN in *in vitro* assays with human cytochrome P450s. Further evaluation of
3 CH₂-oxo-NNK demonstrated that it methylates dGuo in DNA and is thus potentially mutagenic.
4 Together, this work provides the first account of a nitrosamine being converted metabolically to a
5 nitrosamide and furthers our understanding of the metabolism of NNK and NNN.

6

Experimental Procedures:

Caution: *NNN and NNK are carcinogenic in animal models and are IARC Group 1 carcinogens. All nitrosamides are presumed to be carcinogens based on their structure and reactivity. Handle these in a well-ventilated fume hood with personal protective equipment and extreme care.*

Chemicals and Enzymes:

NNK, NNN, 4-oxo-4-(3-pyridyl)-butanol (keto alcohol, **11**), 5-(3-pyridyl)-2-hydroxytetrahydrofuran (lactol, **12**), O⁶-methylguanine (O⁶meGua), and [CD₃]O⁶meGua were synthesized as previously described.^{31–34} 5-(3-Pyridyl)-2-pyrrolidinone (norcotinine, **28**) was obtained from AKos GmbH (Steinen, Germany). P450 2A6 Baculosomes, regeneration system and reaction buffer were available as a Vivid CYP450 Screening Kit from Life Technologies (Carlsbad, CA). Purified P450 2A13 and P450 reductase were a generous gift from Dr. Sharon Murphy (University of Minnesota). All other chemicals and solvents used were obtained from either Sigma Aldrich (Milwaukee, WI) or Thermo Scientific (Waltham, MA) in reagent grade and used without further purification.

General Synthetic Procedures:

NMR spectra were recorded on a Bruker 500 MHz spectrometer. Chemical shifts are reported as parts per million (ppm). Residual solvent peaks were used as an internal reference for ¹H-NMR (7.26 ppm CDCl₃; 2.50 ppm D₆-DMSO) and ¹³C-NMR (77.2 ppm CDCl₃; 39.5 ppm D₆-DMSO). Peak splitting used the following abbreviations: s = singlet, d = doublet, t = triplet, q = quartet, dd = doublet of doublets, dt = doublet of triplets, ddd = doublet of doublet of doublets, bs = broad singlet, and m = multiplet. High resolution mass spectrometry (HRMS) for selected compounds was performed on an LTQ Orbitrap Velos (Thermo Scientific, Carlsbad, CA) and reported as *m/z*. Thin-layer chromatography (TLC) utilized

1 Polygram pre-coated silica gel TLC plate (40 x 80 mm, 0.2 mm thick) with 254 nm fluorescent indicator.
2
3
4
5 2 TLC plates were visualized with permanganate stain when necessary, otherwise UV lamp irradiation
6
7
8 3 sufficed. Flash chromatography was performed on SiliCycle 60 (70-150) mesh silica gel. Reactions were
9
10 4 performed under an atmosphere of N₂ unless specified otherwise.
11
12
13
14
15

16 **6 Methyl 4-oxo-4-(3-pyridyl)-1-butanoate (18):**
17

18
19 7 Sodium cyanide (0.104 g, 2.12 mmol) was suspended in anhydrous *N,N*-dimethylformamide (DMF, 10
20
21 8 mL) and brought to 35 °C. 3-Pyridinecarboxaldehyde (2.27 g, 21.2 mmol, 2 mL) was added dropwise to
22
23 9 the suspension. After 10 min of stirring, the resulting red solution was treated dropwise with methyl
24
25 10 acrylate (1.90 g, 22.1 mmol, 2 mL). Over 4 h, the solution became increasingly yellow and slightly
26
27 11 viscous. The reaction was quenched with acetic acid (100 µL). The resulting yellow solution was diluted
28
29 12 in CH₂Cl₂ and washed with H₂O and brine. The organic layer was dried over MgSO₄, filtered, and
30
31 13 concentrated *in vacuo* to a crude, yellow solid. Purification by column chromatography (50 to 100%
32
33 14 EtOAc in hexanes) yielded pure product as a white, crystalline solid (3.02 g, 73.8%). ¹H-NMR (500 MHz;
34
35 15 CDCl₃): δ 9.21 (dd, *J* = 2.2, 0.8 Hz, 1H, 2-Py), 8.79 (dd, *J* = 4.8, 1.7 Hz, 1H, 6-Py), 8.25 (ddd, *J* = 8.0,
36
37 16 2.2, 1.8 Hz, 1H, 4-Py), 7.43 (ddd, *J* = 8.0, 4.8, 0.9 Hz, 1H, 5-Py), 3.72 (s, 3H, CH₃), 3.33 (t, *J* = 6.5 Hz,
38
39 17 2H, COCH₂CH₂), 2.80 (t, *J* = 6.5 Hz, 2H, COCH₂CH₂).; ¹³C-NMR (126 MHz; CDCl₃): δ 197.1 (CO),
40
41 18 173.2 (CO₂CH₃), 153.8 (2-Py), 149.8 (5-Py), 135.5 (4-Py), 131.9 (3-Py), 123.8 (5-Py), 52.1 (CH₃), 33.8
42
43 19 (COCH₂CH₂CO₂), 27.9 (COCH₂CH₂CO₂) ppm.
44
45
46
47
48
49
50

51 **21 4-Oxo-4-(3-pyridyl)-butanoic acid (19):**
52

53
54 22 Compound **18** (393 mg, 2.03 mmol) was dissolved in 1N NaOH (4 mL) and stirred for 3 h at room
55
56 23 temperature. The solution went from colorless to yellow. The pH was adjusted to ~5-6 with an equal
57
58
59
60

1 volume of 1N HCl and a precipitate formed. The solid was filtered and dried under vacuum to give a
2 white, crystalline solid (251 mg, 69%).

3 ¹H-NMR (500 MHz; DMSO-d₆): δ 12.18 (s, 1H, COOH), 9.14 (d, *J* = 2.1 Hz, 1H, 2-Py), 8.80 (dd, *J* = 4.8,
4 1.6 Hz, 1H, 6-Py), 8.31 (dd, *J* = 8.0, 1.8 Hz, 1H, 4-Py), 7.57 (dd, *J* = 8.0, 4.8 Hz, 1H, 5-Py), 3.30 (t, *J* =
5 6.3 Hz, 2H, COCH₂), 2.60 (t, *J* = 6.3 Hz, 2H, CH₂COOH). ¹³C-NMR (126 MHz; DMSO): δ 198.1 (CO),
6 173.7 (COOH), 153.5 (2-Py), 149.1 (6-Py), 135.4 (4-Py), 131.7 (3-Py), 123.9 (5-Py), 33.4 (COCH₂), 27.7
7 (CH₂COOH) ppm.

9 **Methyl 4-oxo-4-(3-pyridyl)-butanamide (20a):**

10 A solution of **19** (43.57 mg, 0.243 mmol), methylamine hydrochloride (25.41 mg, 0.376 mmol), and *N*-
11 hydroxysuccinimide (NHS, 45.2mg, 0.393 mmol) in anhydrous dimethylsulfoxide (DMSO, 3 mL) was
12 treated with *N,N*-diisopropylethylamine (DIPEA, 75.7 mg, 0.586 mmol, 102 μL) and 1-ethyl-3-(3-
13 dimethylaminopropyl)carbodiimide (EDAC, 196 mg, 1.03 mmol). The reaction was stirred for 22 h at
14 room temperature before diluting with EtOAc. The mixture was washed with H₂O and brine, dried over
15 MgSO₄, filtered, and concentrated *in vacuo* to yield a crude, pink oil. Purification by column
16 chromatography (5:100 MeOH/CH₂Cl₂) yielded pure product as an off-white solid (32.9 mg, 70%).

17 ¹H-NMR (500 MHz; DMSO-d₆): δ 9.13 (d, *J* = 1.2 Hz, 1H, 2-Py), 8.79 (dd, *J* = 4.7, 1.3 Hz, 1H, 6-Py),
18 8.30 (d, *J* = 7.9 Hz, 1H, 4-Py), 7.84 (s, 1H, NH), 7.57 (dd, *J* = 7.9, 4.7 Hz, 1H, 5-Py), 3.27 (t, *J* = 6.6 Hz,
19 2H, COCH₂), 2.57 (d, *J* = 4.6 Hz, 3H, CH₂CONH), 2.48 (t, *J* = 6.6 Hz, 2H, NHCH₃). ¹³C NMR (126
20 MHz; DMSO): δ 198.6 (CO), 171.3 (CONH), 153.3 (6-Py), 149.1 (2-Py), 135.4 (4-Py), 131.8 (3-Py),
21 123.9 (5-Py), 33.7 (COCH₂), 29.0 (CH₂CONH), 25.5 (NHCH₃) ppm.

1 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanedione (CH₂-oxo-NNK, 13):

2 A solution of **20a** (31.4mg, 0.163 mmol) in a 5:1 mixture of acetic anhydride and acetic acid (6 mL) was
3 brought to 0 °C. To this was added NaNO₂ (30.7 mg, 0.445 mmol) all at once. After 4 h, the mixture was
4 poured onto ice-cold H₂O. The aqueous mixture was extracted with CH₂Cl₂. The pooled organics were
5 dried over MgSO₄, filtered, and concentrated *in vacuo* to yield a crude yellow oil. Purification by column
6 chromatography on silica gel (100% EtOAc) yielded pure product as a bright, yellow oil (28.8 mg, 65%).
7 ¹H-NMR (500 MHz; CDCl₃): δ 9.27 (s, 1H, 2-Py), 8.83 (d, *J* = 4.1 Hz, 1H, 6-Py), 8.31 (d, *J* = 7.9 Hz, 1H,
8 4-Py), 7.48 (dd, *J* = 7.8, 4.9 Hz, 1H, 5-Py), 3.67 (t, *J* = 6.0 Hz, 2H, COCH₂), 3.55 (t, *J* = 6.0 Hz, 2H,
9 CH₂CON), 3.14 (s, 3H, CH₃). ¹³C NMR (126 MHz; CDCl₃): δ 196.7 (CO), 175.9 (CONNO), 153.5 (6-
10 Py), 149.4 (2-Py), 135.6 (4-Py), 131.9 (3-Py), 123.8 (5-Py), 33.0 (CH₂CONNO), 28.7 (COCH₂), 25.9
11 (NCH₃) ppm. HRMS Calc: 222.08732, Found: 222.08719

13 CH₂-oxo-NNK from 5'-Hydroxycotinine:

14 5'-Hydroxycotinine (54.23 mg, 0.282 mmol) in CH₂Cl₂ (25 mL) was brought to 0 °C and treated with *p*-
15 toluenesulfonic acid (111 mg, 0.584 mmol). After 5 min of stirring, NaNO₂ (151.2 mg, 2.2 mmol) was
16 added. Stirring was continued for 3 h before pouring onto ice-cold H₂O. The organic layer was separated
17 and the aqueous extracted with CH₂Cl₂. The pooled organics were dried over MgSO₄, filtered, and
18 concentrated *in vacuo* to give a crude orange oil. The product was purified by HPLC using a 150 x 4.6
19 mm, 5 μm, Kinetix HILIC column (Phenomenex) with 1:1 hexanes/CHCl₃ and isopropanol as mobile
20 phases. The gradient was 5% to 20% isopropanol over 10 min at 1 mL/min. The product eluted at ~4 min.
21 The purified product was a bright, yellow oil (4.98 mg, 8%).

1 N'-Nitrosonorcotine (NNC, 15):

A solution of norcotine (31.4 mg, 0.163 mmol) in a 5:1 mixture of acetic anhydride and acetic acid (6 mL) was brought to 0 °C. To this was added NaNO₂ (30.7 mg, 0.445 mmol) all at once. After 4 h, the mixture was poured onto ice-cold H₂O. The aqueous mixture was extracted with CH₂Cl₂. The pooled organics were dried over MgSO₄, filtered, and concentrated *in vacuo* to yield a crude yellow oil. Purification by column chromatography (100% EtOAc) yielded pure product as a bright, yellow oil (92%). ¹H-NMR (500 MHz; CDCl₃): δ 8.56 (d, *J* = 4.3 Hz, 1H, 6-Py), 8.46 (d, *J* = 1.8 Hz, 1H, 2-Py), 7.36 (d, *J* = 8.0 Hz, 1H, 4-Py), 7.30-7.28 (m, 1H, 5-Py), 5.29 (dd, *J* = 9.1, 3.1 Hz, 1H, NCH), 2.97 (dt, *J* = 18.6, 9.4 Hz, 1H, COCH₂), 2.87 (ddd, *J* = 18.5, 9.4, 4.0 Hz, 1H, COCH₂'), 2.66 (dq, *J* = 13.4, 9.4 Hz, 1H, CHCH₂), 2.15-2.09 (m, 2H, CHCH₂') ppm; ¹³C-NMR (126 MHz; CDCl₃): δ 172.7 (CO), 149.4 (6-Py), 147.1 (2-Py), 134.1 (3-Py), 132.7 (4-Py), 123.8 (5-Py), 55.7 (NCH), 26.2 (COCH₂), 22.2 (CHCH₂) ppm. HRMS Calc: 192.07675, Found: 192.07670

2-(3-Pyridyl)-1,3-dithiane (21):

A solution of 3-pyridylcarboxaldehyde (114 mg, 1.06 mmol, 100 μL) and 1,3-propanedithiol (162 mg, 1.50 mmol, 150 μL) in anhydrous tetrahydrofuran (THF, 5 mL) was treated with BF₃•Et₂O (173 mg, 1.22 mmol 300 μL) dropwise at room temperature. The mixture was then heated to reflux at 80 °C and stirred for 24 h before quenching with sat'd NaHCO₃ solution. The aqueous phase was extracted several times with CH₂Cl₂. The pooled organic layers were washed with brine, dried over MgSO₄, filtered, and evaporated *in vacuo* to give crude yellow crystals. Purification by column chromatography (hexanes/EtOAc 1:1) yielded the product as a fine, white powder (199.4 mg, >95%).

¹H-NMR (500 MHz; CDCl₃): δ 8.72 (d, *J* = 2.0 Hz, 1H, 2-Py), 8.57 (dd, *J* = 4.9, 1.5 Hz, 1H, 6-Py), 7.85 (dt, *J* = 7.9, 2.0 Hz, 1H, 4-Py), 7.31 (dd, *J* = 7.9, 4.9 Hz, 1H, 5-Py), 5.21 (s, 1H, S₂CH), 3.10 (ddd, *J* =

1
2
3 1 14.6, 12.3, 2.4 Hz, 2H, SCH₂CH₂ -ax), 2.96 (dt, *J* = 14.0, 3.8 Hz, 2H, SCH₂CH₂ -eq), 2.22 (dtt, *J* = 14.2,
4
5 2 4.6, 2.4 Hz, 1H, SCH₂CH₂ -ax), 2.02-1.93 (m, 1H, SCH₂CH₂ -eq) ppm; ¹³C NMR (126 MHz; CDCl₃): δ
6
7 3 150.1 (6-Py), 149.5 (2-Py), 135.7 (4-Py), 135.4 (3-Py), 124.0 (5-Py), 48.8 (S₂CH), 32.3 (SCH₂CH₂), 25.2
8
9 4 (SCH₂CH₂) ppm.

5

6 ***tert*-Butyl 3-(2-(3-pyridyl)-1,3-dithianyl)-1-propylcarbamate (22):**

7 A solution of **21** (222 mg, 1.13 mmol) and tetramethylethylenediamine (TMEDA, 131.8 mg, 1.13 mmol,
8 170 μL) in anhydrous THF (6 mL) was cooled to -78 °C and treated with *n*-BuLi in hexanes dropwise
9 (1.28 mmol, 800 μL). The resulting dark red solution was stirred at -78 °C for 30 min before dropwise
10 addition of **27** in THF (360 mg, 1.26 mmol, 3 mL). The mixture was stirred at -78 °C for 2 h before
11 allowing the bath to come to room temperature. After 14 h of stirring, the reaction was quenched with
12 H₂O. The aqueous phase was extracted several times with EtOAc. The pooled organics were dried over
13 MgSO₄, filtered, and concentrated *in vacuo* to give a crude, yellow oil. Purification by column
14 chromatography (hexanes/EtOAc 1:1) yielded the product as yellow crystals (290 mg, 72.5%).

15 ¹H-NMR (500 MHz; CDCl₃): δ 9.13 (d, *J* = 2.1 Hz, 1H, 2-Py), 8.52 (dd, *J* = 4.7, 1.6 Hz, 1H, 6-Py), 8.20
16 (ddd, *J* = 8.1, 2.4, 1.6 Hz, 1H, 4-Py), 7.32 (ddd, *J* = 8.1, 4.7, 0.6 Hz, 1H, 5-Py), 4.41 (bs, 1H, NH), 3.03-
17 3.00 (m, 2H, NHCH₂), 2.72-2.61 (m, 4H, SCH₂CH₂), 2.03-2.00 (m, 2H, CCH₂), 1.99-1.91 (m, 2H,
18 SCH₂CH₂), 1.48-1.42 (m, 2H, CH₂CH₂CH₂), 1.40 (s, 9H, C(CH₃)₃) ppm; ¹³C NMR (126 MHz; CDCl₃): δ
19 155.9 (NHCO), 150.8 (2-Py), 148.4 (6-Py), 137.6 (3-Py), 136.8 (4-Py), 123.4 (5-Py), 79.4 (C(CH₃)₃),
20 56.5 (SCS), 42.5 (CCH₂), 40.4 (NHCH₂), 28.5 (CH₃), 27.6 (SCH₂CH₂), 25.1 (SCH₂CH₂), 24.7
21 (CH₂CH₂CH₂) ppm.

22

23

1

3-(2-(3-Pyridyl)-1,3-dithianyl)-1-propylformamide (23):

A solution of **22** (87.6 mg, 0.247 mmol) in CH₂Cl₂ (3 mL) was treated with trifluoroacetic acid (TFA, 1.49 g, 13.1 mmol, 1 mL), which resulted in gas evolution. After 3 h, the solvent was evaporated *in vacuo* to remove excess TFA. The resulting oil was reconstituted in CH₂Cl₂ and washed with sat'd. NaHCO₃ solution and brine. The organic layer was dried over MgSO₄, filtered, and concentrated *in vacuo* to give a yellow oil. This was dissolved in MeOH (5 mL) and treated with triethylamine (29.04 mg, 0.288 mmol, 40 µL) and methyl formate (95.7 mg, 1.60 mmol, 110 µL). The flask was sealed and heated to 55 °C with stirring. After 4 h, the mixture was concentrated *in vacuo* to yield a yellow oil. ¹H-NMR indicated the product was a 9:1 mixture of *cis*- and *trans*-formamide isomers. Purity was sufficient to carry forward without column purification.

¹H-NMR (500 MHz; CDCl₃): δ 9.09 (d, *J* = 2.5 Hz, *trans*-2-Py), 9.08 (d, *J* = 2.1 Hz, 1H, *cis*-2-Py), 8.49 (dd, *J* = 4.8, 1.6 Hz, *trans*-6-Py), 8.48 (dd, *J* = 4.7, 1.5 Hz, 1H, *cis*-6-Py), 8.19-8.16 (m, 1H, 4-Py), 8.07 (d, *J* = 1.1 Hz, 1H, *cis*-NHCHO), 7.92 (d, *J* = 11.9 Hz, *trans*-NHCHO), 7.32-7.29 (m, 1H, 5-Py), 5.94 (s, 1H, *cis*-NHCHO), 5.84 (s, *trans*-NHCHO), 3.18 (q, *J* = 6.7 Hz, 2H, *cis*-CH₂NH), 3.10 (q, *J* = 6.8 Hz, *trans*-CH₂NH), 2.63 (m, 4H, SCH₂), 2.02-1.99 (m, 2H, CCH₂), 1.98-1.85 (m, 2H, SCH₂CH₂), 1.52-1.46 (m, 2H, CH₂CH₂NH). ¹³C-NMR (126 MHz; CDCl₃): δ 164.4 (*trans*-CHO), 161.2 (*cis*-CHO), 150.5 (2-Py), 148.37 (*trans*-6-Py), 148.25 (*cis*-6-Py), 137.45 (*cis*-3-Py), 137.35 (*trans*-3-Py), 136.73 (*cis*-4-Py), 136.67 (*trans*-4-Py), 123.33 (*trans*-5-Py), 123.29 (*cis*-5-Py), 56.28 (*cis*-SCS), 56.17 (*trans*-SCS), 42.3 (*cis*-CCH₂), 42.0 (*trans*-CCH₂), 41.4 (*trans*-NHCH₂), 37.6 (*cis*-NHCH₂), 27.5 (SCH₂), 25.7 (*trans*-CH₂CH₂CH₂), 24.85 (*cis*-SCH₂CH₂), 24.79 (*trans*-SCH₂CH₂), 24.1 (*cis*-CH₂CH₂CH₂) ppm.

22

4-(Formamido)-1-(3-pyridyl)-1-butanone (24):

1 *N*-Chlorosuccinimide (NCS, 101.8 mg, 0.762 mmol) and AgNO₃ (168.5 mg, 0.992 mmol) were
2 suspended in 1:1 MeCN/H₂O (1 mL) and cooled to 0 °C. To this was added a solution of **23** in MeCN
3 (0.247 mmol, 1.5 mL), which resulted in immediate precipitate formation. After 30 min, the reaction was
4 quenched with sat. Na₂SO₃, sat'd NaHCO₃, and brine solutions in succession (1 mL each). The mixture
5 was filtered and extracted with CH₂Cl₂. The pooled organics were dried over MgSO₄, filtered, and
6 concentrated *in vacuo* to give a crude solid. Purification by column chromatography on silica gel (6%
7 MeOH in CHCl₃) yielded pure product as a white solid (20.5 mg, 43% over three steps). ¹H-NMR (500
8 MHz; CDCl₃): δ 9.15 (dd, *J* = 2.2, 0.7 Hz, 1H, 2-Py), 8.78 (dd, *J* = 5.5, 1.7 Hz, 1H, *trans*-6-Py), 8.77 (dd,
9 *J* = 4.8, 1.7 Hz, 1H, *cis*-6-Py), 8.22 (dt, *J* = 8.0, 2.1 Hz, 1H, *trans*-4-Py), 8.21 (dt, *J* = 8.0, 2.0 Hz, 1H, *cis*-
10 4-Py), 8.18 (s, 1H, *cis*-CHO), 8.06 (d, *J* = 11.9 Hz, *trans*-CHO), 7.42 (m, 1H, 5-Py), 5.96 (s, 1H,
11 *NHCHO*), 3.41 (q, *J* = 6.6 Hz, 2H, *cis*-CH₂NH), 3.36 (q, *J* = 6.8 Hz, *trans*-CH₂NH), 3.08 (t, *J* = 6.9 Hz,
12 2H, *cis*-COCH₂), 3.07 (t, *J* = 6.8 Hz, *trans*-COCH₂), 2.01 (quintet, *J* = 6.9 Hz, 2H, *cis*-CH₂CH₂CH₂), 2.01
13 (quintet, *J* = 6.9 Hz, *trans*-CH₂CH₂CH₂); ¹³C-NMR (126 MHz; CDCl₃): δ 198.6 (CO), 161.5 (CHO),
14 153.8 (6-Py), 149.7 (2-Py), 135.5 (4-Py), 132.0 (3-Py), 123.8 (5-Py), 37.8 (CH₂NH), 36.3 (COCH₂), 23.5
15 (CH₂CH₂CH₂) ppm.

16
17 ***tert*-Butyl 3-hydroxypropyl-1-carbamate (26):**

18 A solution of 3-amino-1-propanol (4.94 g, 65.8 mmol, **25**) and triethylamine (7.26 g, 71.7 mmol, 10 mL)
19 in CH₂Cl₂ (175 mL) was treated with Boc anhydride (16.15 g, 74.0 mmol, 17 mL) dropwise, which
20 resulted in vigorous gas evolution. Once bubbling ceased, the reaction was quenched with sat'd NH₄Cl.
21 The organic layer was collected and the aqueous layer was further extracted with CH₂Cl₂. The pooled
22 organics were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. NMR indicated
23 the product was sufficiently pure to be brought directly to the next step.

¹H-NMR (500 MHz; CDCl₃): δ 4.84 (bs, 1H, OH), 3.67 (q, *J* = 5.5 Hz, 2H, HOCH₂), 3.30 (d, *J* = 5.7 Hz, 2H, CH₂NH), 3.07 (bs, 1H, NH), 1.68 (quintet, *J* = 5.7 Hz, 2H, CH₂CH₂CH₂), 1.46 (s, 9H, t-Bu) ppm. ¹³C NMR (126 MHz; CDCl₃): δ 157.2 (CONH₂), 79.6 (C(CH₃)₃), 59.2 (HOCH₂), 36.9 (CH₂NH), 32.9 (CH₂CH₂CH₂), 28.4 (t-Bu) ppm.

tert-Butyl 3-iodopropyl-1-carbamate (27):

Imidazole (Im., 5.42 g, 79.7 mmol) and triphenylphosphine (20.72 g, 79.0 mmol) were dissolved in CH₂Cl₂ (250 mL) and brought to 0 °C. To this was added I₂ (20.50 g, 80.8 mmol) scoopwise, resulting in a dark orange solution. After 20 min of stirring, **26** in CH₂Cl₂ (11.5 g, 65.7 mmol, 50 mL) was added and the solution was allowed to come to room temperature. After 22 h of stirring, the mixture was filtered over Celite and washed with 5% Na₂S₂O₃. The organics were dried over MgSO₄, filtered, and concentrated *in vacuo* to give a yellow solid. Purification by column chromatography (hexanes/EtOAc 4:1) gave pure product as a light yellow solid (83.2% over 2 steps). ¹H-NMR (500 MHz; CDCl₃): δ 4.63 (bs, 1H, NH), 3.19 (m, 4H, NHCH₂/ICH₂), 2.00 (quintet, *J* = 6.6 Hz, 2H, CH₂CH₂CH₂), 1.44 (s, 9H, t-Bu) ppm. ¹³C NMR (126 MHz; CDCl₃): δ 155.9 (CONH), 79.4 (C(CH₃)₃), 41.0 (CH₂NH), 33.4 (CH₂), 28.4 (C(CH₃)₃), 3.1 (ICH₂) ppm.

4-(Nitrosoformamido)-1-(3-pyridyl)-1-butanone (CH₃-oxo-NNK, **14)**

A solution of **24** (31.4mg, 0.163 mmol) in a 5:1 mixture of acetic anhydride and acetic acid (6 mL) was brought to 0 °C. To this was added NaNO₂ (30.7mg, 0.445 mmol) all at once. After 4 h, the mixture was poured onto ice-cold H₂O. The aqueous mixture was extracted with CH₂Cl₂. The pooled organics were dried over MgSO₄, filtered, and concentrated *in vacuo* to yield a crude yellow oil. Purification by column chromatography (100% EtOAc) yielded pure product as a bright, yellow oil (28.8mg, 80%). ¹H-NMR

(500 MHz; CDCl₃): δ 10.00 (s, 1H, CHO), 9.13 (d, J = 1.1 Hz, 1H, 2-Py), 8.80 (dd, J = 4.8, 1.1 Hz, 1H, 6-Py), 8.24 (dd, J = 8.0, 1.6 Hz, 1H, 4-Py), 7.47 (dd, J = 8.0, 4.9 Hz, 1H, 5-Py), 4.11 (q, J = 7.1 Hz, 1H, NCH₂), 2.92 (t, J = 6.9 Hz, 2H, COCH₂), 1.91 (quintet, J = 6.9 Hz, 2H, CH₂CH₂CH₂) ppm. ¹³C NMR (126 MHz; CDCl₃): δ 196.8 (CO), 168.5 (CHO), 153.1 (6-Py), 149.0 (2-Py), 136.0 (4-Py), 132.2 (3-Py), 124.1 (5-Py), 37.6 (NCH₂), 35.8 (COCH₂), 20.9 (CH₂CH₂CH₂) ppm; HRMS Calc: 222.08732, Found: 222.08719

9 Determination of $t_{1/2}$ of Nitrosamides:

HPLC-UV analysis was performed using a 250 x 4.6 mm Gemini-NX C18 column (Phenomenex, Torrance, CA) with the following solvent gradients for the analyses indicated below: (1) isocratic for 5 min at 4% B followed sequentially by a linear gradient to 12% B over 15 min, a 10 min hold, a linear gradient to 30% B over 10 min, and a final linear gradient to 40% B over 2 min; (2) isocratic for 5 min at 12% B followed sequentially by a linear gradient to 30% B over 10 min, a linear gradient to 40% B over 15 min, and a final linear gradient to 70% B over 2 min. In both systems, solvent A was 15 mM NH₄OAc and solvent B was methanol.

For $t_{1/2}$ determination, an aliquot of NNC (180 nmol) in CH₂Cl₂ was dried and reconstituted in 30 μ L of 0.5X P450 2A6 Reaction Buffer (Life Technologies) and incubated for 0 to 30 min at 37 °C. After the desired incubation time, 5 μ L of sample was analyzed by HPLC using gradient 1. A₂₅₄ was monitored and the peaks for NNC and its decomposition products were integrated. NNC eluted at 36.0 min. Peak area for NNC was fit to a single-order exponential plot while using the 0 min incubation peak area as a normalizing factor. The analysis was similarly performed for CH₂-oxo-NNK and CH₃-oxo-NNK, except HPLC gradient 2 was used. CH₂-oxo-NNK and CH₃-oxo-NNK eluted at 33.0 and 29.5 min, respectively.

Decomposition products were identified by retention time comparisons and co-injection with synthetic standards.

***In vitro* detection of CH₂-oxo-NNK using P450 2A13:**

Incubations with P450 2A13 were performed as previously reported.³⁵ Briefly, purified P450 2A13 and cytochrome P450 reductase were reconstituted with dilauroylphosphatidylcholine (DLPC, Sigma Aldrich) for 45 min on ice before diluting with Tris buffer to give a final concentration of 1 μ M P450 2A13, 2 μ M P450 reductase, 0.1 μ g/ μ L DLPC, and 50 mM Tris, pH = 7.4. To initiate the reaction, an aliquot of this (containing 5 pmol P450) was added to a Tris-buffered solution of NNK (4 μ M) and NADPH (0.2 mM). Final reaction volumes were always 100 μ L. The mixture was brought to 37 °C for 1–60 min before quenching with 10 μ L of both Ba(OH)₂ and ZnSO₄. After centrifuging the sample at 8000 g for 4 min, the supernatant was collected and 2 μ L were immediately analyzed by liquid chromatography-positive nanoelectrospray-ionization high-resolution tandem mass spectrometry (LC-NSI⁺-HRMS/MS) with an LTQ Orbitrap Velos (Thermo Scientific, Carlsbad, CA). LC employed a hand-packed, Luna C18 (5 μ m), 100 mm x 75 μ m, 15 μ m orifice capillary column with a multi-step gradient. Initially, 5% B at 1 μ L/min from 0–5 min was used to load the sample. Afterwards, the flow rate was dropped to 0.3 μ L/min and a linear gradient was started from 5% to 20% B over 4 min, followed by a ramp to 55% B over 10 min, and re-equilibration, where solvent A was 5 mM NH₄OAc and solvent B was acetonitrile. CH₂-oxo-NNK and CH₃-oxo-NNK were monitored by both full scan and MS² fragmentation. Full scan was performed at a resolution of 60,000 and the accurate parent mass of both nitrosamides (m/z = 222.08719) was extracted at a mass tolerance of 5 ppm. For MS² fragmentation, parent ions were isolated (2.0 amu isolation width)

1 and fragmented by collision-induced dissociation (CID) with a collision energy of 25 eV, resolution of
2 15,000, and scan time of 30 ms. Accurate product ion masses from characteristic transitions for CH₂-oxo-
3 NNK (m/z 222 \rightarrow m/z 180.06542, -H₃CNNO +OH), CH₃-oxo-NNK (m/z 222 \rightarrow m/z 106.02852), NNK
4 (m/z 208 \rightarrow m/z 178.11002), and keto alcohol **11** (m/z 166 \rightarrow m/z 148.07564) were extracted at a mass
5 tolerance of 5 ppm.

6 7 8 ***In vitro* detection of NNC using P450 2A6:**

9 Incubations with P450 2A6 were performed as described by the manufacturer with modifications.³⁶ After
10 thawing the P450 2A6 Baculosomes and Vivid-NADPH-Regeneration System (Life Technologies) on
11 ice, aliquots were combined and diluted 1:10 and 1:50, respectively, with 0.5X Vivid Reaction Buffer
12 (Life Technologies). For each incubation, an aliquot of the combined-enzyme system (containing 5 pmol
13 P450) was added to a 0.5X Reaction-Buffered solution of NNN (4 μ M) and this new mixture was pre-
14 incubated for 2 min at 37 °C. To initiate the reaction, an aliquot of NADP⁺ (containing 3 nmol) was
15 added. Final reaction volumes were 100 μ L. The incubation and work-up were as described earlier for
16 NNK-P450 2A13 incubations. NNC detection was performed by adapting the NNK-P450 2A13 LC-NSI⁺-
17 HRMS/MS method described above to the accurate parent mass of NNC (m/z 192.07670) in full scan.
18 Likewise, the MS² analysis was used to monitor for the accurate product ion masses from characteristic
19 transitions of NNC (m/z 192 \rightarrow m/z 134.04739, 162.07874), NNN (m/z 178 \rightarrow m/z 148.09941), and lactol
20 **12** (m/z 166 \rightarrow m/z 148.07571).

21 22 ***in vitro* Methylation of dGuo by CH₂-oxo-NNK:**

1 A solution of CH₂-oxo-NNK in CH₂Cl₂ was dried under a stream of N₂ and reconstituted in a phosphate-
2 buffered solution of dGuo (4.34 mM dGuo, 25 mM NaHPO₄, pH = 7.4) so that the molar ratio of CH₂-
3 oxo-NNK to dGuo was 1:1. This was brought to 37 °C and incubated for 18 h. To assess methylation, 200
4 fmol of [CD₃]O⁶meGua was added as internal standard. Samples were brought up to 1 mL with 0.1N HCl
5 and incubated at 90 °C for 30 min. After cooling on ice, the samples were neutralized with 1.0N NaOH
6 and purified by solid-phase extraction (Strata-X polymeric reversed phase, 30 mg, Phenomenex,
7 Torrance, CA). Before sample addition, the cartridge was activated using 1 mL each of MeOH and H₂O.
8 After sample addition, the cartridge was washed with 1 mL of both H₂O and 10% MeOH. The sample
9 was eluted and collected with 1 mL of MeOH. The collected fraction was evaporated to dryness in a
10 Speedvac. The residue was reconstituted in 30 µL of H₂O and analyzed by LC-MS/MS.³⁷

11 We used a well-established liquid chromatography-positive electrospray ionization-tandem mass
12 spectrometry (LC-ESI⁺-MS/MS) method. A 0.5 x 150 mm Zorbax SB-C18, 5 µm column (Agilent, Santa
13 Clara, CA) was eluted with a multi-step gradient and flow rate of 10 µL/min. After a linear gradient from
14 5% to 10% B over 10 min, the eluant was brought to 40% B over 5 min, followed by a wash at 90% B
15 and re-equilibration, where solvent A was 15mM NH₄OAc and solvent B was methanol. MS was
16 performed on a TSQ Vantage triple quadrupole mass analyzer (Thermo Scientific). The SRM transitions
17 were m/z 166.1 → m/z 149.1 and m/z 166.1 → m/z 124.1 for O⁶meGua and m/z 169.1 → m/z 152.1 for
18 [CD₃]O⁶meGua using a collision energy of 30 eV and a 0.2 amu scan width.

19 20 ***in vitro* Methylation of calf thymus DNA by CH₂-oxo-NNK:**

21 A solution of CH₂-oxo-NNK was dried under a stream of N₂ and reconstituted in a phosphate-buffered
22 solution of calf thymus DNA so that the ratio was 3 nmol CH₂-oxo-NNK : 1 µg DNA. This was brought to
23 37 °C and incubated for 18 h. The aqueous sample was extracted twice with equal volumes of
24 CHCl₃:isoamyl alcohol (24:1). The DNA was precipitated by addition of an equal volume of isopropanol

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 and gentle shaking. Isolated DNA was washed with 500 μ L of 70% EtOH and 100% EtOH, and dried
2 under N₂. To assess methylation, isolated DNA was dissolved in 100 μ L of sodium phosphate buffer (25
3 mM, pH = 7.4) and 200 fmol of [CD₃]O⁶meGua was added as internal standard. The samples were then
4 processed and analyzed as described above for dGuo methylation.
5

Results:

Synthesis of Nitrosamides:

Retrosynthetic analysis identified 3-pyridinecarboxaldehyde (**17**) as a common precursor for both NNK-derived nitrosamides (**Scheme 2**). The synthesis of CH₂-oxo-NNK started with the formation of keto ester **18** by using the Stetter reaction³⁸ to couple aldehyde **17** with methyl acrylate, followed by hydrolysis of **18** to keto acid **19**. This method is a convenient alternative to the more commonly used routes to this compound.^{39,40} Keto acid **19** was coupled to methylamine using 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDAC) and *N*-hydroxysuccinimide (NHS) in dimethylsulfoxide (DMSO), resulting in the open-chain and lactam conformers of **20** in a 2:1 ratio, respectively. Nitrosation of the open-chain conformer produced CH₂-oxo-NNK (**13**) as a single rotamer.⁴¹ Nitrosation of the lactam conformer also produced CH₂-oxo-NNK; however, new conditions using a strong acid catalyst were required. These conditions also degraded the product which limited the isolatable yield (<10%). Ultimately, only the open-chain route was synthetically useful for CH₂-oxo-NNK production.

Synthesis of CH₃-oxo-NNK (**Scheme 2**) started with protection of **17** with 1,3-propanedithiol in nearly quantitative yield to give **21**. This was coupled to **27** by classic umpolung chemistry⁴² to give **22** in excellent yield. Compound **27** was prepared in two steps from **25** on a multi-gram scale by first Boc-protecting the amine and then converting the alcohol to an iodide using a modified Appel reaction.⁴³ After Boc removal from **22** and *N*-formylation to achieve **23**, the dithiane group was oxidatively removed⁴⁴ to produce **24** in 43% yield over 3 steps. To complete the synthesis, **24** was nitrosated to give CH₃-oxo-NNK (**14**) in excellent yield.

NNC (**15**) was prepared by nitrosation of norcotinine (**28**) in 92% yield. The three nitrosamides were stored in CH₂Cl₂ at 4 °C. They were stable for at least three months under these conditions.

1 Attempts to store these compounds neat or in H₂O-miscible solvents (MeOH, MeCN, acetone, etc.)
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Attempts to store these compounds neat or in H₂O-miscible solvents (MeOH, MeCN, acetone, etc.)
2 resulted in decomposition.
3

4 **Stability of Nitrosamides**

5 The stabilities of the nitrosamides were determined at pH 7.4, 37 °C, in buffers to be used in our
6 P450 assays. Reactions were followed by HPLC and major products were identified (Figure S1,
7 Supporting Information). Decay curves for each nitrosamide are shown in Figure 1. The half-lives of
8 CH₂-oxo-NNK and CH₃-oxo-NNK were 35.5 min and 6.7 min, respectively. The half-life of NNC under
9 these conditions was 12.3 min. The major product in each case was that expected by nitrosamide
10 hydrolysis, namely keto acid **19** from CH₂-oxo-NNK, keto alcohol **11** from CH₃-oxo-NNK, and hydroxy
11 acid **29** from NNC (Scheme 3).

13 ***In vitro* cytochrome P450-catalyzed metabolism of NNK to CH₂-oxo-NNK:**

14 With synthetic nitrosamide standards in hand and an understanding of their stability, we designed
15 an assay to detect their formation by P450-mediated metabolism. P450 2A13 is the most efficient enzyme
16 for α -hydroxylation of NNK and was chosen for this part of the study.⁴⁵ The NNK concentration was set
17 at 4 μ M, the K_m for production of hydroxylated products. Samples were analyzed immediately after work-
18 up to minimize nitrosamide decomposition. We used an LTQ High-Resolution Orbitrap Velos MS system
19 to enhance sensitivity and minimize background noise. We monitored for the accurate masses of the most
20 abundant product ions of CH₂-oxo-NNK and CH₃-oxo-NNK resulting from MS² fragmentation, and also
21 extracted their accurate parent masses in full scan mode. Similar monitoring was performed for NNK and
22 keto alcohol **11** to ensure catalytic turnover as **11** is the most abundant product from methyl
23 hydroxylation.^{4,45}

When NNK was incubated with the relevant enzymes and cofactors, we detected a peak that matched the accurate parent ion mass, accurate product ion mass resulting from MS² of [M+H]⁺ = 222, and the retention time of synthetic CH₂-oxo-NNK (Figure 2). This peak was detectable as early as the 1-min time point, reached its maximum concentration at 5 min, tapered off by 10–30 min, and was nearly undetectable by 60 min (Figure 2C-F). We did not detect CH₃-oxo-NNK at any time point.

The signal for NNK decreased 8-fold while the signal for keto alcohol **11** simultaneously increased 4-fold over the 60-min period (data not shown). This indicates that NNK metabolism was rapid over the assay time period. CH₂-oxo-NNK was a minor metabolite as its signal was >4000-fold less than that of keto alcohol **11**. No metabolites were observed in control incubations lacking enzyme or cofactors (data not shown).

When identical incubations containing CH₂-oxo-NNK (10 nM) were performed, the peak area was 10-fold higher than in unspiked samples, indicating that CH₂-oxo-NNK was recoverable under our conditions. Based on this, we estimate that CH₂-oxo-NNK was produced at concentrations less than 1 nM in our incubations.

***In vitro* cytochrome P450-catalyzed metabolism of NNN to NNC:**

The NNN incubations were performed in essentially the same way as those with NNK except that P450 2A6 was used instead of P450 2A13 as it is the most efficient enzyme for NNN metabolism.⁴⁶ Catalytic turnover was assessed by measuring lactol **12**, because it is the major product from 5'-hydroxylation of NNN (Scheme 1A).⁴⁷ We detected NNC as early as the 1-min time point (Figure 3). The peak matched the synthetic standard with respect to the accurate parent ion mass in full scan, the accurate mass of the most abundant product ions in the MS² of [M + H]⁺ = 192, and retention time.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The NNC signal was maximal at 5 min and was approximately 1000-fold lower in intensity than
2 that of lactol **12** (Scheme 1A). The concentration of NNC at the 5-min time point was estimated to be ~10
3 nM.

5 ***In vitro* Methylation of dGuo and DNA by CH₂-oxo-NNK:**

6 We tested the ability of CH₂-oxo-NNK to methylate DNA by incubating it with a molar
7 equivalent of dGuo and calf thymus DNA in phosphate buffer for 18 h. Levels of *O*⁶-meGua were 61.7
8 and 802 μmol/mg Gua for the dGuo and DNA reactions, respectively, as determined by LC-MS/MS
9 analysis. (Figures S2, Supporting Information). Formation of 7-meGua was also noted, but it was not
10 quantified.

Discussion:

This study presents the first account of nitrosamines being directly converted to nitrosamides by P450 catalysis. This breaks new ground in our knowledge of nitrosamine metabolism and provides an impetus to determine if this phenomenon applies to all nitrosamines. Specifically, we found that CH₂-oxo-NNK and NNC are novel metabolites of P450-mediated oxidation of NNK and NNN, respectively. We did not observe formation of CH₃-oxo-NNK, perhaps due to its short half-life (6.7 min). These novel metabolites also provide a potentially new mechanism for NNK- and NNN-DNA adduct formation (Scheme 1B). It has long been known that the α -hydroxynitrosamine intermediates **3–6** (Scheme 1A) alkylate DNA, but their short lifetimes raise questions regarding their ability to traverse the hydrolytic cytosol. The detected nitrosamides had half-lives of 12–35 min, 100-fold more than those of the α -hydroxynitrosamines.^{7,8} Additionally, we showed that CH₂-oxo-NNK methylates both dGuo and calf thymus DNA (Figure S2, Supplementary Information). Thus, in the case of NNK, it is plausible that CH₂-oxo-NNK could be partially responsible for the methyl adducts previously thought to be formed purely by α -hydroxynitrosamine **3**.

However, we note that both CH₂-oxo-NNK and NNC are quite minor metabolites of NNK and NNN. It was estimated that CH₂-oxo-NNK and NNC form at concentrations of ~1 nM and 10 nM, respectively, while keto alcohol **11** and lactol **12**, the hydrolysis products of α -hydroxyNNK and α -hydroxyNNN, form at levels ~4000-fold higher. Because P450 2A13 and P450 2A6 are the most efficient enzymes for NNK and NNN oxidation and the formation of their known products keto alcohol **11** and lactol **12** (Scheme 1A) was rapid, it is unlikely that the low levels of CH₂-oxo-NNK and NNC result from low catalytic turnover. Likewise, the positive controls indicate that analyte recovery was achievable under our incubation conditions. We noted that formation of both CH₂-oxo-NNK and NNC started at 1 min, peaked at 5 min, and that both were nearly undetectable by 30 min. This may indicate that as metabolism proceeds, newly formed side products and P450-related reactive-oxygen species are eliminating the nitrosamides via secondary reactions at a rate faster than nitrosamide formation.

1 We were not able to quantify nitrosamide formation in these reactions. We initially attempted
2 quantification by HPLC-radioflow techniques, but this approach was not sensitive enough (LOD = 400
3 fmol on column, data not shown). After nitrosamide detection was achieved by LC-NSI⁺-HRMS, we
4 attempted to trap these products with *N*-acetyl-lysine and *N*-acetyl-cysteine. Though trapping was
5 achieved with synthetic standards, this method was unsuccessful in our assay due to low trapping
6 efficiency and low sample recovery after solid phase extraction, which resulted in no analyte detection
7 even with accurate mass detection (data not shown). Therefore, we settled on estimating formation levels
8 by comparing peak areas to those of spiked positive controls.

9 Despite being minor metabolites, the long half-lives and strong DNA-binding properties of the
10 nitrosamides suggest potential biological relevance. However, determining whether it is more important
11 to be low-forming and stable versus high-forming and unstable would require further study. Additionally,
12 though CH₂-oxo-NNK and NNC are formed to low extents, the nitrosamide pathway may be more
13 efficient for other nitrosamines. For example, studies by Chowdhury et al noted considerable, processive
14 conversion of dimethylnitrosamine and diethylnitrosamine to acid byproducts by P450 2A6.²² It was
15 hypothesized that the α -hydroxynitrosamine intermediate decomposed within the active site and the
16 resulting aldehyde was then oxidized to the acid. The nitrosamide hypothesis was also tested, but
17 detection was unsuccessful. Given our results, it is plausible that nitrosamides were readily produced, but
18 instability limited their detection.

19 Our synthesis of each nitrosamide proceeded essentially as expected, except for a few key
20 findings. First, our method for keto acid **19** (Scheme 2), though not entirely novel,⁴⁸ is considerably more
21 convenient than previously reported routes.^{39,40} The two-step process involves milder conditions, gives
22 reproducible yields, and simpler product purification; the latter step provides >99% pure product after
23 only filtration. Next, the conversion of keto acid **19** to keto amide **20** (Scheme 2) was noteworthy because
24 a previous study⁴⁹ reported compound **20** to be in a ring-chain equilibrium heavily favoring the lactam
25 (~6:1). In our hands, the compounds were readily separable on silica gel and showed no isomerization

1 while stored neat at 2–8 °C. They were clearly distinct by NMR (Figure S3, Supporting Information). The
2 open-chain product had two clean triplets integrating to 2H each while these signals collapsed into non-
3 distinct multiplets integrating to 4H in the ring product. Additionally, the methyl resonance in the lactam
4 is a singlet as opposed to a doublet in the open-chain product. In support of the results reported by
5 Nguyen et al,⁴⁹ when we performed this reaction in solvents other than DMSO, the lactam 5'-
6 hydroxycotinine (**20b**, Scheme 2), predominated. Similarly, **20b** was the major product when harsher
7 amide coupling conditions were used, such as *in situ* acid chloride formation by oxalyl chloride or AlMe₃-
8 mediated amide formation^{50,51} from **18**. It is apparent that DMSO and mild coupling conditions favor the
9 open chain conformer.

10 Our nitrosamides were each isolated exclusively as one rotamer. This contrasts to nitrosamines
11 commonly occur as a mixture of both (*E*)- and (*Z*)-isomers. Past studies indicate that the (*E*)-conformer is
12 electronically favored for most nitrosamides.²⁴ Furthermore, rotation to the (*Z*)-conformer is commonly
13 believed to be the rate-limiting step for nitrosamide decomposition by a pericyclic process²⁴ and thus,
14 may not be isolatable.

15 The order of compound stability was CH₂-oxo-NNK > NNC > CH₃-oxo-NNK. This ranking fits
16 with known factors contributing to nitrosamide decomposition.^{24,26,41} In hydrolytic environments,³⁰
17 nitrosamides with bulkier groups adjacent to the carbonyl group are more stable. This suggests that CH₃-
18 oxo-NNK should be the least stable, consistent with our observations. Likewise, bulky groups adjacent to
19 the nitrogen decrease stability. This is consistent with CH₂-oxo-NNK being most stable and NNC being
20 relatively less stable. The decomposition products suggest that the mechanism is primarily hydrolysis.
21 The products shown in Scheme 3B were all either the major or only identified product. However, for
22 NNC and CH₃-oxo-NNK, we did identify a lactone and ester as minor products, respectively (Figure S1,
23 Supporting Information). These presumably result from the extensively studied 1,3-sigmatropic
24 rearrangement mechanism.²⁸ Though this rearrangement is highly favored when nitrosamides are heated

1 in organic solvents, aqueous conditions seem to favor hydrolysis and are most relevant to the *in vivo*
2 situation.

3 With data supporting the formation of CH₂-oxo-NNK, we were interested in testing one of its
4 possible modes of DNA damage: methylation. Methylation was expected since keto acid is the major
5 product of CH₂-oxo-NNK hydrolysis (Scheme 2). This implies that methane diazohydroxide, a known
6 methylating agent, is also released. Our results clearly demonstrated the formation of O⁶meGua in these
7 reactions, indicating that CH₂-oxo-NNK methylates DNA. In addition to the methyl DNA adducts readily
8 formed by CH₂-oxo-NNK, both CH₂-oxo-NNK and NNC could potentially generate a set of novel DNA
9 adducts (Scheme 1B). Further studies are needed to establish the structures, level, and importance of
10 possible adducts derived from the nitrosamide pathway.

11 In summary, we hypothesized that NNK and NNN are metabolized by P450 2A13 and P450 2A6,
12 respectively, to their corresponding nitrosamides *in vitro*. We tested this by synthesizing CH₂-oxo-NNK,
13 CH₃-oxo-NNK, and NNC and evaluating their stability at pH = 7.4 and 37 °C. They were quite stable
14 relative to the corresponding α -hydroxynitrosamines. We then showed that CH₂-oxo-NNK and NNC are
15 novel, though minor, metabolites of NNK and NNN, respectively, in an *in vitro* P450 model. With the
16 knowledge that CH₂-oxo-NNK has a relatively long half-life and methylates DNA, it could potentially
17 play a role in the mechanism of carcinogenesis by NNK. More broadly, this is the first direct evidence for
18 the conversion of nitrosamines to nitrosamides by P450 catalysis and provides rationale for further studies
19 to determine whether this is a general transformation in nitrosamine metabolism.

Funding Information:

This study was supported by grant no. CA-81301 from the U.S. National Cancer Institute

Acknowledgements:

We thank Bob Carlson for editorial assistance, Dr. Peter Villalta and Xun Ming for mass spectrometry assistance in the Analytical Biochemistry Shared Resource of the Masonic Cancer Center, and Dr. Linda Von Weymarn and the laboratory of Dr. Sharon Murphy for allowing use of their facilities and providing enzymes and advice for the cytochrome P450 incubations. We would also like to thank Dr. Adam T. Zarth and Dr. Anna K. Michel for their valuable discussions and input. The Analytical Biochemistry Shared Resource is partially supported by National Cancer Institute Cancer Center Support Grant CA-77598.

Supporting Information:

Select chromatograms for nitrosamide stability and decomposition, LC-MS/MS chromatograms for dGuo and DNA methylation by CH₂-oxo-NNK, and ¹H-NMR spectra for compounds **20a** and **20b**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Abbreviation List:

CH₂-oxo-NNK – 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanedione

CH₃-oxo-NNK – 4-(Nitrosoformamido)-1-(3-pyridyl)-1-butanone

DCM - Dichloromethane

1		
2		
3	1	DIPEA – <i>N,N'</i> -diisopropylethylamine
4		
5	2	DMF – <i>N,N</i> -dimethylformamide
6		
7	3	DMSO - Dimethylsulfoxide
8		
9	4	EDAC – 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
10		
11	5	5-HC – 5'-hydroxycotinine
12		
13	6	Im – imidazole
14		
15	7	Keto alcohol – 4-oxo-4-(3-pyridyl)-butanol
16		
17	8	Lactol – 5-(3-pyridyl)-2-hydroxytetrahydrofuran
18		
19	9	NCS – <i>N'</i> -chlorosuccinimide
20		
21	10	NHS – <i>N'</i> -hydroxysuccinimide
22		
23	11	NNC – <i>N'</i> -nitrosonorcotinine
24		
25	12	NNK – 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone
26		
27	13	NNN – <i>N'</i> -nitrosonornicotine
28		
29	14	O ⁶ meGua – <i>O</i> ⁶ -methylguanine
30		
31	15	[CD ₃]O ⁶ meGua – <i>O</i> ⁶ -([D ₃)methyl])guanine
32		
33	16	TFA – Trifluoroacetic acid
34		
35	17	TMEDA - Tetramethylethylenediamine
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

References:

- (1) United States Department of Health and Human Services, Center for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health (2014) The Health Consequences of Smoking—50 Years of Progress A Report of the Surgeon General. Atlanta.
- (2) American Cancer Society. (2014) Cancer Facts & Figures 2014. *Cancer Facts Fig.* Atlanta.
- (3) International Agency for Research on Cancer. (2007) Monographs on the Evaluation of Carcinogenic Risks to Humans - Volume 89. *Smokeless Tob. Some Tobacco-specific N-Nitrosamines*. Lyon.
- (4) Hecht, S. S. (1998) Biochemistry, Biology, and Carcinogenicity of Tobacco-Specific N-Nitrosamines. *Chem. Res. Toxicol.* 11, 559–603.
- (5) Hecht, S. S., Castonguay, A., Chung, F. L., and Hoffmann, D. (1984) Carcinogenicity and metabolic activation of tobacco-specific nitrosamines: current status and future prospects. *IARC Sci. Publ.* 763–778.
- (6) Hecht, S. S. (1999) DNA adduct formation from tobacco-specific N-nitrosamines. *Mutat. Res. - Fundam. Mol. Mech. Mutagen.* 424, 127–142.
- (7) Revis, C., Fishbein, J. C., Carolina, N., and April, R. V. (1996) Effects of structure on the reactivity of α -hydroxydialkyl nitrosamines in aqueous. *J. Am. Chem. Soc.* 118, 7412–7413.
- (8) Moohizuki, M., Anjo, T., and Okada, M. (1980) Isolation and characterization of N-alkyl-N-(hydroxymethyl)nitrosamines from N-alkyl-N-(hydroperoxymethyl)nitrosamines by deoxygenation. *Tetrahedron Lett.* 21, 3693–3696.
- (9) Peterson, L. A., and Hecht, S. S. (1991) O6-methylguanine is a critical determinant of 4-

1
2
3 1 (methylnitrosamino)-1-(3-pyridyl)-1-butanone tumorigenesis in A/J mouse lung. *Cancer Res.* 51,
4
5 2 5557–5564.
6
7
8
9 3 (10) Balbo, S., Johnson, C. S., Kovi, R. C., James-Yi, S. A., O'Sullivan, M. G., Wang, M., Le, C. T.,
10
11 4 Khariwala, S. S., Upadhyaya, P., and Hecht, S. S. (2014) Carcinogenicity and DNA adduct
12
13 5 formation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and enantiomers of its metabolite 4-
14
15 6 (methylnitrosamino)-1-(3-pyridyl)-1-butanol in F-344 rats. *Carcinogenesis* 35, 2798–2806.
16
17
18 7 (11) Hecht, S. S., Spratt, T. E., and Trushin, N. (1988) Evidence for 4-(3-pyridyl)-4-oxobutylation of
19
20 8 DNA in F344 rats treated with the tobacco-specific nitrosamines 4-(methylnitrosamino)-1-(3-
21
22 9 pyridyl)-1-butanone and *N'*-nitrosonornicotine. *Carcinogenesis* 9, 161–165.
23
24
25 10 (12) Stepanov, I., Muzic, J., Le, C. T., Sebero, E., Villalta, P., Ma, B., Jensen, J., Hatsukami, D., and
26
27 11 Hecht, S. S. (2013) Analysis of 4-hydroxy-1-(3-pyridyl)-1-butanone (HPB)-releasing DNA adducts
28
29 12 in human exfoliated oral mucosa cells by liquid chromatography-electrospray ionization-tandem
30
31 13 mass spectrometry. *Chem. Res. Toxicol.* 26, 37–45.
32
33
34
35 14 (13) Zarth, A. T., Upadhyaya, P., Yang, J., and Hecht, S. S. (2016) DNA adduct formation from
36
37 15 metabolic 5'-hydroxylation of the tobacco-specific carcinogen *N'*-nitrosonornicotine in human
38
39 16 enzyme systems and in rats. *Chem. Res. Toxicol.* 29, 380–389.
40
41
42 17 (14) Zhao, L., Balbo, S., Wang, M., Upadhyaya, P., Khariwala, S. S., Villalta, P. W., and Hecht, S.
43
44 18 (2013) Quantification of pyridyloxobutyl-DNA adducts in tissues of rats treated chronically with
45
46 19 (*R*)-or (*S*)-*N'*-nitrosonornicotine (NNN) in a carcinogenicity study. *Chem. Res. Toxicol.* 26, 1526–
47
48 20 1535.
49
50
51
52 21 (15) Guttenplan, J. B. (1993) Effects of cytosol on mutagenesis induced by *N*-nitrosodimethylamine, *N*-
53
54 22 nitrosomethylurea and α -acetoxy-*N*-nitrosodimethylamine in different strains of *Salmonella*:
55
56 23 evidence for different ultimate mutagens from *N*-nitrosodimethylamine. *Carcinogenesis* 14, 1013–
57
58
59
60

- 1019.
- (16) Elespuru, R. K., Saavedra, J. E., Kovatch, R. M., and Lijinsky, W. (1993) Examination of α -carbonyl derivatives of nitrosodimethylamine and ethylnitrosomethylamine as putative proximate carcinogens. *Carcinogenesis* 14, 1189–1193.
- (17) Murphy, S. E., Spina, D. A., Nunes, M. G., and Pullo, D. A. (1995) Glucuronidation of 4-((hydroxymethyl)nitrosamino)-1-(3-pyridyl)-1-butanone, a metabolically activated form of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, by phenobarbital-treated rats. *Chem. Res. Toxicol.* 8, 772–9.
- (18) Wiench, K., Frei, E., Schroth, P., and Wiessler, M. (1992) 1-c-glucuronidation of N-nitrosodiethylamine and N-nitrosomethyl-N-pentylamine in vivo and in primary hepatocytes from rats pretreated with inducers. *Carcinogenesis* 13, 867–872.
- (19) Wiessler, M., and Rossnagel, G. (1987) Alpha-Glucuronides of N-nitrosomethylbenzylamine, in *The Relevance of N-Nitroso Compounds to Human Cancer, Exposure and Mechanism* (Bartsch, H., O'Neill, I. K., and Schulte-Hermann, R., Eds.), pp 170–172. IARC Scientific Reports 84, Lyon, France.
- (20) von Weymarn, L. B., Retzlaff, C., and Murphy, S. E. (2012) CYP2A6- and CYP2A13-catalyzed metabolism of the nicotine $\Delta^5(1')$ iminium ion. *J. Pharmacol. Exp. Ther.* 343, 307–315.
- (21) Bell-parikh, L. C., and Guengerich, F. P. (1999) Kinetics of cytochrome P450 2E1-catalyzed oxidation of ethanol to acetic acid via acetaldehyde. *J. Biol. Chem.*, 274, 23833–23840.
- (22) Chowdhury, G., Calcutt, M. W., and Peter Guengerich, F. (2010) Oxidation of N-nitrosoalkylamines by human cytochrome p450 2A6: Sequential oxidation to aldehydes and carboxylic acids and analysis of reaction steps. *J. Biol. Chem.* 285, 8031–8044.

- 1
2
3 1 (23) Chowdhury, G., Calcutt, M. W., Nagy, L. D., and Guengerich, F. P. (2012) Oxidation of methyl
4
5 2 and ethyl nitrosamines by cytochromes P450 2E1 and 2B1. *Biochemistry* 51, 9995-10007.
6
7
8 3 (24) Chow, Y. L. (1979) Chemistry of *N*-Nitrosamides and Related *N*-Nitrosamino Acids, in *N*-
9
10 4 *Nitrosamines*, pp 13–37. American Chemical Society, Washington D.C.
11
12
13 5 (25) White, E. H. (1955) The Chemistry of the *N*-Alkyl-*N*-nitrosoamides. II. A New Method for the
14
15 6 Deamination of Aliphatic Amines. *J. Am. Chem. Soc.* 77, 6011–6014.
16
17
18 7 (26) White, E. H., and Dolak, L. A. (1966) *N*-Nitroamides and *N*-nitrocarbamates. IV. Rates of
19
20 8 decomposition. A case of steric acceleration. *J. Am. Chem. Soc.* 88, 3790–3795.
21
22
23 9 (27) Darbeau, R. W., White, E. H., Song, F., Darbeau, N. R., and Chou, J. (1999) A study of essentially
24
25 10 free carbocations derived via diazonium and oxo diazonium ions in the liquid phase. *J. Org. Chem.*
26
27 11 64, 5966–5978.
28
29
30 12 (28) White, E. H., Field, K. W., Hendrickson, W. H., Dzadzic, P., Roswell, D. F., Paik, S., and Mullen,
31
32 13 P. W. (1992) Inert-molecule-separated ion pairs. Stereochemical, I8O, and product studies. *J. Am.*
33
34 14 *Chem. Soc.* 114, 8023–8031.
35
36
37 15 (29) Darbeau, R. W., Pease, R. S., and Gible, R. E. (2001) A study of electronic effects on the kinetics
38
39 16 of thermal deamination of *N*-nitrosoamides. *J. Org. Chem.* 66, 5027–5032.
40
41
42 17 (30) Moss, R. A. (1966) The solvolysis of alkyl diazotates. I. Partition between carbonium ions and
43
44 18 diazoalkanes in aqueous base. *J. Org. Chem.* 31, 1082–1087.
45
46
47 19 (31) Loh, T.-P., Zhou, J.-R., Li, X.-R., and Sim, K.-Y. (1999) A novel reductive aminocyclization for
48
49 20 the syntheses of chiral pyrrolidines: stereoselective syntheses of (S)-nornicotine and 2-(2'-
50
51 21 pyrrolidyl)-pyridines. *Tetrahedron Lett.* 40, 7847–7850.
52
53
54 22 (32) Loozen, H. J. J., Godefroi, E. F., and Besters, J. S. M. M. (1975) A Novel and efficient route to 5-
55
56
57
58
59
60

- 1
2
3 1 arylated gamma-Lactones. *J. Org. Chem.* 40, 892–894.
4
5
6 2 (33) Amin, S., Desai, D., Hecht, S. S., and Hoffmann, D. (1996) Synthesis of tobacco-specific *N*-
7
8 3 nitrosamines and their metabolites and results of related bioassays. *Crit. Rev. Toxicol.* 26, 139–47.
9
10
11 4 (34) Balsiger, R. W., and Montgomery, J. A. (1960) Synthesis of potential anticancer agents. XXV.
12
13 5 Preparation of 6-alkoxy-2-aminopurines. *J. Org. Chem.* 25, 1573–1575.
14
15
16 6 (35) von Weymarn, L. B., Zhang, Q. Y., Ding, X., and Hollenberg, P. F. (2005) Effects of 8-
17
18 7 methoxypsoralen on cytochrome P450 2A13. *Carcinogenesis* 26, 621–629.
19
20
21 8 (36) Wong, H. L., Murphy, S. E., Wang, M., and Hecht, S. S. (2003) Comparative metabolism of *N*-
22
23 9 nitrosopiperidine and *N*-nitrosopyrrolidine by rat liver and esophageal microsomes and cytochrome
24
25 10 P450 2A3. *Carcinogenesis* 24, 291–300.
26
27
28 11 (37) Upadhyaya, P., Lindgren, B. R., and Hecht, S. (2009) Comparative Levels of *O*⁶-Methylguanine ,
29
30 12 pyridyloxobutyl- , and pyridylhydroxybutyl-DNA adducts in lung and liver of rats treated
31
32 13 chronically with the tobacco-specific carcinogen 4- (methylnitrosamino) -1- (3-pyridyl) -1-
33
34 14 butanone. *Drug Metab. Dispos.* 37, 1147–1151.
35
36
37 15 (38) Stetter, H., and Schreckenber, M. (1973) A new method for addition of aldehydes to activated
38
39 16 double bonds. *Angew. Chemie Int. Ed. English* 12, 81.
40
41
42 17 (39) McKennis, H., Turnbull, L. B., Wingfield, H. N., and Dewey, L. J. (1958) Metabolites of nicotine
43
44 18 and a synthesis of nornicotine. *J. Am. Chem. Soc.* 80, 1634–1636.
45
46
47 19 (40) Wingfield Jr., H. N. (1959) 4-(3-Pyridyl)-4-ketobutyric Acid. *J. Org. Chem.* 24, 872–873.
48
49
50 20 (41) White, E. H. (1955) The chemistry of the *N*-alkyl-*N*-nitrosamides. I. Methods of preparation. *J. Am.*
51
52 21 *Chem. Soc.* 77, 6008–6010.
53
54
55 22 (42) Corey, E. J.; Seebach, D. (1965) Carbanions of 1,3-Dithianes. Reagents for C-C bond formation by
56
57
58
59
60

- 1 nucleophilic displacement and carbonyl addition. *Chemie Int. Ed.* 4, 1075–1077.
- 2
- 3 (43) Appel, R. (1975) Tertiary phosphane/tetrachloromethane, a versatile reagent for chlorination,
- 4
- 5 dehydration, and P-N linkage. *Angew. Chemie Int. Ed. English* 14, 801–811.
- 6
- 7
- 8 (44) Corey, E. J., and Erickson, B. W. (1971) Oxidative hydrolysis of 1, 3-dithiane derivatives to
- 9
- 10 carbonyl compounds using *N*-halosuccinimide reagents. *J. Org. Chem.* 36, 3553–3560.
- 11
- 12
- 13 (45) Jalas J.R., Hecht, S. S., and Murphy S.E. (2005) Cytochrome P450 enzymes as catalysts of
- 14
- 15 metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), a tobacco-specific
- 16
- 17 carcinogen. *Chem. Res. Toxicol.* 18, 95–110.
- 18
- 19
- 20 (46) Patten, C. J., Smith, T. J., Friesen, M. J., Tynes, R. E., Yang, C. S., and Murphy, S. E. (1997)
- 21
- 22 Evidence for cytochrome P450 2A6 and 3A4 as major catalysts for *N'*-nitrososornicotine α -
- 23
- 24 hydroxylation by human liver microsomes. *Carcinogenesis* 18, 1623–1630.
- 25
- 26
- 27 (47) Wong, H. L., Murphy, S. E., and Hecht, S. S. (2005) Cytochrome P450 2A-catalyzed metabolic
- 28
- 29 activation of structurally similar carcinogenic nitrosamines: *N'*-Nitrososornicotine enantiomers, *N*-
- 30
- 31 nitrosopiperidine, and *N*-nitrosopyrrolidine. *Chem. Res. Toxicol.* 18, 61–69.
- 32
- 33
- 34 (48) Stetter, H., and SchreckenBerg, M. (1974) A new method for the addition of aldehydes to activated
- 35
- 36 double bonds III. Addition of aromatic and heterocyclic aldehydes to α,β -unsaturated nitriles. *Chem.*
- 37
- 38 *Ber.* 107, 210.
- 39
- 40
- 41 (49) Nguyen, T., Dagne, E., Gruenke, L., Bhargava, H., and Castagnoli, N. J. (1981) The tautomeric
- 42
- 43 structures of 5-hydroxycotinine, a secondary mammalian metabolite of nicotine. *J. Org. Chem.* 46,
- 44
- 45 758–760.
- 46
- 47
- 48 (50) Basha, A., Lipton, M., and Weinreb, S. M. (1977) A mild, general method for conversion of esters
- 49
- 50 to amides. *Tetrahedron Lett.* 18, 4171–4172.
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 1 (51) Lipton, Michael F., Basha, Anwer, Weinreb, S. M. (1979) Conversion of esters to amides with
4
5 2 dimethylaluminum amides: *N,N*-dimethylcyclohexanecarboxamide. *Org. Synth.* 59, 49.
6
7
8
9 3
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends:

2

Figure 1: Stability of CH₂-oxo-NNK (blue diamonds), CH₃-oxo-NNK (green triangles), and NNC (red squares) in reaction buffer at 37 °C. The half-lives were determined to be 35.5, 6.7, and 12.3 min, respectively, by HPLC-UV. Nitrosamide peak areas were normalized to the 0-min peak area and fit to a first-order exponential. Relative amounts of each nitrosamide were determined at each time point in triplicate with error bars denoting the standard deviation.

8

Figure 2: LC-NSI-HRMS chromatograms resulting from the NNK-P450 2A13 incubations. For all sections, the top chromatogram is the accurate parent mass extracted from full scan for CH₂-oxo-NNK and CH₃-oxo-NNK. The middle and bottom chromatogram is the accurate product ion masses extracted from MS² fragmentation for CH₂-oxo-NNK and CH₃-oxo-NNK, respectively. Sections are as follows: (A) CH₂-oxo-NNK standard, (B) CH₃-oxo-NNK standard, and NNK-P450 2A13 incubations containing all relevant enzymes and cofactors with incubation times of (C) 1 min, (D) 5 min, (E) 10 min, and (F) 60 min. RT = retention time; MA = Mass Area.

16

Figure 3: LC-NSI-HRMS chromatograms resulting from the NNN-P450 2A6 incubations. For all sections, the top chromatogram is the accurate parent mass extracted from full scan for NNC. The middle and bottom chromatograms are two accurate product ion masses extracted from MS² fragmentation for NNC. Sections are as follows: (A) NNC standard, and NNC-P450 2A6 incubations containing all relevant enzymes and cofactors with incubation times of (B) 1 min, (C) 5 min, and (D) 10 min. RT = retention time; MA = Mass Area.

Figure 1:




Figure 2:





Figure 3:



Scheme Legends:

2


Scheme 1: (A) Established *in vivo* metabolism of NNK (**1**) and NNN (**2**) by P450 2A13- or P450 2A6-mediated oxidation, respectively. Oxidation results in unstable α -hydroxynitrosamines (**3** - **6**) which spontaneously decompose to diazohydroxides (**7** - **9**). These either hydrolyze to products excreted in the urine (**10** - **12**) or react with DNA to form adducts. (B) Proposed P450-mediated oxidation of NNK (**1**) and NNN (**2**) to nitrosamides (**13** - **15**) through retention of the α -hydroxynitrosamines **3**, **4**, and **6** within the P450 active site. If formed *in vivo*, we anticipate these species would also form adducts with DNA.

Scheme 2: Synthesis of Nitrosamides (A) Methyl Acrylate, NaCN, DMF, 40 °C, 4h; (B) NaOH, H₂O, RT, 3h; (C) EDAC, NHS, MeNH₂•HCl, DMSO, RT, 22h; (D) NaNO₂, Ac₂O:HOAc, 0 °C, 4h; (E) HS(CH₂)₃SH, BF₃•OEt₂, THF, 80 °C, 24h; (F) (i) n-BuLi, TMEDA, THF, -78 °C, 1h; (ii) **27**, THF, -78 °C to RT, 16h; (G) 25% TFA, CH₂Cl₂, RT, 3h; (H) HCO₂Me, Et₃N, MeOH, 55 °C, 4h; (I) AgNO₃, NCS, MeCN:H₂O (1:1), 0 °C, 30 min; (J) Boc₂O, Et₃N, CH₂Cl₂, RT, 30 min; (K) I₂, PPh₃, Im., CH₂Cl₂, 0 °C-RT, 22h.


Scheme 3: (A) Mechanism of hydrolysis of nitrosamides. Hydrolysis results in a carboxylic acid and an alcohol via a transient diazohydroxide that decomposes to a diazonium ion. (B) The hypothesized decomposition products of CH₂-oxo-NNK, CH₃-oxo-NNK, and NNC in assay buffer (pH = 7.4) at 37 °C.

1 Scheme 1.

A.




B.



2

Scheme 2:



Scheme 3:

