<u>LETTERS</u>

1,3-Dipolar Cycloadditions of Diazo Compounds in the Presence of Azides

Matthew R. Aronoff,^{†,§} Brian Gold,^{†,§} and Ronald T. Raines^{*,†,‡}

[†]Department of Chemistry and [‡]Department of Biochemistry, University of Wisconsin—Madison, Madison, Wisconsin 53706, United States

Supporting Information

ABSTRACT: The diazo group has untapped utility in chemical biology. The tolerance of stabilized diazo groups to cellular metabolism is comparable to that of azido groups. However, chemoselectivity has been elusive, as both groups undergo 1,3-dipolar cycloadditions with strained alkynes. Removing strain and tuning dipolarophile electronics yields diazo group selective 1,3-dipolar cycloadditions that can be performed in the presence of an azido group. For example, diazoacetamide but not its azido congener react with dehydroalanine residues, as in the natural product nisin.



Recent developments¹ in the Huisgen azide–alkyne 1,3dipolar cycloaddition² have led to an apogee of interest in the azido group. Now, azides dominate the landscape of chemoselective reactions in chemical biology.^{3,4} Similar but distinct reactivity makes the diazo group an attractive alternative, especially when multicomponent selectivity is desirable.

The diazo group has been of tremendous utility to the field of organic chemistry for well over a century.⁵ This functional group is capable of varied reactivity, including alkylation, carbene generation, nucleophilic addition, homologation, and ring expansions.^{5d} A stabilized diazo group can also serve as a reporter for chemical biology, undergoing cycloadditions with the cyclooctynes now in common use in strain-promoted azide–alkyne cycloadditions (SPAACs) (Scheme 1).⁶

With its compact size, facile preparation, and stability in an aqueous environment, a pendant diazoacetamide group is wellsuited for biological investigations.⁷ This type of diazo group is capable of faster cycloadditions with common cyclooctynes than is its azido analogue.⁸ Nevertheless, the reactivity of these two groups is redundant, and the overlap encouraged us to search for a cycloaddition that is selective for only one.

We sought a reaction that not only differentiated the reactivity of diazo and azido groups but also was compatible with physiological conditions. Known strategies exist for diazo-specific reactions in aqueous conditions, such as metal—carbenoid additions/insertions,⁹ esterifications,¹⁰ and carbonyl additions,¹¹ but each of these processes has drawbacks that prevent broad application in a biological context. Frustrated by the lack of selectivity generated with SPAAC-type reactions between azido and diazo groups, we sought a simple cycloaddition reaction that would favor the characteristics of a diazo group.

In theory, orthogonal cycloadditions could be generated through modulation of the electronics of the reacting partners. In normal-electron-demand (NED) cycloadditions, the dipole has relatively high nucleophilicity; in inverse-electron-demand (IED) cycloadditions, the dipole has relatively high electrophilicity.¹² Selectivity from electronic demand has been attained using a diversity of azides¹³ as well as nitrones.¹⁴ Here, we report a strategy to differentiate a diazo group and analogous azido group through chemoselective NED cycloadditions.

Diazo compounds undergo uncatalyzed cycloadditions with numerous dipolarophiles other than strained alkynes.^{Sb,6,8,15} The decreased HOMO–LUMO gap imparted by strain enables efficient reactions with azides. For example, electron-rich azides react more rapidly than do electron-deficient azides in a cycloaddition with DIBAC.^{13,16} DIBAC also reacts faster than do other cyclooctynes with the diazoacetamide functionality.⁶

To search for preferential reactivity of the more electron-rich diazo group over the azido group, we performed a competition experiment. One equivalent each of both a diazoacetamide (1) and its azide analogue (2) were combined with 1 equiv of DIBAC and allowed to react until the octyne was consumed completely (Scheme 1). Despite the ~5-fold faster rate of the diazo group, which was consistent with calculated values of ΔG^{\ddagger} (Figure 1), a significant percentage of the azido group reacted as well. The modest rate difference did not produce selectivity.

From this precedent, we reevaluated the need for strain to effect 1,3-dipolar cycloadditions with diazoacetamide 1. Specifically, we hypothesized that we could replace strain with electronic activation. In this modality, dipolarophiles possessing a lower energy LUMO, generated from bond polarization of adjacent electron-withdrawing groups—rather than through strain—would complement the higher nucleophilicity of diazo compounds.

We first examined ethyl propiolate \mathbf{b} and its alkene analogue, ethyl acrylate \mathbf{a} , under conditions identical to those in the competition reaction with DIBAC. In this case, selective reactivity for the diazoacetamide over the azide was observed,

Received: January 27, 2016

Scheme 1. (Left) 1,3-Dipolar Cycloaddition of a 1:1 Mixture of Diazoacetamide 1 and Azidoacetamide 2 with a Strained Dipolarophile ($R = CH_2CH_2NH_2$) Was Not Selective; (Right) 1,3-Dipolar Cycloaddition with Unstrained Dipolarophiles Was Selective for the Diazo Compound^a



^{*a*}Conditions: 1:1 CH₃CN/H₂O; \leq 72 h; ambient temperature.



Figure 1. Computational analysis of cycloadditions with *N*-methyl-2diazoacetamide and *N*-methyl-2-azidoacetamide. Optimized geometries, activation energies, distortion–interaction energies, and changes in dipolarophile charge (*italics*) were calculated at the M06-2X/6-31+G(2d,p) level of theory. Energies (kcal/mol) and NBO charges on dipolarophiles (*italics*) include solvation corrections (water) on gas-phase geometries using IEFPCM model (radii = UFF).

even in the presence of a 5-fold excess of dipolarophile (Scheme 1). These experimental results are consistent with a computational analysis, which suggests that the cycloaddition of the diazo group proceeds by a transition state (TS) of lower ΔG^{\ddagger} (Figure 1). Although the dipoles of both 1 and 2 favor NED cycloadditions, the greater nucleophilicity of the diazo group of 1 leads to chemoselectivity. Cycloaddition of the diazo group appears to be less synchronous, as its TS exhibits greater transfer of charge from the dipole to the dipolarophile as well as a longer terminal N···C bond distance (Figure 1). Notably, the relative reactivity of the diazo group is amplified in aqueous conditions (Figure S3),^{6,8} which favor the more polar TS of the diazo group.¹⁷

Ethyl acrylate (a) reacted more quickly than did ethyl propiolate (b) with diazoacetamide 1 ($k = 1.6 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$ and $k = 5.8 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$, respectively). This result was surprising, as previous reports had suggested similar reaction rates for alkenes and alkynes, despite different FMO energies.¹⁸ The difference that we observed is likely due to a smaller distortion energy (ΔE_{dist}) for dipolarophile **a**, along with a slightly more favorable interaction energy (ΔE_{int}) (Figure 1).¹⁹ Although alkynes are easier to bend than alkenes, an alkyne must be bent to a larger degree to reach a TS geometry.²⁰ The difference in dipolarophile distortion energy of 1.7 kcal/mol is similar to the value of 2.1 kcal/mol reported previously for the cycloadditions of diazomethane with ethylene and acetylene calculated at the B3LYP/6-31G(d) level of theory.¹⁸ These results are consistent with the smaller HOMO-LUMO gap of alkenes relative to alkynes.

In light of the faster rates observed for the alkene,²⁰ we focused our attention on ethyl acrylate and related alkenyl compounds.²¹ Upon examining the reaction kinetics of a carbonyl series of electron-deficient alkenes, we verified that the rate of the reaction was dependent upon the electron-withdrawing capabilities of the substituent. Cycloadditions with a variety of dipolarophiles (**a**–**f**, see Supporting Information) produced rate constants comparable to those of early SPAACs (that is, $k \approx 10^{-2} \text{ M}^{-1} \text{ s}^{-1}$).^{4b,22}

We next sought to challenge the selectivity in a simple context. To do so, we synthesized a compact molecule containing both a diazo group and an azido group. We reacted diazo azide 3 with an excess of ethyl acrylate **a**. The results were gratifying—the diazo group reacted completely to form the pyrazoline product 3**a**, while the azido group remained intact and free for subsequent functionalization or derivation (Scheme 2).

Scheme 2. Competition for a Dipolarophile by a Diazo Group and Azido Group in the Same $Molecule^{a}$



"The reaction was selective for the diazo group. Conditions: CH $_3$ CN/ H $_2$ O 1:1; 24 h.

Lastly, we considered applications of this chemoselective process within the arena of chemical biology. Acrylate esters are too reactive for most biological applications.⁶ The tempered reactivity of acrylamides, however, enables their metabolic incorporation into biomolecules.²³ For example, the 2-acetamidoacrylamide group in dehydroalanine (Dha) is the product of a post-translational elimination of H₂O from serine or H₂S from cysteine.²⁴ The alkenyl group of dehydroalanine is in conjugation with both an amide carbonyl group and an amide nitrogen. We hypothesized that this functionality might be a good substrate for reaction with a diazo group.

We first tested the reactivity of diazoacetamide **1** with an *O*acyl derivative of Dha, methyl 2-acetamidoacrylate. The reaction rate exceeded that of previously tested alkenes by an order of magnitude, proceeding with $k = 6.7 \times 10^{-2} \text{ M}^{-1} \text{ s}^{-1}$. This enhanced rate was recapitulated in computational investigations, as calculated free energies of activation with a 2-acetamido-*N*-methylacrylamide model system were similar to those predicted for *N*-alkylacrylamides (Figure 2 and Table 1S).



Figure 2. Computational analysis of the cycloaddition of 2-acetamido-*N*-methylacrylamide with *N*-methyl-2-diazoacetamide. Optimized geometries and free energies were calculated at the M06-2X/6-31+G(2d,p) level of theory. Energies include solvation corrections (water) on gas-phase geometries using IEFPCM model (radii = UFF).

The Curtin–Hammett principle predicts a ΔG^{\ddagger} of 29.1 kcal/ mol for the lower energy C-5 conformation, but enforcement of the C-7 conformation would provide a lower overall ΔG^{\ddagger} of 27.4 kcal/mol.

Inspired by the reactivity of diazoacetamide 1 with 2acetamido-*N*-methylacrylamide, we investigated its reactivity with a natural product. Dehydroalanine is found in lantibiotics, including the peptide antibiotic nisin.²⁵ Nisin is well suited to test the biological relevance of the cycloaddition with the diazo group, as it contains two Dha residues and one dehydrodrobutyrine (Dhb) residue (which is derived from threonine) along with myriad common peptidic functional groups (Figure S5, Supporting Information).

We began our work with nisin by assessing the ability of diazoacetamide 1 to label a purified sample of the natural product. After 16 h, most of the peptide had experienced a single cycloaddition, and some had reacted twice (Figure 3). Notably, the full mass of the diazo compound had been added to the peptide, ruling out the possibility of reaction by esterification, as is possible with diazo groups.¹⁰

Finally, we tested the utility of this cycloaddition for naturalproduct discovery. To do so, we reacted a crude mixture of denatured protein solids containing 2.5% w/w nisin with diazoacetamide—biotin conjugate. We observed a change in mass expected for the labeling of nisin by cycloaddition (Figure S33, Supporting Information). No reaction was observed with the analogous azidoacetamide—biotin conjugate.

In conclusion, we report on a simple means to differentiate the reactivity of an azido group and a diazo group. Whereas strain is necessary to obtain useful cycloaddition rates with azides, electronic activation of alkenes allows for rapid reactivity with diazoacetamides. This insight enables cycloadditions to dehydroalanine, such as that in nisin. To our knowledge, this example is the first of a cycloaddition to a natural amino acid residue. The chemoselectivity of the diazo–Dha cycloaddition could aid in applications of known natural products and the discovery of new ones.



Figure 3. MALDI–TOF spectra of the products of the cycloaddition of nisin with N-benzyl-2-diazoacetamide (1) and N-benzyl-2-azidoacetamide (2). Conditions: 10 mM sodium phosphate buffer (pH 7.3), 16 h, ambient temperature.

Organic Letters

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.or-glett.6b00278.

Synthetic methods and analytical data along with additional computational data, Cartesian coordinates, energies, and imaginary frequencies for TSs (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: rtraines@wisc.edu.

Author Contributions

[§]M.R.A. and B.G. contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by Grant No. R01 GM044783 (NIH) and made use of the National Magnetic Resonance Facility at Madison, which is supported by Grant No. P41 GM103399 (NIH). The purchase of the Bruker microflex LRF in 2015 was funded by the Bender gift to the Department of Chemistry. Computational resources were supported in part by Grant No. CHE-0840494 (NSF). We are grateful to K. A. Andersen and C. L. Jenkins (University of Wisconsin—Madison) for contributive discussions.

REFERENCES

(1) (a) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596–2599. (b) Tornøe, C. W.; Christensen, C.; Meldal, M. J. Org. Chem. **2002**, *67*, 3057–3064.

(2) For key historical references, see: (a) Dimroth, O. Liebigs Ann.
1909, 364, 183–226. (b) Huisgen, R. Proc. Chem. Soc. 1961, 357–396.
(c) Wittig, G.; Krebs, A. Chem. Ber. 1961, 94, 3260–3275.
(d) Huisgen, R. Angew. Chem., Int. Ed. Engl. 1963, 2, 565–598.
(e) Huisgen, R.; Szeimies, G.; Möbius, L. Chem. Ber. 1967, 100, 2494–2507.

(3) For examples, see: (a) Agard, N. J.; Prescher, J. A.; Bertozzi, C. R. I. Am. Chem. Soc. 2004, 126, 15046-15047. (b) Baskin, J. M.; Prescher, J. A.; Laughlin, S. T.; Agard, N. J.; Chang, P. V.; Miller, I. A.; Lo, A.; Codelli, J. A.; Bertozzi, C. R. Proc. Natl. Acad. Sci. U. S. A. 2007, 104, 16793-16797. (c) Ning, X.; Guo, J.; Wolfert, M. A.; Boons, G.-J. Angew. Chem., Int. Ed. 2008, 47, 2253-2255. (d) Poloukhtine, A. A.; Mbua, N. E.; Wolfert, M. A.; Boons, G.-J.; Popik, V. V. J. Am. Chem. Soc. 2009, 131, 15769-15776. (e) Debets, M. F.; van Berkel, S. S.; Schoffelen, S.; Rutjes, F. P. J. T.; van Hest, J. C. M.; van Delft, F. L. Chem. Commun. 2010, 46, 97-99. (f) Jewett, J. C.; Sletten, E. M.; Bertozzi, C. R. J. Am. Chem. Soc. 2010, 132, 3688-3690. (g) Dommerholt, J.; Schmidt, S.; Temming, R.; Hendriks, L. J. A.; Rutjes, F. P. J. T.; van Hest, J. C. M.; Lefeber, D. J.; Friedl, P.; van Delft, F. L. Angew. Chem., Int. Ed. 2010, 49, 9422-9425. (h) Varga, B. R.; Kállay, M.; Hegyi, K.; Béni, S.; Kele, P. Chem. - Eur. J. 2012, 18, 822-828. (i) Gold, B.; Dudley, G. B.; Alabugin, I. V. J. Am. Chem. Soc. 2013, 135, 1558-1569. (j) Ni, R.; Mitsuda, N.; Kashiwagi, T.; Igawa, K.; Tomooka, K. Angew. Chem., Int. Ed. 2015, 54, 1190-1194.

(4) For recent reviews, see: (a) Sletten, E. M.; Bertozzi, C. R. Acc. Chem. Res. 2011, 44, 666–676. (b) Debets, M. F.; van Berkel, S. S.; Dommerholt, J.; Dirks, T.; Rutjes, F. P. J. T. Acc. Chem. Res. 2011, 44, 805–815. (c) Schilling, C. I.; Jung, N.; Biskup, M.; Schepers, U.; Bräse, S. Chem. Soc. Rev. 2011, 40, 4840–4871. (d) Patterson, D. M.; Nazarova, L. A.; Prescher, J. A. ACS Chem. Biol. 2014, 9, 592–605.

(5) For relevant background, see: (a) Pechmann, H. v. Ber. Dtsch. Chem. Ges. 1894, 27, 1888–1891. (b) Cowell, G. W.; Ledwith, A. Q. Rev., Chem. Soc. 1970, 24, 119–167. (c) Maas, G. In Organic Synthesis, Reactions and Mechanisms; Springer: Berlin, 1987; pp 75–253. (d) Regitz, M. Diazo Compounds: Properties and Synthesis; Elsevier: Philadelphia, 2012.

(6) Andersen, K. A.; Aronoff, M. R.; McGrath, N. A.; Raines, R. T. J. Am. Chem. Soc. 2015, 137, 2412–2415.

(7) Caution! Unstabilized diazo compounds are dangerous, and their use should never be attempted in the context of chemical biology. See: (a) Lewinn, E. B. *Am. J. Med. Sci.* **1949**, *218*, 556–562. (b) Schoental, R. *Nature* **1960**, *188*, 420–421. (c) Lewis, C. E. *J. Occup. Environ. Med.* **1964**, *6*, 91–92.

(8) McGrath, N. A.; Raines, R. T. Chem. Sci. 2012, 3, 3237–3240. (9) (a) Antos, J. M.; Francis, M. B. J. Am. Chem. Soc. 2004, 126, 10256–10257. (b) Antos, J. M.; McFarland, J. M.; Iavarone, A. T.; Francis, M. B. J. Am. Chem. Soc. 2009, 131, 6301–6308. (c) Tishinov, K.; Schmidt, K.; Häussinger, D.; Gillingham, D. G. Angew. Chem., Int. Ed. 2012, 51, 12000–12004. (d) Ball, Z. T. Acc. Chem. Res. 2013, 46, 560–570. (e) Fei, N.; Haussinger, D.; Blumli, S.; Laventie, B.-J.; Bizzini, L. D.; Zimmermann, K.; Jenal, U.; Gillingham, D. Chem. Commun. 2014, 50, 8499–8502. (f) Ball, Z. T. Curr. Opin. Chem. Biol. 2015, 25, 98–102.

(10) (a) McGrath, N. A.; Andersen, K. A.; Davis, A. K. F.; Lomax, J. E.; Raines, R. T. *Chem. Sci.* **2015**, *6*, 752–755. (b) Mix, K. A.; Raines, R. T. *Org. Lett.* **2015**, *17*, 2358–2361.

- (11) Zhang, Y.; Wang, J. Chem. Commun. 2009, 5350-5361.
- (12) (a) Houk, K. N.; Sims, J.; Duke, R. E.; Strozier, R. W.; George, J. K. J. Am. Chem. Soc. 1973, 95, 7287–7301. (b) Houk, K. N.; Sims, J.; Watts, C. R.; Luskus, L. J. J. Am. Chem. Soc. 1973, 95, 7301–7315.
- (13) Dommerholt, J.; van Rooijen, O.; Borrmann, A.; Guerra, C. F.; Bickelhaupt, F. M.; van Delft, F. L. *Nat. Commun.* **2014**, *5*, 5378.
- (14) MacKenzie, D. A.; Pezacki, J. P. Can. J. Chem. 2014, 92, 337-340.

(15) Jung, M. E.; Min, S.-J.; Houk, K. N.; Ess, D. J. Org. Chem. 2004, 69, 9085–9089.

(16) Xie, S.; Lopez, S. A.; Ramström, O.; Yan, M.; Houk, K. N. J. Am. Chem. Soc. 2015, 137, 2958–2966.

(17) Ess, D. H.; Jones, G. O.; Houk, K. N. Adv. Synth. Catal. 2006, 348, 2337–2361.

(18) Ess, D. H.; Houk, K. N. J. Am. Chem. Soc. 2008, 130, 10187–10198.

(19) For a tutorial review of the distortion/interaction model (which is also known as the activation strain model), see: Fernández, I.; Bickelhaupt, F. M. *Chem. Soc. Rev.* **2014**, *43*, 4953–4967.

(20) Liu, F.; Liang, Y.; Houk, K. N. J. Am. Chem. Soc. 2014, 136, 11483-11493.

(21) This decision echoes back to the 19th century origin of cycloaddition chemistry, when Buchner discovered the cycloaddition of diazo ethylacetate and an α,β -unsaturated carboxylic ester: Buchner, E. Ber. Dtsch. Chem. Ges. **1888**, 21, 2637–2647.

(22) Lang, K.; Chin, J. W. ACS Chem. Biol. 2014, 9, 16-20.

(23) Lee, Y.-J.; Wu, B.; Raymond, J. E.; Zeng, Y.; Fang, X.; Wooley, K. L.; Liu, W. R. ACS Chem. Biol. 2013, 8, 1664–1670.

(24) For a recent review of dehydroalanine, see: Siodłak, D. Amino Acids 2015, 47, 1–17.

(25) (a) Gross, E.; Morell, J. L. J. Am. Chem. Soc. 1971, 93, 4634– 4635. (b) Wiedemann, I.; Breukink, E.; van Kraaij, C.; Kuipers, O. P.; Bierbaum, G.; de Kruijff, B.; Sahl, H.-G. J. Biol. Chem. 2001, 276, 1772–1779. (c) Lubelski, J.; Rink, R.; Khusainov, R.; Moll, G. N.; Kuipers, O. P. Cell. Mol. Life Sci. 2008, 65, 455–476. (d) Ortega, M. A.; Hao, Y.; Zhang, Q.; Walker, M. C.; van der Donk, W. A.; Nair, S. K. Nature 2014, 517, 509–512.