

149. Carbenoid Reactions of Dimethyl Diazomalonate with Aromatic Thioketones and 1,3-Thiazole-5(4H)-thiones

by Grzegorz Mlostón¹⁾*

Department of Organic & Applied Chemistry of the University of Łódź, Narutowicza 68, PL-90-136 Łódź

and Heinz Heimgartner*

Organisch-chemisches Institut der Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich

(14. VII. 96)

Dimethyl diazomalonate (**4**) and thiobenzophenone (**2a**) do not react in toluene even after warming to 50°. After addition of catalytic amounts of Rh₂(OAc)₄, a smooth reaction under N₂ evolution afforded a mixture of thiiranedicarboxylate **5** and (diphenylmethylidene)malonate **6** (Scheme 2). A reaction mechanism via an intermediate 'thiocarbonyl ylide' **7**, formed by the addition of the carbenoid species **8** to the S-atom of **2a**, is plausible. Similar reactions were carried out with 9H-xanthene-9-thione (**2b**), 9H-thioxanthene-9-thione (**2c**, Scheme 4), and 1,3-thiazole-5(4H)-thione **18** (Scheme 6). In the cases of **2b** and **2c**, spirocyclic 1,3-dithiolanetetracarboxylates **14a** and **14b**, respectively, were obtained as the third product. Reaction mechanisms for their formation are proposed in Scheme 5: S-transfer from intermediate thiirane **12** to the carbenoid species yielded thioxomalonate **15** which underwent a 1,3-dipolar cycloaddition with 'thiocarbonyl ylide' **16**. An alternative is the formation of 'thiocarbonyl ylide' **17** via carbene addition to **15**, followed by 1,3-dipolar cycloaddition with **2b** and **2c**, respectively.

1. Introduction. – Diazoalkanes and their aryl-substituted derivatives react efficiently with thiocarbonyl compounds to give 2,5-dihydro-1,3,4-thiadiazoles as primary products (cf. [1–3]). Most of these adducts are unstable at ambient temperature and eliminate N₂ spontaneously or after slight warming to give reactive 'thiocarbonyl ylides' which undergo various reactions such as 1,3-dipolar cycloadditions [4–6], ring closure to thiiranes [7] [8], dimerization to 1,4-dithianes [9–11], and 1,4-H shifts [12]. Recently, we described that α -diazo ketones, -amides, and -esters are less reactive in 1,3-dipolar cycloadditions with thioketones and, therefore, these reactions required higher temperature [13] [14]. Generally, in reactions with carbonyl-substituted diazo compounds, the primary adducts have not been isolated until now, and elimination of N₂ occurred already under the reaction conditions²⁾.


Recently, we described results of reactions of methyl 2-diazo-3,3,3-trifluoropropanoate (**1**) with thiocarbonyl compounds, and we stated that only the most reactive thiobenzophenone (**2a**) reacted under standard conditions with evolution of N₂ to yield thiirane **3a** [16] (Scheme 1). Under similar conditions, the less reactive 9H-xanthene-9-thione (**2b**) and 9H-thioxanthene-9-thione (**2c**) reacted with **1** only the presence of catalytic amounts of rhodium(II) acetate, and the only products detected in the reaction

¹⁾ Stay at the Institute of Organic Chemistry of the University of Zürich (March–June 1995) as a Swiss Federal Scholar (Bundesstipendiat).

²⁾ Recently, the primary, stable cycloadduct of a sterically crowded diazo amide and 2,2,4,4-tetramethyl-3-thioxocyclobutanone could be isolated [15].

mixtures were thiiranes **3b** and **3c**, respectively. In the case of 4,4-dimethyl-2-phenyl-1,3-thiazole-5(4*H*)-thione, the corresponding thiirane was isolated as a minor product, and olefinic compounds, formed by its desulfurization, were the major components of the reaction mixture. It is likely that the formation of thiiranes **3** proceeds *via* ring closure of intermediate ‘thiocarbonyl ylides’. In the noncatalyzed reaction, the precursor of this species is the [2+3] cycloadduct of **1** and **2a**. However, in the catalyzed decomposition of **1**, the formation of a carbenoid must be the first step of the conversion³⁾. The electrophilic carbenoid then adds to the S-atom of the thiocarbonyl group to generate the ‘thiocarbonyl ylide’.

Scheme 1




Diazomalonates are known as very useful reagents in organic syntheses; especially important are their applications as precursors of carbenes and carbenoids [18–20]. On the other hand, it is well known that diazomalonates are practically unreactive as 1,3-dipoles. Some years ago, *Huisgen* and coworkers found that thioketones are very reactive dipolarophiles (‘superdipolarophiles’ [3] [21]), and for this reason, we decided to test them as probably the best dipolarophiles in reactions with dimethyl diazomalonate (**4**).

2. Results and Discussion. – According to *Huisgen*, thiobenzophenone (**2a**) is one of the most reactive thiocarbonyl compounds in 1,3-dipolar cycloadditions. In our hands, heating of a solution of equimolar amounts of **2a** and **4** in toluene to 50° did not lead to any change of the mixture. However, addition of a catalytic amount of $\text{Rh}_2(\text{OAc})_4$ resulted in a slow evolution of N_2 . After 6 h, the reaction was terminated; the initially blue color of the mixture disappeared, and an almost equimolar amount of N_2 was set free. The ^1H -NMR spectrum of the mixture showed the presence of two new products in a ratio of *ca.* 7:3 (2*s* at 3.48 and 3.57 ppm, resp. (MeO)). After chromatographic workup, the two products were isolated and identified as thiirane **5** (MeO at 3.50 ppm) and the known alkylidenemalonate **6** [22] (MeO at 3.60 ppm) (Scheme 2).

³⁾ Unlike reactions of carbenes and carbenoids with other $\text{C}=\text{X}$ bonds ($\text{X} = \text{C}, \text{N}, \text{O}$), reactions with thiocarbonyl groups are very scarcely described (*cf.* [17]).


Scheme 2



The formation of **5** results from the cyclization of an intermediate ‘thiocarbonyl ylide’ **7**, formed by the addition of carbenoid **8** to the S-atom of **2a**. Desulfurization of **5** then yields **6**. The chemistry of ‘thiocarbonyl ylides’ of type **7** with two electron-withdrawing substituents is almost unknown⁴⁾. In contrast to similar intermediates with alkyl and/or aryl groups, **7** underwent no cycloaddition with the ‘superdipolarophile’ **2a**. A similar behavior was already found for ‘thiocarbonyl ylides’ with an ester and a CF₃ group [16]. *Kaufman* and *Weininger* described a photolytical reaction of **2a** with diazomalonate and stated that the major product was diethyl (diphenylmethylidene)malonate [23]. According to the same authors, refluxing of **2a** and diethyl diazomalonate in diglyme gave the same product in good yield. It is likely that, under photolytical conditions and thermally at 162°, elimination of the S-atom from primarily formed thiirane occurs rapidly. The advantage of our Rh-catalyzed reaction is that under relatively mild conditions, thiirane **5** can be prepared in fairly good yield.

A similar generation of the ‘thiocarbonyl ylide’ intermediate **9** has been used by *Tokitoh et al.* [24]: the reaction of di(*tert*-butyl)thioketene and **4** in the presence of Rh₂(OAc)₄ at 50° yielded the 2-alkylidene-1,3-oxathiole **10**, which has been shown to be in an equilibrium with **9** (Scheme 3). At 70° in CCl₄, **10** rearranged to give allene episulfide **11**.


Scheme 3



⁴⁾ Recent experiments concerning the generation of ‘thiocarbonyl-S-dichloromethanides’ will be published elsewhere.

In reactions with diazo compounds, 9*H*-xanthene-9-thione (**2b**) and 9*H*-thioxanthene-9-thione (**2c**) are less reactive dipolarophiles than **2a** [21]. Therefore, reactions of both thioketones with **4** were performed at 60° in the presence of Rh₂(OAc)₄ catalyst. After 2–3 days, the equimolar amount of N₂ was produced, and, after chromatographic workup, the products **12–14** were isolated (*Scheme 4*). Similar to the reactions with **2a**, the less polar fractions were identified as thiirane **12** and malonate **13**. The elucidation of the structure of the most polar **14**, a new type of product, based on mass and ¹³C-NMR spectroscopy.


Scheme 4



In the ¹H-NMR spectra, **12** and **13** showed only one and **14** two MeO signals. The CI-MS of **14a** and **14b** indicated that both products contain two malonate moieties and two S-atoms in addition to the xanthene and thioxanthene fragment, respectively. The ¹³C-NMR spectrum of **14a** revealed 3 signals for quaternary C-atoms at 96.1, 69.5, and 67.8 ppm. A similar set of signals was found for **14b**. All these data are in good agreement with the structure of 1,3-dithiolane derivatives.


The formation of the unexpected products **14a,b** may be formulated as shown in *Scheme 5*. The crucial step is the formation of the new thiocarbonyl compounds **15**. We propose that **15** is formed by S-transfer from thiirane **12** to the carbene species. Subsequent interception of 'thiocarbonyl ylide' **16** by **15** results in the formation of **14**. However, a second route to **14** should also be taken into account: reaction of the carbene species **8** with *in situ* generated **15** could give 'thiocarbonyl ylide' **17** which, *via* cycloaddition with **2**, can also lead to **14**. The trapping of 'thiocarbonyl ylides' by thiones is a characteristic reaction of these S-centered 1,3-dipoles and known as *Schönberg* reaction [25]. A desulfurization of a thiirane by carbenoid **8**, similar to **12** → **13**, has been postulated by *Takano et al.*, but in their experiments, all attempts to isolate or to trap the resulting thiocarbonyl compound were unsuccessful [26].

Scheme 5



From our previous studies it is known that 1,3-thiazole-5(4*H*)-thiones of type **18** are much less reactive as dipolarophiles compared with aromatic thioketones [16]. The $\text{Rh}_2(\text{OAc})_4$ -catalyzed reaction of **4** and **18** at 60° yielded (1,3-thiazol-5-ylidene)malonate **19** in almost quantitative yield as the sole product (Scheme 6).

Scheme 6



It is worth noting that in the IR spectrum (KBr) of **19** two ester-CO absorptions at 1725 and 1705 cm^{-1} were observed. The nonequivalent ester groups also show remarkable differences in the ^{13}C -NMR spectrum: the low-field signal appears at 172.3 and the high-field one at 166.4 ppm. These differences can be explained by the assumption that the ester group *cis* to the S-atom is coplanar with the C=C bond (i.e., an α,β -unsaturated ester) whereas the *trans*-oriented ester group, because of steric hindrance with the geminal dimethyl groups, is twisted out of the C=C plane (cf. [16]).

The formation of **19** corresponds very well with results obtained from reactions of **4** and analogous diazo compounds with tetrahydrofuran-2-thiones in refluxing benzene

[26] [27]. In this case, despite of the lower temperature in our experiments, the desulfurization of primarily formed thiirane was complete. This simple methodology for the formation of methylenide derivatives of heterocyclic thiones, which is closely related to *Eschenmoser's* 'sulfide contraction via alkylative coupling' [28], has already been used in natural-product syntheses [27]. Our present studies show that this rather general strategy can be performed as a 'one-pot reaction' under relatively mild conditions.

In contrast to [24], using the 'carbenoid procedure', we could not detect products of 1,5-electrocyclization of ester-substituted 'thiocarbonyl-ylide' intermediates, neither in reactions with aromatic thioketones nor with 1,3-thiazole-5(4*H*)-thione **18**.

We thank the analytical sections of our institutes for spectra and analyses, the *Polish State Committee for Scientific Research* (grant No. 3 TO9A 257 10), the *Swiss National Science Foundation*, and *F. Hoffmann-La Roche AG*, Basel, for financial support. *G. M.* thanks the *Swiss Federal Commission for Foreign Students* for a scholarship.

Experimental Part

1. *General*. See [5]. Thioketones **2a–c** were prepared from the corresponding ketones by treatment with *Lawesson reagent* [29]; 4,4-dimethyl-2-phenyl-1,3-thiazole-5(4*H*)-thione (**18**) was synthesized according to [30]. Dimethyl diazomalonate (**4**) was prepared from dimethyl malonate by treatment with tosyl azide in benzene as described in [31].

2. *Reactions of Thiocarbonyl Compounds with Dimethyl Diazomalonate (4)*. 2.1. *General Procedure*. At r.t., **4** (177.0 mg, 1.1 mmol) was added to a soln. of the thiocarbonyl compound (1.0 mmol) in toluene (0.5 ml). No N₂ evolution was observed, neither at r.t. nor at 50°. After addition of a catal. amount (*ca.* 0.1 mmol) of Rh₂(OAc)₄·H₂O at 50°, N₂ evolution started and was followed volumetrically. After 6–42 h, the evolution of N₂ ceased. The mixtures were filtered through a short SiO₂ column (*ca.* 3 cm) to remove the catalyst and dark decomposition products. After evaporation, the residues were analyzed by ¹H-NMR (1,1,2,2-tetrachloroethane as a standard) to establish yields of the products. Prep. TLC (SiO₂) and crystallization yielded pure products.

2.2. *With Thiobenzophenone (2a)*. For 6 h; prep. TLC with pentane/CH₂Cl₂ 6:4: two fractions.

Dimethyl 3,3-Diphenylthiirane-2,2-dicarboxylate (5): 46% (NMR); isolated 130 mg (40%). Colorless crystals. M.p. 92–93° (pentane/CH₂Cl₂). IR (KBr): 1745vs (C=O), 1490m, 1447m, 1432s, 1275m (br.), 1250–1200s (br.), 1125vs, 1052vs, 975m, 952m, 765s (br.), 732m, 708s, 695s. ¹H-NMR: 7.45–7.4 (m, 4 arom. H); 7.3–7.25 (m, 6 arom. H); 3.50 (s, 2 MeO). ¹³C-NMR: 166.0 (s, 2 C=O); 138.9 (s, 2 arom. C); 129.1, 128.0, 127.9 (3d, 10 arom. CH); 62.7 (s, C(2)); 57.3 (s, C(3)); 53.2 (q, 2 MeO). CI-MS: 329 (25, [M + I]⁺), 297 (100, [M – S]⁺), 282 (95, [M – MeS]⁺), 265 (31). Anal. calc. for C₁₈H₁₆O₄S (328.37): C 65.84, H 4.91, S 9.76; found: C 65.55, H 5.35, S 10.05.

Dimethyl (Diphenylmethylenide)malonate (6): 18% (NMR); isolated 40 mg (13%). Colorless crystals. M.p. 119–121° (EtOH at –10°; [21]: 122–123°). IR (KBr): 1730vs (C=O), 1700vs (C=O), 1600s (C=C), 1440s, 1425s, 1330vs, 1300s, 1245vs (br.), 1165s, 1080vs, 770s, 695s. ¹H-NMR: 7.35–7.3 (m, 6 arom. H); 7.25–7.15 (m, 4 arom. H); 3.60 (s, 2 MeO). ¹³C-NMR: 166.3 (s, 2 C=O); 156.4 (s, Ph₂C); 139.8 (s, 2 arom. C); 129.2, 129.0, 128.1 (3d, 10 arom. CH); 125.4 (s, C(CO₂Me)₂); 52.1 (q, 2 MeO). EI-MS: 296 (92, M⁺), 265 (73, [M – MeO]⁺), 236 (100, [M – CO₂Me]⁺), 205 (18), 197 (30), 178 (42), 165 (19), 105 (31).

2.3. *With 9H-Xanthene-9-thione (2b)*. For 40 h; prep. TLC with hexane/CH₂Cl₂ 1:1: three fractions.

Dimethyl Spiro[thiirane-2,9'-[9H]xanthene]-3,3-dicarboxylate (12a): 31% (NMR); isolated 80 mg (22%). Colorless crystals. M.p. 150–152° (hexane/CH₂Cl₂). IR (KBr): 1740vs (br., C=O), 1725vs (br., C=O), 1475s, 1450vs, 1435s, 1312s, 1265vs (br.), 1235vs, 1055m, 767s, 760s. ¹H-NMR: 7.5–7.05 (m, 8 arom. H); 3.39 (s, 2 MeO). ¹³C-NMR: 164.6 (s, 2 C=O); 154.7, 119.3 (2s, 4 arom. C); 129.5, 127.3, 123.0, 116.4 (4d, 8 arom. CH); 59.9 (s, C(3)); 53.2 (q, 2 MeO); 51.1 (s, C(2)). EI-MS: 342 (5, M⁺), 310 (69, ([M – S]⁺), 283 (35, [M – CO₂Me]⁺), 282 (100), 279 (44), 268 (15), 251 (32), 224 (29), 211 (19), 194 (13), 180 (17), 163 (13). Anal. calc. for C₁₈H₁₄O₅S (342.35): C 63.15, H 4.12, S 9.36; found: C 62.95, H 3.98, S 9.11.

Dimethyl (9H-Xanthene-9-ylidene)malonate (13a): 32% (NMR); isolated 80 mg (26%). Pale yellow crystals. M.p. 95–97° (EtOH). IR (KBr): 1750s (C=O), 1710vs (br., C=O), 1610m (br., C=C), 1450s, 1340m, 1250vs (br.), 1210m (br.), 1173m, 1155m, 1075s, 775m. ¹H-NMR: 7.7–7.1 (m, 8 arom. H); 3.75 (s, 2 MeO). ¹³C-NMR: 166.2 (s, 2 C=O); 152.6, 120.9 (2s, 4 arom. C); 137.8 (s, Ar₂C); 130.9, 127.0, 123.2, 116.9 (4d, 8 arom. CH); 120.0 (s,

C(CO₂Me)₂; 52.6 (*q*, 2 MeO). EI-MS: 310 (100, *M*⁺), 279 (72, [*M* – MeO]⁺), 251 (25), 221 (23), 220 (22), 211 (28), 194 (22), 180 (16), 163 (20). Anal. calc. for C₁₈H₁₄O₅ (310.31): C 69.67, H 4.54; found: C 69.56, H 4.21.

Tetramethyl Spiro[1,3-dithiolane-4,9'-[9H]xanthene]-2,2,5,5-tetracarboxylate (14a): 17% (NMR); isolated 70 mg (14%, 25% rel. to **4**). Colorless viscous oil. IR (CHCl₃): 1745vs (br., C=O), 1475m, 1440m, 1320m, 1285m, 1240s (br.). ¹H-NMR: 8.32 (*d*, *J* = 1.3, 2 arom. H); 7.35–7.3 (*m*, 2 arom. H); 7.15–7.05 (*m*, 4 arom. H); 3.95 (*s*, 2 MeO); 3.44 (*s*, 2 MeO). ¹³C-NMR: 167.3, 165.9 (2s, 4 C=O); 151.6, 119.7 (2s, 4 arom. C); 131.8, 130.2, 122.6, 116.2 (4d, 8 arom. CH); 96.1, 69.5, 67.8 (3s, C(2), C(4), C(5)); 54.6, 53.4 (2*q*, 4 MeO). CI-MS: 522 (100, [*M* + NH₄]⁺), 505 (18, [*M* + 1]⁺), 473 (45, [*M* – MeO]⁺), 343 (90), 311 (19), 310 (34), 282 (14), 213 (44). Anal. calc. for C₂₃H₂₀O₈S₂ (504.51): C 54.75, H 3.99, S 12.71; found: C 54.29, H 3.53, S 12.51.

2.4. *With 9H-Thioxanthene-9-thione (2c)*. For 72 h (from time to time, new portions of catalyst were added); prep. TLC with hexane/CH₂Cl₂ 1:1: two fractions.

Dimethyl Spiro[thiirane-2,9'-[9H]thioxanthene]-3,3-dicarboxylate (12b): 42% (NMR); isolated 110 mg (31%). Colorless crystals. M.p. 124–126° (EtOH/CH₂Cl₂). IR (KBr): 1740vs (C=O), 1435s, 1270vs (br.), 1220vs (br.), 1115 *m*, 1050s, 760s, 740s. ¹H-NMR: 7.6–7.55 (*m*, 2 arom. H); 7.5–7.45 (*m*, 2 arom. H); 7.3–7.2 (*m*, 4 arom. H); 3.38 (*s*, 2 MeO). ¹³C-NMR: 165.0 (*s*, 2 C=O); 136.8, 132.8 (2s, 4 arom. C); 128.9, 127.8, 126.5, 126.2 (4d, 8 arom. CH); 59.2, 56.3 (2s, C(2), C(3)); 53.1 (*q*, 2 MeO). EI-MS: 358 (5, *M*⁺), 326 (100, [*M* – S]⁺), 298 (51), 295 (52), 284 (14), 267 (41), 240 (18), 236 (20), 227 (16), 208 (29), 196 (20), 163 (16), 152 (21). Anal. calc. for C₁₈H₁₄O₄S₂ (358.41): C 60.32, H 3.94, S 17.89; found: C 60.08, H 3.66, S 17.98.

Tetramethyl Spiro[1,3-dithiolane-4,9'-[9H]thioxanthene]-2,2,5,5-tetracarboxylate (14b): 24% (NMR); isolated 105 mg (20%, 36% rel. to **4**). Colorless viscous oil⁵. IR (CHCl₃): 1745vs (br., C=O), 1460 *m*, 1435s, 1230vs (br.), 1180m. ¹H-NMR: 8.47 (*d*, *J* = 1.3, 2 arom. H); 7.3–7.15 (*m*, 6 arom. H); 3.89 (*s*, 2 MeO); 3.50 (*s*, 2 MeO). ¹³C-NMR: 167.0, 165.9 (2s, 4 C=O); 133.5, 130.7 (2s, 4 arom. C); 133.0, 128.0, 125.7, 125.3 (4d, 8 arom. CH); 96.0, 76.2, 67.0 (3s, C(2), C(4), C(5)); 54.4, 53.3 (2*q*, 4 MeO). CI-MS: 538 (20, [*M* + NH₄]⁺), 537 (37, [*M* + NH₃]⁺), 521 (44, [*M* + 1]⁺), 507 (100), 489 (13), 358 (34), 298 (22).

2.5. *With 4,4-Dimethyl-2-phenyl-1,3-thiazole-5(4H)-thione (18)*. For 9 h; prep. TLC with hexane/CH₂Cl₂ 1:1: *Dimethyl (4,5-Dihydro-4,4-dimethyl-2-phenyl-1,3-thiazol-5-ylidene)malonate (19)* in 87% (NMR) yield; isolated 250 mg (78%). Colorless plates. M.p. 137–138° (EtOH). IR (KBr): 1725vs (C=O), 1705vs (C=O), 1620m (C=C), 1550s, 1450s, 1440s, 1280s, 1235vs (br.), 1190vs, 1010s, 990m, 955s, 940m, 770s, 750m, 690s. ¹H-NMR: 7.9–7.85 (*m*, 2 arom. H); 7.5–7.4 (*m*, 3 arom. H); 3.88, 3.82 (2s, 2 MeO); 1.62 (*s*, 2 Me). ¹³C-NMR: 172.3, 166.4 (2s, 2 C=O); 164.8 (*s*, C=N); 160.8 (*s*, C(5')); 132.4 (*s*, 1 arom. C); 131.6, 128.8, 128.1 (3d, 5 arom. CH); 115.1 (*s*, C(CO₂Me)₂); 85.0 (*s*, C(4')); 52.7, 52.6 (2*q*, 2 MeO); 26.9 (*q*, 2 Me). EI-MS: 319 (60 *M*⁺), 304 (100), 288 (17), 272 (60), 260 (60), 216 (22), 184 (53), 169 (15), 154 (45), 152 (85), 145 (36), 104 (32), 73 (42). Anal. calc. for C₁₆H₁₇NO₄S (319.17): C 60.17, H 5.37, N 4.39, S 10.04; found: C 59.97, H 5.46, N 4.21, S 10.21.

REFERENCES

- [1] R. M. Kellogg, *Tetrahedron* **1976**, *32*, 2165.
- [2] R. Huisgen, C. Fulka, I. Kalwinski, L. Xi, G. Mlostoń, J. R. Moran, A. Pröbstl, *Bull. Soc. Chim. Belg.* **1984**, *93*, 511.
- [3] L. Fišera, R. Huisgen, I. Kalwinski, E. Langhals, X. Li, G. Mlostoń, K. Polborn, J. Rapp, W. Sicking, R. Sustmann, *Pure Appl. Chem.* **1996**, *68*, 789.
- [4] V. Alcazar, I. Tapia, J. R. Moran, *Tetrahedron* **1990**, *46*, 1057; J. R. Moran, I. Tapia, V. Alcazar, *ibid.* **1990**, *46*, 1783.
- [5] G. Mlostoń, A. Linden, H. Heimgartner, *Helv. Chim. Acta* **1991**, *74*, 1386.
- [6] G. Mlostoń, A. Linden, H. Heimgartner, *Helv. Chim. Acta* **1996**, *79*, 31.
- [7] J. Buter, S. Wassenaar, R. M. Kellogg, *J. Org. Chem.* **1972**, *37*, 4045.
- [8] G. Mlostoń, M. Petit, A. Linden, H. Heimgartner, *Helv. Chim. Acta* **1994**, *77*, 435; M. Petit, A. Linden, G. Mlostoń, H. Heimgartner, *ibid.* **1994**, *77*, 1076.
- [9] I. Kalwinski, X. Li, J. Gottstein, R. Huisgen, *J. Am. Chem. Soc.* **1981**, *103*, 7032.
- [10] G. Mlostoń, J. Romański, E. B. Rusanov, A. N. Tshernega, J. G. Shermolovich, *Russian J. Org. Chem.* **1995**, *31*, 1027.
- [11] M. Kägi, A. Linden, H. Heimgartner, G. Mlostoń, *Helv. Chim. Acta* **1993**, *76*, 1715.
- [12] R. Huisgen, G. Mlostoń, *Heterocycles* **1990**, *30*, 737.

⁵) This compound decomposed during storage and was not obtained in anal. pure form.

- [13] I. Kalwinski, R. Huisgen, *Tetrahedron Lett.* **1981**, 22, 3941.
- [14] M. Kägi, G. Mlostoń, A. Linden, H. Heimgartner, *Helv. Chim. Acta* **1994**, 77, 1299; M. Kägi, A. Linden, G. Mlostoń, H. Heimgartner, *ibid.* **1996**, 79, 855.
- [15] M. Kägi, unpublished results.
- [16] G. Mlostoń, T. Gendek, H. Heimgartner, *Helv. Chim. Acta* **1996**, 79, 1537.
- [17] a) A. Padwa, F. Hornbuckle, *Chem. Rev.* **1991**, 91, 263; b) 'Houben-Weyl, Methoden der Organischen Chemie', Ed. M. Regitz, Thieme Verlag, Stuttgart, 1989, Vol. E19b.
- [18] B. W. Peace, D. S. Wulfman, *Synthesis* **1973**, 137.
- [19] E. A. Shapiro, A. B. Dyatkin, O. M. Nefedov, *Russian Chem. Rev.* **1993**, 62, 485.
- [20] H. Zollinger, 'Diazo Chemistry II', VCH Verlagsgesellschaft mbH, Weinheim, 1995.
- [21] R. Huisgen, E. Langhals, *Tetrahedron Lett.* **1989**, 30, 5369.
- [22] M. N. Elison, S. K. Fedukovitch, G. I. Nikishin, *Izv. Akad. Nauk SSSR, Ser. Khim.* **1990**, 2783.
- [23] J. A. Kaufman, S. J. Weininger, *J. Chem. Soc., Chem. Commun.* **1969**, 593.
- [24] N. Tokitoh, T. Suzuki, A. Itami, M. Goto, W. Ando, *Tetrahedron Lett.* **1989**, 30, 1249.
- [25] G. Mlostoń, R. Huisgen, *Heterocycles* **1985**, 23, 2201.
- [26] S. Takano, S. Tomita, M. Takahashi, K. Ogasawara, *Synthesis* **1987**, 1116.
- [27] T. Honda, H. Ishige, J. Araki, S. Akimoto, K. Hirayama, M. Tsubuki, *Tetrahedron* **1992**, 48, 79.
- [28] M. Roth, P. Dubs, E. Götschi, A. Eschenmoser, *Helv. Chim. Acta* **1971**, 54, 710.
- [29] B. S. Pedersen, S. Scheibye, N. H. Nilsson, S.-O. Lawesson, *Bull. Soc. Chim. Belg.* **1978**, 87, 223.
- [30] C. Jenny, H. Heimgartner, *Helv. Chim. Acta* **1986**, 69, 374; P. Wipf, C. Jenny, H. Heimgartner, *ibid.* **1987**, 70, 1001.
- [31] B. W. Peace, F. Carman, D. S. Wulfman, *Synthesis* **1971**, 658.