

Synthesis of N^2 -Alkyl-8-oxo-7,8-dihydro-2'deoxyguanosine Derivatives and Effects of These **Modifications on RNA Duplex Stability**

Arunkumar Kannan and Cynthia J. Burrows*

Department of Chemistry, University of Utah, 315 South 1400 East, Salt Lake City, Utah 84112-0850, United States

burrows@chem.utah.edu

Received November 2, 2010



 N^2 -Alkyl analogues of 8-oxo-7,8-dihydro-2'-deoxyguanosine (OG) were synthesized (alkyl = propyl, benzyl) via reductive amination of the protected OG nucleoside and incorporated into various positions of an RNA strand. Thermal stability studies of duplexes containing A or C opposite a single modified base revealed only moderate destabilization. Both OG as well as its N^2 -alkyl analogues can pair opposite A or C with nearly equal stability, potentially offering a new means of modulating RNA-protein interactions in the minor vs major grooves.

Chemical modification of nucleosides has been a successful strategy for antiviral,¹ antimetabolite,² antitumor,^{3,4} and diagnostic agents.⁴ Modified nucleosides in oligomers exist naturally due to cellular reactions on both the bases and the sugars of DNA and RNA and have also been introduced into oligomers synthetically. Modified nucleosides are common in the RNA field due to applications in investigating reaction mechanisms,^{5,6} imparting favorable properties on siRNAs,⁷

- (1) Duong, A.; Mousa, S. A. Drugs Today 2009, 45, 751-761.
- (2) Foss, F. M. Best. Pract. Res. Clin. Haematol. 2004, 17, 573-584.
- (3) Jordheim, L. P.; Dumontet, C. Biochim. Biophys. Acta, Rev. Cancer 2007, 1776, 138-159.
- (4) Kassis, A. I.; Adelstein, S. J.; Mariani, G. Q. J. Nucl. Med. 1996, 40, 301-319
- (5) Maydanovych, O.; Easterwood, L. M.; Cui, T.; Veliz, E. A.; Pokharel, S.; Beal, P. A. Methods Enzymol. 2007, 424, 369-386.
- (6) Suydam, I. T.; Strobel, S. A. J. Am. Chem. Soc. 2008, 130, 13639-13648.
- (7) Terrazas, M.; Kool, E. T. Nucleic Acids Res. 2009, 37, 346-353. (8) Watts, J. K.; Deleavey, G. F.; Damha, M. J. Drug Discov. Today 2008,
- 13.842-855.
- (9) Rist, M. J.; Marino, J. P. Curr. Org. Chem. 2002, 6, 775-793.
- (10) Hougland, J. L.; Piccirilli, J. A. Methods Enzymol. 2009, 468, 107-125.

720 J. Org. Chem. 2011, 76, 720–723

probing RNA structure and function,⁹⁻¹¹ and exploring interactions between RNA and proteins or small molecules.

Both sugar and backbone modifications have been explored to improve nuclease stability and target identification in antisense and siRNA approaches;^{12,13} however, examples of base modification in these applications are few by comparison.^{14–16} The ability to alter the hydrophobicity and steric properties of the major and minor groove appeared to us as a possible means of modulating both interstrand interactions as well as nucleic acid-protein interactions, 7,17,18 thereby expanding the range of nucleic acid modifications in the design of DNA and RNA therapeutics.

Inspiration for the present work came from one of the major products of oxidative damage to the DNA, namely 8-oxo-7,8-dihydro-2'-deoxyguanosine (OG).¹⁹⁻²¹ Introduction of an oxo group at C8 of the purine increases the propensity of the purine to flip from the normal anti conformation to syn, where it exposes the Hoogsteen face of the purine to base pairing. Like its parent guanine, OG(anti) accepts cytosine as a Watson-Crick partner, while the complementary base for OG(syn) is adenosine (Figure 1). During the anti-syn conformational change, the N^2 -amino group and the C8-oxo groups exchange positions between the minor and major grooves (Figure 1). As a result, addition of an alkyl or aryl group to the exocyclic amine should enable the placement of the substituent in either groove, in a way that will be governed by the identity of the base opposite.

OG can be synthetically incorporated into DNA and RNA oligomers via the corresponding phosphoramidite. While we, among others, have been studying OG and its incorporation into oligomers, including chemical characterization and enzymology with DNA processing enzymes, 22,23 OG is rarely studied in RNA. Thermal denaturation studies,^{22,24,25} NMR studies,²⁶ and X-ray crystallographic

- (13) Rozners, E. *Curr. Org. Chem.* 2006, *10*, 675–692.
 (14) Chiu, Y. L.; Rana, T. M. *RNA* 2003, *9*, 1034–1048.
 (15) Somoza, A.; Silverman, A.; Miller, R.; Chelliserrykattil, J.; Kool, E.
- Chem. Eur. J. 2008, 14, 7978–7987.
 (16) Xia, J.; Noronha, A.; Toudjarska, I.; Li, F.; Akinc, A.; Braich, R.; Frank-Kamenetsky, M.; Rajeev, K. G.; Egli, M.; Manoharan, M. ACS Chem. Biol. 2006, 1, 176–183.
- (17) Peacock, H.; Fostvedt, E.; Beal, P. A. ACS Chem. Biol. 2010, 5, 1115-1124.
- (18) Liu, Y.; Xu, J.; Karimiahmadabadi, M.; Zhou, C.; Chattopadhyaya, J. J. Org. Chem. 2010, 75, 7112–7128.

(19) Wood, M. L.; Esteve, A.; Morningstar, M. L.; Kuziemko, G. M.; Essigmann, J. M. Nucleic Acids Res. 1992, 20, 6023-6032.

- (20) Wang, D.; Kreutzer, D. A.; Essigmann, J. M. Mutat. Res., Fundam. Mol. Mech. Mutagen. 1998, 400, 99-115.
- (21) Ames, B. N.; Shigenaga, M. K.; Hagen, T. M. Proc. Natl. Acad. Sci. U.S.A. 1993, 90, 7915-7922.

(22) Burrows, C. J.; Muller, J. G.; Kornyushyna, O.; Luo, W.; Duarte, V.; Leipold, M. D.; David, S. S. Environ. Health Perspect. 2002, 110 (Suppl 5), 713-717

- (23) Krahn, J. M.; Beard, W. A.; Miller, H.; Grollman, A. P.; Wilson, S. H. Structure 2003, 11, 121-127.
- (24) Ober, M.; Mueller, H.; Pieck, C.; Gierlich, J.; Carell, T. J. Am. Chem. Soc. 2005, 127, 18143-18149.

(25) Hamm, M. L.; Billig, K. Org. Biomol. Chem. 2006, 4, 4068–4070.
(26) Kouchakdjian, M.; Bodepudi, V.; Shibutani, S.; Eisenberg, M.; Johnson, F.; Grollman, A. P.; Patel, D. J. Biochemistry 1991, 30, 1403–1412.

Published on Web 12/30/2010

DOI: 10.1021/jo102187y © 2010 American Chemical Society

⁽¹¹⁾ Sinkeldam, R. W.; Greco, N. J.; Tor, Y. Chem. Rev. 2010, 110, 2579–2619.

⁽¹²⁾ Shukla, S.; Sumaria, C. S.; Pradeepkumar, P. I. ChemMedChem 2010, 5, 328-349.



FIGURE 1. 8-Oxo-7,8-dihydro-2'-deoxyguanosine (OG) pairing with A and C.

structures^{23,27,28} of the OG(syn):A(anti) base pair show that the B-form DNA helix is only very slightly perturbed by the presence of this noncanonical pair. Because the DNA–RNA duplex synthesized during transcription is A-form, the transcription studies suggest that both the Watson–Crick OG(anti):C(anti) and the OG(syn):A(anti) base pairs can be stably accommodated in A-form RNA.^{29,30}

In the G:C or OG:C Watson-Crick base pair, only one hydrogen on the N^2 nitrogen of the purine is involved in hydrogen-bonding to the opposite base; the other hydrogen projects into the minor groove. In the OG-A base pair, the entire exocyclic amino group of OG projects into the major groove. Thus, replacement of one of the hydrogens of OG's N^2 group by an alkyl should have a very minimal effect on the hydrogen bonding ability in a duplex. However, the presence of a bulky substituent could have other effects on local solvation or steric interactions that are not easily predicted. Therefore our current focus was to study the effects of different substitutions that vary in steric bulk at the N^2 position of OG on the stability of duplex formation in RNA. Chemical modifications in siRNA are an important way to address issues such as stability, delivery and reducing off-target effects.^{14,31,32} The ability of OG to base pair in two alternative conformations, syn and anti, could play a major role in siRNA therapeutics. The switching of steric bulk between minor (OG(anti)) vs major (OG(syn)) grooves may prevent undesirable interactions depending on pairing with C or A, respectively, in the complementary strand.

In previous work, we reported the stability of OG and its oxidation products in DNA strands.²² The Beal laboratory has shown that chemical modification of Gs in caspase-2 siRNAs can reduce their ability to bind to off-target proteins such as Protein Kinase R (PKR) while maintaining caspase-2 gene knockdown capability.³³ Hence, in our current experiments, we have synthesized various N^2 -alkyl derivatives of OG phosphoramidites and incorporated them into a caspase-2 siRNA sequence by replacing selected uracils with

the modified OGs. The caspase-2 sequence selected has Us situated near the 5' end, near the 3' end and in the middle of the guide strand sequence. Accordingly, we chose positions 4 (5' seed region), 11 (middle; at the Argonaute2 cleavage site) and 16 (near the 3' terminus) for incorporation of modified bases (vide infra). These three positions would allow us to understand the effects of the modified nucleosides on stability of RNA duplexes in detail. Furthermore, by forming the RNA duplex with either A or C opposite modified guanosines, we could evaluate the duplex stability with both Watson-Crick (OG:C) and Hoogsteen (OG:A) base pairs in these three regions. The 2'-deoxynucleoside of OG was selected for study because the absence of the 2'-OH group might readily facilitate both the syn and anti conformations in comparison to the ribonucleoside analogue.³⁰ Previous studies have shown that the presence of a few 2'-deoxynucleosides in an RNA strand does not dramatically alter its physical or biological properties.¹⁴ In the current work, the N^2 substituents are propyl or benzyl in comparison to H to evaluate a small vs medium-sized hydrophobic group added to either the major or minor groove.

Initially, attempts were made to follow a synthetic scheme similar to that of alkyl-substituted 2-aminopurines by substitution of the diacylated-2-bromo-2'-deoxyinosine nucleoside.¹⁷ Unfortunately, substitution of 2-bromoinosine with ethyl, propyl and benzyl amine all proceeded in low yields. Furthermore, diazotization of deoxyguanosine to yield the 2-bromo-2'-deoxyinosine base led to many different products as in previous work.^{34,35}

As a second approach, we attempted reductive amination of aldehydes with guanosine; reaction of 2'-deoxyguanosine with the respective aldehyde in the presence of NaBH₃CN led to a \sim 60% yield along with a small amount of dialkylated side product (\sim 20%). However, subsequent conversion into 8-oxo-7,8-dihydro-2'-deoxyguanosine yielded many side products complicating the purification.

Thus, the most promising procedure was to install the 8-oxo moiety first and then carry out reductive amination to prepare the N^2 -alkyl derivatives of OG; these were then finally converted into the corresponding phosphoramidites. The overall procedure, shown in Scheme 1, involved conversion of 8-bromo-2'-deoxyguanosine (2) to 8-O-benzyl-2'deoxyguanosine (3) in the presence of sodium benzylate and DMSO. Compound 3 is highly unstable in acidic conditions and yields OG (4) upon stirring with 1 M HCl for 2 h. Reductive amination of OG in the presence of sodium cyanoborohydride in 20% aqueous methanol and the appropriate aldehyde yielded the respective N^2 -alkylated products (5) in good yield. Modest changes in the concentration of NaBH₃CN or the aldehyde did not alter the yield of the monoalkylated or dialkylated products (5 vs 6) significantly; however, a large excess of the aldehyde and longer reaction times yielded predominately the dialkylated compound. The mono- N^2 -alkylated product was purified chromatographically, and the 5'-OH was protected by DMTrCl in pyridine to yield (7). This compound was then converted into its phosphoramidite (8) using triethylamine as base in CH₂Cl₂ under anhydrous conditions.³⁶ The modified phosphoramidites

⁽²⁷⁾ Brieba, L. G.; Eichman, B. F.; Kokoska, R. J.; Doublie, S.; Kunkel, T. A.; Ellenberger, T. *EMBO J.* **2004**, *23*, 3452–61.

⁽²⁸⁾ Hsu, G. W.; Ober, M.; Carell, T.; Beese, L. S. *Nature* 2004, 431, 217–221.
(29) Kim, S. K.; Lee, S. H.; Kwon, O. S.; Moon, B. J. J. Biochem. Mol.

 ⁽²⁰⁾ Kin, S. K. 262.
 (30) Kim, S. K.; Yokoyama, S.; Takaku, H.; Moon, B. J. *Biolog. Med.*

Chem. Lett. **1998**, *8*, 939–944. (31) Chiu, Y.-L.; Ali, A.; Chu, C.-y.; Cao, H.; Rana, T. M. Chem. Biol. **2004**, *11*, 1165–1175.

⁽³²⁾ Chiu, Y. L.; Rana, T. M. Mol. Cell 2002, 10, 549-561.

 ⁽³³⁾ Puthenveetil, S.; Whitby, L.; Ren, J.; Kelnar, K.; Krebs, J. F.; Beal,
 P. A. Nucleic Acids Res. 2006, 34, 4900–4911.

⁽³⁴⁾ Suzuki, T.; Yamaoka, R.; Nishi, M.; Ide, H.; Makino, K. J. Am. Chem. Soc. 1996, 118, 2515–2516.

⁽³⁵⁾ Glaser, R.; Rayat, S.; Lewis, M.; Son, M.-S.; Meyer, S. J. Am. Chem. Soc. **1999**, *121*, 6108–6119.



were introduced at either one, two or three positions via standard solid-phase synthesis. Coupling efficiencies for the modified phosphoramidites were similar to those of unmodified bases. The deprotected oligonucleotides were purified by ion-exchange chromatography, analyzed by negative ion ESI-MS, and hybridized to form RNA duplexes.

The effect of the 2'deoxy-OG nucleosides with and without N^2 -alkyl substituents on duplex stability was investigated via thermal denaturation (T_M) studies at pH 7.4 in 100 mM NaCl. In the initial set of experiments, the nucleoside analogs were incorporated into the duplex at each of three different positions (4, 11 or 16) of the guide strand sequence from caspase 2 with C, A and G in the strand opposite (Figure 2). The modified strands containing OG and its analogues were also compared with the unmodified strand containing a U:A base pair. In the second set of experiments, the nucleoside analogs were incorporated into the duplex at more than one position, that is, (4,11), (11,16), (4,16), or (4,11,16) of the antisense strands opposite A or C (Figure 3).

 $T_{\rm M}$ measurements showed that replacement of a single U:A base pair with a single O:C (O = OG with R = H) at positions 16 or 11 decreased the $T_{\rm M}$ by ~2 °C whereas at position 4 it decreased by 5 °C. Indeed, position 4 proved to be consistently more sensitive to the effects of modified bases. Introduction of O:A at the same positions showed a further slight decrease in $T_{\rm M}$ of ~1 °C. However, on introduction of guanosine opposite the modified bases, which is expected to be a destabilizing mispair, the $T_{\rm M}$ decreased considerably ($\Delta T_{\rm m} \approx 8$ °C). In DNA, the OG(syn):G(anti) base pair can form, but is somewhat less stable;³⁷ therefore, it was of interest to know the likelihood of mispairing with G if OG was present in the guide strand of an siRNA.





FIGURE 2. T_M values of RNA duplexes singly modified at positions 4, 11, or 16 of the caspase-2 guide sequence compared to native U:A base pairs in these positions. The passenger strand has either Y = C (green bars), A (blue bars), or G (red bars) opposite the modified site X, in which the R group on N^2 of OG is H (X = O), propyl (X = P), or benzyl (X = B).



FIGURE 3. $T_{\rm M}$ values of RNA duplexes doubly and triply modified at positions 4, 11 and/or 16 of the caspase-2 guide sequence compared to native U:A base pairs in these positions. The passenger strand has either Y = C (green bars) or A (blue bars) opposite the modified site X, in which the R group on N^2 of OG is H (X = O), propyl (X = P), or benzyl (X = B).

Introduction of alkyl substituents onto the N^2 position of OG, either propyl (P) and benzyl (B), showed similar $T_{\rm M}$ patterns, of which the benzyl modification was somewhat more destabilizing than the propyl analogue or OG itself. In brief, the order of stability of these duplex oligonucleotides can be listed as: Unmodified > O > P > B. From the pattern observed from the data, it was clear that the larger the N^2 group, the greater the destabilizing effect, although all duplexes with single modifications were still quite stable $(T_{\rm M} \approx 60 \,^{\circ}{\rm C})$. The small difference in $T_{\rm M}$ between the pairing against A and C further supports the concept that OG can switch conformations according to the complementary base, projecting the N^2 substituent into either the major or minor groove. Interference with RNA-protein contacts at these sites is expected to disrupt complexes with double-stranded RNA binding motifs in key proteins.

⁽³⁶⁾ Bodepudi, V.; Shibutani, S.; Johnson, F. Chem. Res. Toxicol. 1992, 5, 608–617.

⁽³⁷⁾ Thiviyanathan, V.; Somasunderam, A.; Hazra, T. K.; Mitra, S.; Gorenstein, D. G. *J. Mol. Biol.* **2003**, *325*, 433–442.

In the next set of experiments (Figure 3), the alkylated OGs were incorporated into the duplex at 2 or 3 positions of the caspase-2 guide strand sequence with either As or Cs in the strand opposite. As expected, this decreased the stability even further with most of the T_M values of strands having modifications at more than one position being around 55 °C. These strands still form stable duplexes, however, and the differences in T_M values between pairing against A or C remained very similar.

In conclusion, we have developed an optimized route to N^2 -alkyl derivatives of 8-oxo-7,8-dihydro-2'deoxyguanosine phosphoramidites and incorporated them into the caspase-2 siRNA guide sequence at one, two or three positions in which a U site was replaced by an OG nucleotide. The thermal stabilities of duplexes containing OG(anti):C(anti) vs OG(syn):A(anti) were consistent with the conclusion that OG can exist in either the syn or anti orientation depending on the identity of the base opposite. Introduction of alkyl groups onto the exocyclic amino group led to only modest decreases in stability. No major sequence effects were evident although modification at position 4, a 5'GXU-3' sequence context, led to slightly greater destabilizing effects than at positions 11 (5'-UXG-3') or 16 (5'-UXC-3'). The overall stabilities of the duplexes and the small differences in stabilities of N^2 -alkyl-8-oxo-2'-deoxyguanosine on pairing with A and C suggests that these modifications can be successfully incorporated into siRNA duplexes with a view to switching between modes that might prevent unwanted protein interactions while permitting silencing activity.

Experimental Section

*N*²**-Propyl-8-oxo-7,8-dihydro-2'-deoxyguanosine** (**5**a). To a suspension of $4^{3,31}$ (950 mg, 3.3 mmol) and NaBH₃CN (628 mg, 9.9 mmol) in methanol (80 mL) was added propanal (8.16 mL, 114 mmol) in one portion, and the mixture was heated at 50 °C overnight under nitrogen. After removal of the solvent under reduced pressure, the residue was purified by column chromatography using 20% MeOH–CH₂Cl₂ to yield 805 mg (75%) of **5a**. ¹H NMR ((CD₃)₂SO): δ 10.4–9.9 (2 H, bs), 6.95 (1 H, bs), 6.03 (1 H, t, *J* = 6.04 Hz), 5.17 (1 H, bs), 5.7 (1 H, bs), 4.36–4.30 (1H, m), 4.31–4.27 (1H, m), 3.58–3.47 (1H, m), 3.65–3.52 (2H), 3.42–3.31 (1 H, m), 3.04–2.76 (1H, m), 1.88–1.88 (1 H, m), 0.89 (3H, t, *J* = 7.5 Hz) ¹³C NMR ((CD₃)₂SO): δ 151.7, 146.9, 124.2, 98.5, 94.2, 87.1, 80.7, 71.2, 64.1, 62.4, 42.2, 35.6, 22.1, 11.3. ESI-MS (*m*/*z*): calcd for C₁₃H₁₉N₅O₅Na (M + Na⁺) 348.1284; found 348.1286.

5'-O-4,4'-Dimethoxytrityl-N²-propyl-8-oxo-7,8-dihydro-2'deoxyguanosine (7a). To a solution of 5a (400 mg, 1.23 mmol) in dry pyridine (6 mL) cooled in ice water was added 4,4'-DMTrCl (500 mg, 1.47 mmol). The cooling bath was removed and stirring was continued at RT for 15 min, followed by cooling in ice and quenching with water (50 mL). The solution was extracted with CH_2C1_2 (5 \times 20 mL), and the combined organic layers were washed with H₂O (2×20 mL) and then dried over MgSO₄. The solvent was evaporated under reduced pressure, and the crude residue was purified by column chromatography over silica gel using MeOH-CH₂Cl₂ containing 2% triethylamine to inhibit detritylation yielding 7a (385 mg, 50%). ¹H NMR (CD₂Cl₂): δ 10.22 (2 H), 7.50-6.89 (13 H, m), 6.41(1 H, bs), 6.21 (1 H, t, J =6.05 Hz), 4.55-4.51 (1H, m), 4.01-3.98 (1H, m), 3.88-3.80 (1H, m], 3.86 and 3.85 (6 H, 2 s), 3.57-3.30 (2 H, 2 m), 3.18-3.06 (2 H, m), 2.23-2.19 (2 H, m), 1.50-1.43 (2 H, m), 0.89 (3H, t, J = 7.5 Hz); ¹³C NMR (CD₂Cl₂): δ 176.4, 154.5, 152.4, 147.6, 140.6, 128.9, 128.8, 127.7, 127.5, 127.3, 99.5, 87.6, 81.4, 71.9, 63.0, 44.4, 43.4, 36.0, 25.3. ESI-MS (m/z): calcd for C₃₄H₃₇N₅O₇ Na $(M + Na^+)$ 650.2591; found 650.2604.

3'-O-[(Diisopropylamino)-(2-cyanoethoxy)phosphino]-5'-O-(4,4'-dimethoxytrityl)-N²-propyl-8-oxo-7,8-dihydro-2'-deoxyguanosine (8a). To a mixture of 7a (0.112 g, 0.18 mmol) dried over P2O5 in a vacuum desiccator for 24-48 h and then coevaporated with 1:1 dry CH₂Cl₂ and benzene prior to reaction, plus dry Et₃N (0.044 g, 0.43 mmol) in dry CH₂Cl₂ (1 mL) under N₂ was added 2-cyanoethyl N,N-diisopropyl-phosphoramidochloridite (0.050 g, 0.25 mmol). The reaction was monitored by TLC analysis for 20 min; the solvent was evaporated, and a mixture of dry THF-benzene (1:4; 25 mL) was introduced, stirred for 10 min and filtered under N₂ to remove the $Et_3N \cdot HCl$. The process was then repeated twice using dry benzene as the solvent. The material was purified by column chromatography using 5% MeOH-CH2Cl2 containing 2% triethylamine. The resultant viscous foamy material, when dried over P_2O_5 in a vacuum desiccator at RT overnight, led to a yield of 105 mg of 8a (70%). ¹H NMR (CD₂Cl₂): 7.42-6.74 (13H, m), 6.25 (1H, t, J = 6.05 Hz), 4.72 (1H, m), 4.21-4.08 (2H, m), 3.79(6H, s), 3.59-3.52 (4H, m,), 3.42-3.14 (2H, m), 3.12-3.01 (4H, m), 2.75–2.70 (2H, m), 2.40–2.23 (2H, m), 1.48–1.05 (14H, m), 0.85, (3H, t, J = 7.5). ¹³C NMR (CD₂Cl₂): δ 158.8, 152.9, 149.2, 147.9, 136.3, 130.3, 128.5, 128.4, 127.9, 126.9, 113.2, 98.6, 86.2, 81.3, 64.2, 58.6, 55.6, 55.4, 45.5, 43.3, 24.6, 22.6, 20.7, 11.6. ³¹P NMR (CD₂Cl₂): 147.69, 147.27. ESI-MS (*m*/*z*): calcd for $C_{43}H_{54}N_7O_8PNa (M + Na^+) 850.3669$; found 850.3662.

Acknowledgment. We thank Prof. P. A. Beal (UC, Davis) for valuable comments and the NIH for financial support (GM080784).

Supporting Information Available: Additional experimental procedures, copies of ¹H, ¹³C and ³¹P NMR spectra for new nucleoside derivatives, sequence information, $T_{\rm m}$ curves and mass spectral data for oligomers. This material is available free of charge via the Internet at http://pubs.acs.org.