

Nickel-Catalyzed Decarbonylative Addition of Anhydrides to Alkynes

Yuichi Kajita, Takuya Kurahashi,* and Seijiro Matsubara*

Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Kyoto 615-8510, Japan

Received August 19, 2008; E-mail: matsubar@orgxn.mbox.media.kyoto-u.ac.jp; tkuraha@orgxn.mbox.media.kyoto-u.ac.jp

The transition metal-catalyzed insertion reaction of an unsaturated carbon–carbon bond into a carbon–oxygen bond of an oxacyclic compound is a useful transformation to prepare more complicated oxacyclic compounds in a single step (Scheme 1). However, there is no precedent for such a potentially valuable methodology. We postulated that oxidative addition of cyclic anhydrides to nickel(0) and subsequent decarbonylation are facile and might allow insertion of alkynes. Thus, we attempted the decarbonylative addition of phthalic anhydride to alkyne using nickel(0) catalyst to form isocoumarin,^{1,2} which displays a wide range of biological activities.³


Our investigation began with an attempted addition of phthalic anhydride (**1a**) to 4-octyne (**2a**) with 10 mol % of Ni(cod)₂ and 40 mol % of PMe₃ in refluxing acetonitrile (80 °C) for 12 h.⁴ However, this resulted in low conversion of **1a** and led to isocoumarin **3aa** in 12% yield (Table 1, entry 1).⁵ A detailed examination of the reaction conditions revealed that, on addition of 20 mol % of ZnCl₂, the reaction proceeded smoothly to furnish **3aa** in 96% isolated yield (entry 2).⁶ Among the Lewis acids examined, ZnCl₂ gave the best yield of the product (entries 3–7). An addition of quaternary ammonium salts was also found to be effective for the reaction (entries 8–12). An electrostatic interaction between the carbonyl moiety of **1a** and the quaternary ammonium might behave as a general Lewis acid.⁷

With the optimized conditions in hand,^{8–10} we next investigated the scope of the reaction using various anhydrides (Table 2). The decarbonylative addition of maleic anhydride **1b** to 4-octyne provides α -pyrone **3ba** in 87% yield (Table 2, entry 1). The reaction of 2,3-naphthalenedicarboxylic anhydride (**1c**) with **2a** gave the product **3ca** in 82% yield (entry 2). Unsymmetrical anhydrides **1d** and **1e** provide the corresponding products **3da** and **3ea** in 61% and 53% yields, along with formation of the respective regioisomers (entries 3 and 4). The addition of **1f** to **2a** proceeded smoothly to provide **3fa** in 94% yield (entry 5). The addition of diphenyl-substituted maleic anhydride **1g** to 4-octyne resulted in lower yield (23%) even with prolonged reaction time (entry 6).

The addition reaction is also compatible with aryl-substituted alkyne (Table 3, entry 1) and afforded the corresponding isocoumarin **3ab** in 87% yield. The reaction of **1a** with unsymmetrical alkynes such as **2c** and **2d** gave the products consisting of regioisomers in a 1/1 ratio (entries 2 and 3). The addition of **1a** to **2e**, containing sterically hindered groups, gave adduct **3ae** as a major product with a regioselectivity ratio of 2/1 (entry 4). Bulky and electron-rich trimethylsilyl-substituted alkynes such as **2f** and **2g** reacted with **1a** to provide adducts with complete regiocontrol in excellent yields (entries 5 and 6). Terminal alkynes, such as 1-octyne and phenylacetylene, failed to participate in the reaction, presumably due to rapid oligomerization of alkynes.















A plausible reaction pathway to account for the formation of isocoumarins **3** based on the observed results is outlined in Scheme 2. In view of the mechanism of the previously reported nickel-catalyzed cross-coupling reaction of anhydrides with organometallic reagents,¹¹ it is reasonable to consider that the catalytic cycle of

Table 1. Nickel-Catalyzed Decarbonylative Addition of Phthalic Anhydride to 4-Octyne^a

					
entry	additive	yield (%) ^b	entry	additive	yield (%) ^b
1	—	12	7	LiCl	76
2	ZnCl ₂	96	8	Bu ₄ NPF ₆	54
3	Zn(OTf) ₂	92	9	Bu ₄ NCl	72
4	ZnBr ₂	91	10	Bu ₄ NOTf	67
5	ZnI ₂	90	11	Bu ₄ NBr	65
6	BPh ₃	92	12	Bu ₄ NI	52

^a All reactions were carried out using Ni(cod)₂ (10 mol %), PMe₃ (40 mol %), additive (20 mol %), phthalic anhydride (0.5 mmol), and 4-octyne (0.6 mmol) in 2 mL of refluxing acetonitrile (80 °C). ^b Isolated yields.

Table 2. Nickel-Catalyzed Decarbonylative Addition of Anhydrides to 4-Octyne^a

entry	1	time (h)	3 (yield ^b)
1		15	 3ba (87%)
2		18	 3ca (82%)
3		18	 3da (61%)  3da' (29%)
4		18	 3ea (53%)  3ea' (8%)
5		15	 3fa (94%)
6		18	 3ga (23%)



^a All reactions were carried out using Ni(cod)₂ (10 mol %), PMe₃ (40 mol %), ZnCl₂ (20 mol %), **1** (0.5 mmol), and **2a** in 2 mL of refluxing acetonitrile (80 °C). ^b Isolated yields.

the present reaction should consist of oxidative addition of an anhydride O–CO bond to nickel.¹² Subsequent decarbonylation and coordination of alkyne take place, in which the steric repulsive interaction is minimal between the bulkier R¹ and the PMe₃ ligand on the nickel, to give nickel(II) intermediate **4a**. The alkyne would then insert into the aryl-nickel bond to give nickelacycle **5**, which undergoes reductive elimination to give **3** and regenerates the

Table 3. Nickel-Catalyzed Decarbonylative Addition of Phthalic Anhydride to Alkynes^a

entry	2 (eq)	time (h)	3	yield (%) ^b
1	2b (1.5)	12	3ab	87
2	2c (1.5)	15	3ac	91 (1/1) ^c
3	2d (1.5)	15	3ad	75 (1/1) ^c
4	2e (1.5)	15	3ae	89 (2/1) ^c
5	2f (6.0)	18	3af	93 ^d
6	2g (1.5)	24	3ag	86 ^d

^a All reactions were carried out using Ni(cod)₂ (10 mol %), PMe₃ (40 mol %), **1a** (0.5 mmol), and alkyne in 2 mL of refluxing acetonitrile (80 °C). ^b Isolated yields. ^c Ratio of regioisomers. ^d Only a single regioisomer was formed.

Scheme 1. Transition-Metal-Catalyzed Decarbonylative Addition of Cyclic Anhydrides to Alkynes**Scheme 2.** Plausible Mechanism for the Nickel-Catalyzed Decarbonylative Addition of Phthalic Anhydrides to Alkynes

starting nickel(0). In-situ IR spectra analysis demonstrated that the stoichiometric reaction of Ni(cod)₂/PMe₃ with **1a** and **2a** without ZnCl₂ resulted in gradual consumption of **1a** without formation of **3aa**. Importantly, constant generation of **3aa** was observed simultaneously on addition of ZnCl₂ (Figure S1, Supporting Information). These results imply that reductive elimination is specifically promoted by the addition of ZnCl₂. The origin of the effect of ZnCl₂ is likely to result from the coordination of a Lewis acid to a carbonyl

group, which may generate electron-poor alkenylnickel through a conjugated system.⁵ⁱ

In conclusion, the decarbonylative addition reaction of phthalic anhydrides to alkynes is successfully demonstrated using a nickel catalyst in association with a Lewis acid as a cocatalyst. Further efforts to expand the scope of the chemistry and studies of the detailed mechanism are currently underway in our laboratories.

Acknowledgment. This work was supported by Grants-in-Aid for Young Scientist (B) (19750031) from the Ministry of Education, Culture, Sports, Science, and Technology, Japan. T.K. also acknowledges NOVARTIS Foundation (Japan) for the Promotion of Science, and the Takeda Pharmaceutical Company Award in Synthetic Organic Chemistry, Japan. We thank Mr. Okada and Mr. Deuchi, Mettler-Toledo, for in situ IR analysis.

Supporting Information Available: Experimental procedures including spectroscopic and analytical data of new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For review of synthesis, see: (a) Barry, R. D. *Chem. Rev.* **1964**, *64*, 229. (b) Jones, J. B.; Pinder, A. R. *J. Chem. Soc.* **1958**, 2612. (c) Napolitano, E. *Org. Prep. Proced. Int.* **1997**, *29*, 631.
- (2) For transition metal catalyzed synthesis, see: (a) Larock, R. C.; Doty, M. J.; Han, X. *J. Org. Chem.* **1999**, *64*, 8770. (b) Hua, R.; Tanaka, M. *New J. Chem.* **2001**, *25*, 179. (c) Nakamura, Y.; Ukita, T. *Org. Lett.* **2002**, *4*, 2317. (d) Subramanian, V.; Batchu, V. R.; Barange, D.; Pal, M. *J. Org. Chem.* **2005**, *70*, 4778. (e) Cherry, K.; Parrain, J.-L.; Thibonnet, J.; Duchêne, A.; Abarbri, M. *J. Org. Chem.* **2005**, *70*, 6669. (f) Kishimoto, Y.; Mitani, I. *Synlett* **2005**, 2141. (g) Fukuyama, T.; Higashibepu, Y.; Yamaura, R.; Ryu, I. *Org. Lett.* **2007**, *9*, 587. (h) Luo, T.; Schreiber, S. L. *Angew. Chem., Int. Ed.* **2007**, *46*, 8250. (i) Woo, J. C. S.; Walker, S. D.; Faul, M. M. *Tetrahedron Lett.* **2007**, *48*, 5679.
- (3) For recent examples, see: (a) Agata, N.; Nogi, H.; Milhollen, M.; Kharbanda, S.; Kufe, D. *Cancer Res.* **2004**, *64*, 8512. (b) Pochet, L.; Frédéric, R.; Masereel, B. *Curr. Pharm. Design* **2004**, *10*, 3781.
- (4) Kajita, Y.; Matsubara, S.; Kurahashi, T. *J. Am. Chem. Soc.* **2008**, *130*, 6058.
- (5) The nickel-catalyzed addition reaction of **1a** with **2a** gave **3aa** in 72% when the reaction was carried out at elevated temperature (250 °C, 6 h, in an autoclave).
- (6) (a) Ikeda, S.; Mori, N.; Sato, Y. *J. Am. Chem. Soc.* **1997**, *119*, 4779. (b) Mahadevan, V.; Getzler, Y. D. Y. L.; Coates, G. W. *Angew. Chem., Int. Ed.* **2002**, *41*, 2781. (c) Kamijo, S.; Yamamoto, Y. *Angew. Chem., Int. Ed.* **2002**, *41*, 3230. (d) Getzler, Y. D. Y. L.; Mahadevan, V.; Lobkovsky, E. B.; Coates, G. W. *J. Am. Chem. Soc.* **2002**, *124*, 1174. (e) Fontaine, F.-G.; Zargarian, D. *J. Am. Chem. Soc.* **2004**, *126*, 8786. (f) Ogoshi, S.; Ueta, M.; Arai, T.; Kurosawa, H. *J. Am. Chem. Soc.* **2005**, *127*, 12810. (g) Rubina, M.; Conley, M.; Gevorgyan, V. *J. Am. Chem. Soc.* **2006**, *128*, 5818. (h) Nakao, Y.; Yada, A.; Ebata, S.; Hiyama, T. *J. Am. Chem. Soc.* **2007**, *129*, 2428. (i) Shen, Q.; Hartwig, J. F. *J. Am. Chem. Soc.* **2007**, *129*, 7734. (j) Nakao, Y.; Kanyiva, K. S.; Hiyama, T. *J. Am. Chem. Soc.* **2008**, *130*, 2448. (k) Baxter, R. D.; Montgomery, J. *J. Am. Chem. Soc.* **2008**, *130*, 9662.
- (7) Orglmeister, E.; Mallat, T.; Baiker, A. *J. Catal.* **2005**, *233*, 333.
- (8) For detailed examination of the reaction conditions, see Supporting Information Table S1.
- (9) The reaction of **1a** and **2a** using air-stable trimethylphosphonium tetrafluoroborate (40 mol %) and *N,N*-diisopropylethylamine (100 mol %) in place of trimethylphosphine under an existence of ZnCl₂ (20 mol %) as an additive also affords **3aa** in 86% isolated yield. For detailed examination of the reaction conditions, see Supporting Information Table S2. Netherton, M. R.; Fu, G. C. *Org. Lett.* **2001**, *3*, 4295.
- (10) While the majority of the addition reactions have been run on only a 0.5 mmol scale, a larger scale reaction (10 mmol scale) also proceeded smoothly even with reduced catalyst loading (5 mol %) without any difficulty (94% isolated yield), which illustrates the robustness of the reaction. In-situ IR spectra analysis revealed that the reaction proceeded exclusively to give the addition product and was completed in 36 h (Figure S2, Supporting Information).
- (11) (a) O'Brien, E. M.; Bercot, E. A.; Rovis, T. *J. Am. Chem. Soc.* **2003**, *125*, 10498. (b) Johnson, J. B.; Bercot, E. A.; Rowley, J. M.; Coates, G. W.; Rovis, T. *J. Am. Chem. Soc.* **2007**, *129*, 2718.
- (12) (a) Trost, B. M.; Chen, F. *Tetrahedron Lett.* **1971**, *12*, 2603. (b) Sano, K.; Yamamoto, T.; Yamamoto, A. *Chem. Lett.* **1984**, 941. (c) Sano, K.; Yamamoto, T.; Yamamoto, A. *Bull. Chem. Soc. Jpn.* **1984**, *57*, 2741. (d) Yamamoto, T.; Sano, K.; Yamamoto, A. *J. Am. Chem. Soc.* **1987**, *109*, 1092. (e) Fischer, R.; Walther, D.; Kempe, R.; Sieler, J.; Schönecker, B. *J. Organomet. Chem.* **1993**, *447*, 131.

JA806569H