


Synthesis of 2-Aryl- and 2-Vinylpyrrolidines via Copper-Catalyzed Coupling of Styrenes and Dienes with Potassium β -Aminoethyl Trifluoroborates

Chanchamnan Um and Sherry R. Chemler*

Department of Chemistry, State University of New York at Buffalo, Buffalo, New York 14260, United States

S Supporting Information

ABSTRACT: 2-Arylpyrrolidines occur frequently in bioactive compounds, and thus, methods to access them from readily available reagents are valuable. We report a copper-catalyzed intermolecular carboamination of vinylarenes with potassium *N*-carbamoyl- β -aminoethyltrifluoroborates. The reaction occurs with terminal, 1,2-disubstituted, and 1,1-disubstituted vinylarenes bearing a number of functional groups. 1,3-Dienes are also good substrates, and their reactions give 2-vinylpyrrolidines. Radical clock mechanistic experiments are consistent with the presence of carbon radical intermediates and do not support participation of carbocations.



Functionalized pyrrolidines are important nitrogen heterocycles found in numerous bioactive compounds of both natural and synthetic origin. 2-Arylpiperidines, in particular, are ubiquitous.^{1–3} The need to access these important moieties and related heterocycles has inspired the development of a number of methods.^{4–26} Many of these methods, however, utilize strong bases⁵ and water-sensitive organometallic reagents.^{7,8} Additional intermolecular metal-catalyzed couplings^{18–20} and intramolecular metal-catalyzed and Bronsted acid catalyzed cyclizations^{21–25} have been developed, providing various routes to 2-arylpiperidines and related products. These latter methods, however, are mainly limited to sulfonamides.


More recently, readily available vinylarenes have been used to directly access 2-arylpiperidines and related saturated heterocycles via intermolecular coupling with bifunctional heteroatom-substituted reagents that can undergo polar/radical [3 + 2]-type bond-forming reaction sequences under mild reaction conditions.^{26–31} The products of these reactions, by virtue of the required substrates, often contain additional functional groups (Schemes 1a,b) that may not be desired in all applications, and primarily *N*-sulfonyl piperidine synthesis has been reported.

To address the existing limitations as well as to explore an orthogonal reactivity mode, we envisaged that a β -aminoethyl carbon radical could serve as a bifunctional three-atom unit to effect a net intermolecular carboamination in the presence of an oxidant (Scheme 1c). The resulting products of the coupling with vinylarenes are simple 2-arylpiperidines whose applications in medicinal chemistry endeavors can be readily envisioned. Herein is presented our development of this method and its extension to 1,3-dienes. The method is ideal for the synthesis of *N*-carbamoylpiperidines. Carbamates are generally considered to be more attractive than sulfonamides as the latter are higher molecular weight and often require more strenuous conditions to reveal the parent amine. The utility of carbamates in medicinal chemistry has been noted.³²


Scheme 1. Polar/Radical Piperidine Syntheses

Previous 2-Arylpiperidine Synthesis from Alkenes:


a. From β -amino aldehydes using SOMO-catalysis



b. From α,β -unsaturated amides via photoredox



c. This Work: Alkene carboamination with β -aminoalkylborates




We recently disclosed an oxidative alkyl Heck-type reaction between alkylboron reagents and vinylarenes.³³ In this transformation, alkyl radicals, generated in situ from [Cu(II)] oxidation of the alkylboron reagent, add to vinylarenes. Oxidation of the resulting benzyl radical then provides the observed higher substituted vinylarene products. We hypothesized that under the reaction conditions the benzylic radical, or carbocation derived thereof, could be intercepted with an amine, resulting in 2-arylpiperidine formation (Scheme 1c). The ready availability of *N*-carbamoyl- β -aminoethylboron reagents, due to the respective contributions of Overman and Molander,^{34,35} presented an excellent opportunity to test this hypothesis.

Copper(II) 2-ethylhexanoate [Cu(eh)₂] is a readily available copper salt that has previously been shown to activate potassium alkyltrifluoroborates in coupling reactions with

Received: April 29, 2016

radical acceptors.^{33,36} However, an attempt at the copper(II) 2-ethylhexanoate-catalyzed coupling of potassium *N*-Cbz- β -aminoethyltrifluoroborate **1a** with 4-methoxystyrene in the presence of MnO₂ (2.55 equiv) as stoichiometric oxidant did not result in pyrrolidine formation (Table 1, entry 1). Upon changing the catalyst to [Cu(1,10-phenanthroline)](OTf)₂,³³ oxidative coupling readily occurred to give 82% yield of 2-arylpyrrolidine **2a** (Table 1, entry 2). The Boc analogue **1b** also provided the coupling product **2b**, but in lower yield (49%, entry 3). The boronic acid analogue **1c** underwent the coupling reaction but also less efficiently (Table 1, entry 4, 52% yield).³⁷ Lowering the copper loading from 20 mol % to 15 mol % led to a less efficient reaction with *N*-Cbz- β -aminoethyltrifluoroborate **1a** (Table 1, entry 5, 67% yield). Ag₂CO₃ (2 equiv) could also serve as a stoichiometric oxidant in place of MnO₂ (Table 1, entry 6).

Table 1. Effect of Alkylborane Structure, Catalyst Loading, and Oxidant on Reaction Efficiency^a




entry	PG	BX _n	oxidant	CuX ₂ (mol %)	yield ^c (%)
1 ^b	Cbz	BF ₃ K 1a	MnO ₂ (2.55 equiv)	Cu(eh) ₂ (20 mol %)	NR
2	Cbz	BF ₃ K 1a	MnO ₂ (2.55 equiv)	Cu(OTf) ₂ (20 mol %)	82
3	Boc	BF ₃ K 1b	MnO ₂ (2.55 equiv)	Cu(OTf) ₂ (20 mol %)	46
4	Cbz	B(OH) ₂ 1c	MnO ₂ (2.55 equiv)	Cu(OTf) ₂ (20 mol %)	52
5 ^d	Cbz	BF ₃ K 1b	MnO ₂ (2.55 equiv)	Cu(OTf) ₂ (15 mol %)	67
6	Cbz	BF ₃ K 1b	Ag ₂ CO ₃ (2 equiv)	Cu(OTf) ₂ (20 mol %)	63

^a25 mol % of 1,1-phenanthroline was used unless otherwise noted. ^b1,10-Phenanthroline was not used. ^cIsolated yield. ^d20 mol % of 1,10-phenanthroline was used. MnO₂ (85% by weight) was used in these reactions.

A number of styrenes underwent this reaction (Scheme 2). Styrenes with electron-donating substituents were most reactive. While most reactions were performed at the 0.125 mmol scale of **1a** (Scheme 2), a 1.25 mmol scale for the efficient (70% yield) production of **2f** was also performed. Ethers, alkyl substituents, sulfonamides and amides were tolerated. Halide-substituted styrenes underwent the reaction, but longer reaction times were required. Substrates bearing two potentially reactive alkenes (terminal styrene, indole, terminal alkyl-substituted alkene) favored reaction at the terminal styrene. A styrene bearing a methyl ester also underwent the reaction but with lower efficiency. A 2-fluoropyridyl in place of a substituted phenyl was also tolerated, although the reaction was less efficient. 1-Phenyl-4-pentene, lacking conjugation, did not react.


Disubstituted styrenes were next evaluated (Scheme 3). 1-Alkyl- and 1-arylstyrenes provided tertiary amines and spirocycles **3a–e**. Coupling with indene gave *cis*-fused bicyclic pyrrolidine **3f**. Both *trans* and *cis*-stilbene gave the same *trans*-pyrrolidine adduct **3g**. 1-Phenyl-1-cyclohexene, a trisubstituted alkene, was unreactive (not shown).

Scheme 2. Annulation with Terminal Styrenes



^aTwo equiv of styrene was used. ^bReaction run for 48 h. ^cReaction run at 95 °C.

Scheme 3. Annulation with Disubstituted Styrenes




^aReaction run in anhydrous 1,4-dioxane at 120 °C. ^bReaction run for 36 h.


Dienes and an enyne also underwent the coupling reaction to give the corresponding 2-vinyl- and 2-propargylpyrrolidines **4** and **5** (Scheme 4). *trans*-Ethyl cinnamate and an α,β -unsaturated diene provided β -amino acid esters **6** and **7** with good diastereoselectivity for the 2,3-*trans* pyrrolidines (Scheme 4).

Hexarotene methyl ester, the methyl ester of a retinoid anticancer agent that has recently shown promise in the treatment and prevention of Alzheimer's disease,³⁸ provided pyrrolidine **8** from the coupling reaction along with two alkyl Heck-type diastereomers **9** (Scheme 5).

Scheme 4. Reactions of Dienes and a Dienoate




Scheme 5. Reaction of Bexarotene Methyl Ester



These intermolecular coupling reactions appear to occur through copper-catalyzed/ MnO_2 mediated stepwise oxidative coupling sequence (Scheme 6). Copper(II)-catalyzed oxidation

Scheme 6. Proposed Reaction Mechanism




of the alkylborane to its corresponding alkyl radical initiates the process.^{33,36,39} The alkyl radical then adds to the styrene to produce a stabilized benzylic radical intermediate. This intermediate can combine with $[\text{Cu}(\text{II})]$ to form an alkylcopper(III) intermediate capable of undergoing C–N bond formation via reductive elimination (path I).⁴⁰ Alternatively, the benzylic radical could be further oxidized by MnO_2 to provide a benzylic carbocation that is then trapped by the pendant amine (path II).³¹ Oxidation of $[\text{Cu}(\text{I})]$ back to $[\text{Cu}(\text{II})]$ with MnO_2 then closes the catalytic cycle. At the onset of our mechanistic investigation, we could not differ-

entiate between path I and path II because, on the basis of oxidation potentials, MnO_2 is capable of oxidizing both $[\text{Cu}(\text{I})]$ and a benzylic radical, although $[\text{Cu}(\text{I})]$ is the more easily oxidized species.⁴¹

To investigate the mechanism further, a series of vinylcyclopropanes were submitted to the coupling reaction with *N*-Cbz- β -aminoethyltrifluoroborate 1a (Scheme 7). Reaction with

Scheme 7. Radical Clock Mechanism Probes



2-(buta-1,3-dienyl)cyclopropylbenzene provided a mixture of pyrrolidine diastereomers 10, and no cyclopropane ring-opened products were detected (Scheme 7). Reaction with 2-((phenylcyclopropyl)vinyl)benzene provided both pyrrolidine diastereomers 11 and dihydronaphthalene 12.³³ The regiochemistry of 12 indicates that cyclopropane ring opening occurred at the phenyl-bearing carbon. This supports a radical cyclopropane ring-opening mechanism over a metal-mediated ring opening where the less substituted organometallic would have been preferred.⁴² To differentiate between a carbocation and a radical cyclopropane opening, the 2-*tert*-butoxy-3-(1-phenylvinyl)cyclopropylbenzene radical clock was applied (Scheme 7). Newcomb has demonstrated this kind of radical clock will open at the oxygen-bearing carbon in carbocationic mechanisms and at the phenyl-bearing carbon in radical mechanisms.⁴³ In the event, a mixture of 4-phenylnaphthylene 13 and dihydronaphthalene 14 were obtained in this reaction. Naphthalene 13 is likely formed by elimination of *t*-BuOH from 14. The regioselectivity in these reactions support involvement of radical intermediates and do not provide evidence for carbocation intermediates. The lack of ring-opened product from reaction with the cyclopropyl diene probe could indicate that C–N bond formation is favored over ring-opening and/or radical addition to the arene when the carbon is less hindered. With increased steric hindrance at carbon, ring-opening and/or radical addition to the arene becomes competitive. While pyrrolidine formation without ring-opening is feasible, it is also possible the ring could open and subsequently close prior to pyrrolidine formation.⁴⁴

In summary, we have developed new conditions for the synthesis of simple 2-arylpyrrolidines from vinylarenes and dienes. The scope of the alkene partner is broad. Radical clock experiments support a purely radical mechanism, likely

involving C–N bond formation through a copper(III) intermediate. Our future efforts involve reaction refinement and scope expansion.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.6b01259](https://doi.org/10.1021/acs.orglett.6b01259).

Experimental procedures and characterization of new compounds (PDF)

NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: schemler@buffalo.edu.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the National Institute of General Medical Science for support of this work (R01 078383) and the Bureau of Educational and Cultural Affairs of the U.S. Department of States for a Fulbright fellowship to C.U. We thank Mr. Shuklendu Karyakarte (UB chemistry) for assistance with some NMR spectra.

■ REFERENCES

- (1) Klapars, A.; Campos, K. R.; Waldman, J. H.; Zewge, D.; Dormer, P. G.; Chen, C.-Y. *J. Org. Chem.* **2008**, *73*, 4986.
- (2) Cho-Schultz, S.; Patten, M. J.; Huang, B.; Elleraas, J.; Gajiwala, K. S.; Hickey, M. J.; Wang, J.; Mehta, P. P.; Kang, P.; Gehring, M. R.; Kung, P.-P.; Sutton, S. C. *J. Comb. Chem.* **2009**, *11*, 860.
- (3) Elliott, R. L.; Ryther, K. B.; Anderson, D. J.; Raszkievicz, J. L.; Campbell, J. E.; Sullivan, J. P.; Garvey, D. S. *Bioorg. Med. Chem. Lett.* **1995**, *5*, 991.
- (4) Wagner, F. F.; Comins, D. L. *Tetrahedron* **2007**, *63*, 8065.
- (5) Campos, K. R.; Klapars, A.; Waldman, J. H.; Dormer, P. G.; Chen, C. Y. *J. Am. Chem. Soc.* **2006**, *128*, 3538.
- (6) Wu, S.; Lee, S.; Beak, P. *J. Am. Chem. Soc.* **1996**, *118*, 715.
- (7) Brinner, K. M.; Ellman, J. A. *Org. Biomol. Chem.* **2005**, *3*, 2109.
- (8) Rajender Reddy, L.; Das, S. G.; Liu, Y.; Prashad, M. *J. Org. Chem.* **2010**, *75*, 2236.
- (9) Leemans, E.; Mangelinckx, S.; De Kimpe, N. *Chem. Commun.* **2010**, *46*, 3122.
- (10) Zuo, Z.; Ahneman, D. T.; Chu, L.; Terrett, J. A.; Doyle, A. G.; MacMillan, D. W. C. *Science* **2014**, *345*, 437.
- (11) Broere, D. L. J.; de Bruin, B.; Reek, J. N. H.; Lutz, M.; Dechert, S.; van der Vlugt, J. I. *J. Am. Chem. Soc.* **2014**, *136*, 11574.
- (12) Yus, M.; Soler, T.; Foubelo, F. *J. Org. Chem.* **2001**, *66*, 6207.
- (13) Coldham, I.; Robinson, S. P.; Baxter, C. A. *Synlett* **2012**, *23*, 2405.
- (14) Bunrit, A.; Dahlstrand, C.; Olsson, S. K.; Srifa, P.; Huang, G.; Orthaber, A.; Sjoberg, P. J. R.; Biswas, S.; Himo, F.; Samec, J. S. M. *J. Am. Chem. Soc.* **2015**, *137*, 4646.
- (15) Cui, Z.; Yu, H.-J.; Yang, R.-F.; Gao, W.-Y.; Feng, C.-G.; Lin, G.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 12394.
- (16) Gribkov, D. V.; Pastine, S. J.; Schnurch, M.; Sames, D. *J. Am. Chem. Soc.* **2007**, *129*, 11750.
- (17) Henry, C. E.; Xu, Q.; Fan, Y. C.; Martin, T. J.; Belding, L.; Dudding, T.; Kwon, O. *J. Am. Chem. Soc.* **2014**, *136*, 11890.
- (18) Scarborough, C. C.; Stahl, S. S. *Org. Lett.* **2006**, *8*, 3251.
- (19) Shi, M.; Liu, L.-P.; Tang, J. *Org. Lett.* **2006**, *8*, 4043.
- (20) Rao, W.; Chan, P. W. H. *Chem. - Eur. J.* **2008**, *14*, 10486.
- (21) Schlummer, B.; Hartwig, J. F. *Org. Lett.* **2002**, *4*, 1471.
- (22) Chen, J.; Zhou, L.; Yeung, Y.-Y. *Org. Biomol. Chem.* **2012**, *10*, 3808.
- (23) Xu, T.; Qiu, S.; Liu, G. *J. Organomet. Chem.* **2011**, *696*, 46.
- (24) O'Broin, C. Q.; Fernandez, P.; Martinez, C.; Muniz, K. *Org. Lett.* **2016**, *18*, 436.
- (25) Hennessy, E. T.; Betley, T. A. *Science* **2013**, *340*, 591.
- (26) Jui, N. T.; Garber, J. A. O.; Finelli, F. G.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2012**, *134*, 11400.
- (27) Gesmundo, N. J.; Grandjean, J.-M. M.; Nicewicz, D. A. *Org. Lett.* **2015**, *17*, 1316.
- (28) Grandjean, J.-M. M.; Nicewicz, D. A. *Angew. Chem., Int. Ed.* **2013**, *52*, 3967.
- (29) Michaelis, D. J.; Shaffer, C. J.; Yoon, T. P. *J. Am. Chem. Soc.* **2007**, *129*, 1866.
- (30) Zhao, B.; Peng, X.; Zhu, Y.; Ramirez, T. A.; Cornwall, R. G.; Shi, Y. *J. Am. Chem. Soc.* **2011**, *133*, 20890.
- (31) Lu, D.-F.; Zhu, C.-L.; Jia, Z.-X.; Xu, H. *J. Am. Chem. Soc.* **2014**, *136*, 13186.
- (32) Ghosh, A. K.; Brindisi, J. *Med. Chem.* **2015**, *58*, 2895.
- (33) Liwosz, T. W.; Chemler, S. R. *Org. Lett.* **2013**, *15*, 3034.
- (34) Kamatani, A.; Overman, L. E. *J. Org. Chem.* **1999**, *64*, 8743.
- (35) Molander, G. A.; Jean-Gerard, L. *J. Org. Chem.* **2007**, *72*, 8422.
- (36) Sorin, G.; Mallorquin, R. M.; Contie, Y.; Baralle, A.; Malacria, M.; Goddard, J. P.; Fensterbank, L. *Angew. Chem., Int. Ed.* **2010**, *49*, 8721.
- (37) Attempted couplings with other aminoalkyl trifluoroborates were not successful (see the Supporting Information).
- (38) (a) Qu, L.; Tang, X. *Cancer Chemother. Pharmacol.* **2010**, *65*, 201. (b) Habchi, J.; Arosio, P.; Perni, M.; Costa, A. R.; Yagi-Utsumi, M.; Joshi, P.; Chia, S.; Cohen, S. I. A.; Muller, M. B. D.; Linse, S.; Nollen, E. A. A.; Dobson, C. M.; Knowles, T. P. J.; Vendruscolo, M. *Sci. Adv.* **2016**, *2*, e1501244.
- (39) (a) Tang, S.; Liu, C.; Lei, A. *Chem. Soc. Rev.* **2015**, *44*, 1070. (b) Zhao, W.; Montgomery, J. *Angew. Chem., Int. Ed.* **2015**, *54*, 12683.
- (40) Clark, S. J.; Roche, C. *Chem. Commun.* **2005**, 5175.
- (41) (a) Milazzo, G.; Caroli, S.; Sharma, V. K. *Tables of Standard Electrode Potentials*; Wiley: Chichester, 1978. (b) Bard, A. J.; Parsons, R.; Jordan, J. *Standard Potentials in Aqueous Solutions*; Marcel Dekker: New York, 1985. (c) Connelly, N. G.; Geiger, W. E. *Chem. Rev.* **1996**, *96*, 877. (d) Wayner, D. D. M.; McPhee, D. J.; Griller, D. *J. Am. Chem. Soc.* **1988**, *110*, 132.
- (42) Wender, P. A.; Dyckman, A. J.; Husfeld, C. O.; Kadereit, D.; Love, J. A.; Rieck, H. *J. Am. Chem. Soc.* **1999**, *121*, 10442.
- (43) (a) Newcomb, M.; Chestney, D. L. *J. Am. Chem. Soc.* **1994**, *116*, 9753. (b) Le Tadic-Biadatti, M.-H.; Newcomb, M. *J. Chem. Soc., Perkin Trans. 2* **1996**, 1467. (c) Faulkner, A.; Race, N. J.; Scott, J. S.; Bower, J. F. *Chem. Sci.* **2014**, *5*, 2416.
- (44) Benkovics, T.; Du, J.; Guzei, I. A.; Yoon, T. P. *J. Org. Chem.* **2009**, *74*, 5545.