

Communication

# Decarboxylative Fluorination of Aliphatic Carboxylic Acids via Photoredox Catalysis

Sandrine Ventre, Filip R. Petronijevi#, and David W. C. MacMillan

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • DOI: 10.1021/jacs.5b02244 • Publication Date (Web): 16 Apr 2015

Downloaded from <http://pubs.acs.org> on April 19, 2015

## Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



**ACS Publications**  
High quality. High impact.

# Decarboxylative Fluorination of Aliphatic Carboxylic Acids via Photoredox Catalysis

Sandrine Ventre, Filip R. Petronijevic and David W. C. MacMillan\*

Merck Center for Catalysis at Princeton University, Princeton, New Jersey 08544  
 dmacmill@princeton.edu


RECEIVED DATE

**Abstract.** The direct conversion of aliphatic carboxylic acids to the corresponding alkyl fluorides has been achieved via visible light-promoted photoredox catalysis. This operationally simple, redox-neutral fluorination method is amenable to a wide variety of carboxylic acids. Photon-induced oxidation of carboxylates leads to the formation of carboxyl radicals, which upon rapid CO<sub>2</sub>-extrusion and F• transfer from a fluorinating reagent yield the desired fluoroalkanes with high efficiency. Experimental evidence indicates that an oxidative quenching pathway is operable in this broadly applicable fluorination protocol.


The capacity of fluorine atoms to engender a variety of useful properties in pharmaceuticals, agrochemicals, and performance materials has driven significant research efforts towards the invention of novel fluorination reactions.<sup>1-3</sup> Over the last two decades, significant progress has been made towards the production of sp<sup>2</sup> C–F bonds;<sup>4</sup> however, catalytic methods for sp<sup>3</sup> C–F formation have only recently become available.<sup>5-8</sup> In particular, metal-mediated radical C–H abstraction-fluorination protocols have been developed to form tertiary aliphatic,<sup>6</sup> benzylic,<sup>7</sup> as well as allylic<sup>8</sup> C–F centers, a strategy that is founded upon the selective functionalization of weak C–H bonds. Despite these important advances, the development of a general sp<sup>3</sup> C–F bond-forming platform that is (i) highly regioselective, (ii) bond strength independent, (iii) operationally simple, and (iv) able to employ readily available, inexpensive starting materials, remains a challenging goal.

Harnessing visible light as a safe, renewable, and inexpensive source of chemical energy to facilitate the construction of complex organic molecules has emerged recently as a powerful theme in organic chemistry.<sup>9</sup> In this context, our group has recently introduced a visible light-mediated alkylation of α,β-unsaturated carbonyl compounds with alkyl radicals, generated via a CO<sub>2</sub>-extrusion mechanism involving carboxylic acids (Eq 1).<sup>10</sup> Based on these findings, we wondered if a similar photon-induced decarboxylation strategy might be employed as a general platform for the construction of C–F bonds. Precedent for this transformation has been demonstrated via the silver-mediated Hunsdiecker reaction<sup>11</sup> and the work of Sammis to achieve a light-promoted decarboxylative fluorination of 2-aryloxyacetic acids to generate α-oxyfluoro motifs.<sup>12</sup> Despite these seminal studies, a general light-mediated strategy for the fluorination of a wide range of aliphatic carboxylic acids has not yet been reported. In this communication, we further advance the visible light activation concept to describe the first broadly applicable protocol for the photon-mediated decarboxylative fluorination of sp<sup>3</sup> carbon bearing carboxylic acids, using a blue LED light source and a commercial photocatalyst (Eq 2).


**Design Plan.** Drawing from the mechanistic insights gained in the course of our decarboxylative alkylation<sup>10</sup> and arylation<sup>13</sup> methods, we envisioned that a broad range of aliphatic carboxylic acids could be employed as viable precursors to fluoroalkanes. The specific mechanistic details of our proposed visible light-mediated photoredox decarboxylative fluorination are outlined in Scheme 1. Irradiation of heteroleptic iridium(III) photocatalyst Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub> (**1**) with visible light



## Decarboxylative Conjugate Addition Using Aliphatic Carboxylic Acids (Eq 1)




## Fluorination Enabled by Decarboxylative Radical Coupling (Eq 2)



leads to the formation of a long-lived ( $\tau = 2.3 \mu\text{s}$ )<sup>14</sup> excited state  $^*\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})^+$  (**2**), which can undergo oxidative quenching ( $E_{1/2}^{\text{red}} = -0.89 \text{ V vs. SCE in CH}_3\text{CN}$ )<sup>14</sup> in the presence of an appropriate electron acceptor. We hypothesized that an initial reduction of a sacrificial quantity of Selectfluor® (**3**) ( $E_{1/2}^{\text{red}} = +0.33 \text{ V vs. SCE in CH}_3\text{CN}$ )<sup>15</sup> by  $^*\text{Ir}(\text{III})$  **2** via a single electron transfer (SET) process should generate the strongly oxidizing  $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})^{2+}$  (**5**). Indeed, the earlier work of Sammis clearly delineated that such a possibility was viable with an electrophilic source of fluorine.<sup>12</sup> We further assumed that base-mediated formation of an alkyl carboxylate followed by an SET oxidation ( $E_{1/2}^{\text{red}} = +1.16 \text{ V for hexanoate}$ )<sup>16</sup> using the transiently formed Ir(IV) species **5** ( $E_{1/2}^{\text{red}} = +1.69 \text{ V vs. SCE in CH}_3\text{CN}$ )<sup>14</sup> would be thermodynamically feasible. This process is envisioned to generate a carboxyl radical, which upon immediate extrusion of CO<sub>2</sub> should provide the SOMO species **7**.<sup>17</sup> Concurrently, reduction of Ir(IV) **5** would regenerate the ground-state photocatalyst **1**, thus completing the photoredox cycle. At this stage, direct F-transfer from Selectfluor® to the alkyl radical **7** is proposed to forge the desired fluoroalkane bond (**8**) with concomitant formation of the corresponding Selectfluor® radical cation **4**. We assume that radical cation **4** would replace Selectfluor® in subsequent photoredox cycles as a suitable electron acceptor in the conversion of excited state  $^*\text{Ir}(\text{III})$  (**2**) to the requisite  $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})^{2+}$  (**5**) species.

**Results.** We first explored the proposed decarboxylative fluorination reaction in the context of *N*-benzoyl-4-piperidine carboxylic acid and Selectfluor® (Table 1). Examination of a range of photocatalysts and bases revealed that a combination of  $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$  (**1**) and disodium hydrogen phosphate (1 equiv.) was superior with respect to reaction

**Scheme 1.** Mechanism for Decarboxylative Fluorination.

efficiency. The use of more oxidizing photocatalysts, such as  $\text{Ru}(\text{phen})_3^{2+}$  or  $\text{Ru}(\text{bpz})_3^{2+}$  (9), resulted in slower or no reaction (entries 1, 3, and 4). Moreover, the use of the Sammis protocol ( $\text{Ru}(\text{bpy})_3^{2+}$  (10), NaOH, developed for the decarboxylative fluorination of  $\alpha$ -oxy acids)<sup>12c</sup> gave no observable product. Further improvement was achieved via the use of two equivalents of  $\text{Na}_2\text{HPO}_4$  as a base, presumably due to the larger extent of acid deprotonation (entries 5 and 6; 80% vs. 90% yield). The critical role of acetonitrile and water as solvent mixture was demonstrated by the absence of any fluorinated product when either of these two solvents was used independently (entries 7 and 8). This observation is likely due to the low solubility of (i) Selectfluor® in acetonitrile and (ii) alkyl carboxylic acids in water, whereas the miscible mixture of acetonitrile and water allowed for both substrates to be employed in a homogeneous solution. Lastly, control experiments confirmed the requirement of a photocatalyst, base, and a light source in this new fluorination protocol (entries 9–11).

**Reaction Scope.** Having identified optimal conditions for what we hoped would be a general photocatalytic  $\text{CO}_2$ -extrusion fluorination protocol, we aimed to define the scope of the carboxylic acid precursor. As shown in Table 2, a wide range of differentially substituted alkyl carboxylates were readily converted to the corresponding alkyl fluorides. It is of note that primary, secondary, and tertiary alkyl carboxylic acids are all well-tolerated, with no observed decrease in efficiency with less substituted acids. Substrates with a heteroatom in the vicinity of the carboxylic group ( $\alpha$  or  $\beta$ ) underwent faster  $\text{CO}_2$ -extrusion/fluorination (precursors to **18** (99% yield) and **28–30**, (92%, 90% and 90% yields respectively)), with reaction times in the 1–3 h range. In addition, 2 equiv. of Selectfluor® could be used without any decrease in reaction efficiency with these substrates. The same observation was made for benzylic and homobenzylic carboxylic acids (precursors to **14** (87% yield), **17** (82% yield), and **22** (92% yield), presumably due to stabilization of the transiently formed radical intermediate. It is important to note that unactivated acids were also found to be competent substrates for this fluorination protocol (products **20** (70% yield), **27** (83% yield), and **34** (79% yield)). However, when 1,4-phenyldipropionic acid was employed as a substrate, no formation of difluoride **11** was observed, due to the very low solubility of the dicarboxylic acid in the acetonitrile/water mixture. To our delight, when the corresponding preformed disodium salt was employed, the desired difluoride was isolated in 71% yield. Similarly, 4-*tert*-butylcyclohexanecarboxylic acid

**Table 1.** Initial Studies and Reaction Optimization.

carboxylic acid

fluoroalkane

Selectfluor®

entry	photocatalyst	base	solvent <sup>a</sup>	yield <sup>b</sup>
1	$\text{Ru}(\text{bpz})_3(\text{PF}_6)_2$	$\text{Na}_2\text{HPO}_4$ (1 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	64%
2	$\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$	$\text{CsF}$ (1 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	68%
3	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	$\text{NaOH}$ (1 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	0%
4	$\text{Ru}(\text{phen})_3(\text{PF}_6)_2$	$\text{Na}_2\text{HPO}_4$ (1 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	0%
5	$\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$	$\text{Na}_2\text{HPO}_4$ (1 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	80%
6	$\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$	$\text{Na}_2\text{HPO}_4$ (2 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	90%
7	$\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$	$\text{Na}_2\text{HPO}_4$ (2 equiv.)	$\text{CH}_3\text{CN}$	0%
8	$\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$	$\text{Na}_2\text{HPO}_4$ (2 equiv.)	$\text{H}_2\text{O}$	0%
9	none	$\text{Na}_2\text{HPO}_4$ (2 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	0%
10	$\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$	none	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	0%
11 <sup>c</sup>	$\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbbpy})\text{PF}_6$	$\text{Na}_2\text{HPO}_4$ (2 equiv.)	$\text{CH}_3\text{CN}/\text{H}_2\text{O}$	0%

<sup>a</sup>Acetonitrile/water 1:1 (v/v) was used as solvent. <sup>b</sup>Yields determined by  $^1\text{H}$  NMR using 3,5-bis(trifluoromethyl)bromobenzene as an internal standard. <sup>c</sup>Reaction performed in the absence of light.

provided higher yield of the corresponding fluorocyclohexane **20** when the ratio of the acetonitrile/water medium was adjusted to 3:1. It is interesting to note that while the fluoride **28**, derived from ribosic acid, was formed in high yield, the corresponding glucopyranouronic acid derivative did not react under these reaction conditions. We believe that this result can be rationalized by the change in bond strength and oxidation potential of the  $\text{C}-\text{CO}_2$  moiety as it exists in either the axial anomer position (with ribosic acid) or the anomeric equatorial topography (as expected with glucopyranouronic acid). We assume that the axial  $\text{C}-\text{CO}_2$  bond is weaker, and the rate of decarboxylation is faster (in competition with back electron transfer) due to the hyperconjugative stabilization by the ring oxygen lone pair in the case of the ribosic carboxylic acid system. In contrast, we presume that back electron transfer is competitive with decarboxylation in the case of glucopyranouronic acid, thereby preventing the formation of an  $\alpha$ -oxy radical intermediate. Interestingly, no elimination of the fluoride group was observed when this decarboxylative fluorination protocol was used to generate a  $\beta$ -fluoro carbonyl under basic conditions. More specifically, when a 1,4-keto acid substrate was employed, the desired  $\beta$ -fluoroketone **25** was isolated in 96% yield, without observation of the corresponding  $\alpha,\beta$ -unsaturated product. Finally, less reactive substrates, such as unactivated primary (precursors to products **12** and **15**) or tertiary carboxylic acids (precursors to **32** and **34**) proved to be viable substrates; however, longer reaction times (12–15 h) were required for full conversion of the starting material. This observation is in agreement with the rate of oxidation of the carboxylate (in the case of primary acids) or the rate of formation of the carboxylate (in the case of cyclic tertiary acids),

**Table 2.** Decarboxylative Fluorination: Scope of Carboxylic Acids.<sup>a</sup>

RCO <sub>2</sub> H	product	RCO <sub>2</sub> H	product	RCO <sub>2</sub> H	product
	 <b>11</b> 77% yield <sup>e</sup>		 <b>12</b> 71% yield		 <b>13</b> 81% yield
	 <b>14</b> 87% yield <sup>b,g</sup>		 <b>15</b> 79% yield <sup>d</sup>		 <b>16</b> 92% yield
	 <b>17</b> 82% yield <sup>b</sup>		 <b>18</b> 99% yield <sup>b,g</sup>		 <b>19</b> 82% yield <sup>e,i</sup>
	 <b>20</b> 70% yield <sup>e,h,j</sup>		 (±)- <b>21</b> 80% yield <sup>b</sup>		 <b>22</b> 92% yield <sup>b,f,g</sup>
	 (±)- <b>23</b> 71% yield <sup>e,f</sup>		 (±)- <b>24</b> 85% yield <sup>b,g</sup>		 (±)- <b>25</b> 96% yield <sup>b,f,g</sup>
	 <b>26</b> 90% yield		 (±)- <b>27</b> 83% yield		 <b>28</b> 92% yield <sup>b,g,i</sup>
	 (±)- <b>29</b> 90% yield <sup>e,f,g</sup>		 (±)- <b>30</b> 90% yield <sup>b,g</sup>		 <b>31</b> 90% yield <sup>b,g</sup>
	 <b>32</b> 76% yield <sup>e</sup>		 <b>33</b> 88% yield <sup>b,g</sup>		 <b>34</b> 79% yield <sup>e</sup>

<sup>a</sup>Isolated yields, see Supporting Information for experimental details. <sup>b</sup>Reaction time 1 h. <sup>c</sup>Reaction time 3 h. <sup>d</sup>Reaction time 12 h. <sup>e</sup>Reaction time 15 h. <sup>f</sup>Reaction run with Ru(bpz)<sub>3</sub>(PF<sub>6</sub>)<sub>2</sub> instead of Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub>. <sup>g</sup>Reaction run using 2 equiv. of Selectfluor. <sup>h</sup>Reaction run in acetonitrile/water 3:1. <sup>i</sup>Obtained as a single diastereomer. <sup>j</sup>Obtained as 2.5:1 *trans/cis* mixture of diastereomers (major diastereomer shown).


critical steps ahead of the formation of the radical intermediate. Finally, we have performed a series of Stern-Volmer fluorescence quenching studies in an effort to gather evidence with regard to the mechanistic proposal outlined in Scheme 1.<sup>19</sup> As revealed in Figure 1, we observed that the emission intensity of the excited state of the heteroleptic catalyst Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub> is diminished in the presence of Selectfluor® (Figure 1). In contrast, fluorescence quenching was not observed when a solution of sodium 4-*tert*-butylcyclohexanecarboxylate was exposed to the photoexcited \*Ir(III) species. These experiments strongly indicate that the

reduction of Selectfluor® is likely the initiation point of the photoredox catalytic cycle, as proposed in Scheme 1. Moreover, we presume that the requisite oxidation of the aliphatic carboxylate is performed by the resulting Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)<sup>2+</sup> species (also delineated in Scheme 1).

In summary, we have developed a photoredox-assisted decarboxylative fluorination protocol and demonstrated its utility over a wide range of carboxylic acid substrates. In contrast to previously described methods, this redox-neutral reaction does not require activated substrates. In addition, its operational simplicity and mild reaction conditions allow for the synthesis of



**Figure 1.** Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub> Emission Quenching with Selectfluor® and Sodium 4-*tert*-butylcyclohexanecarboxylate.



a diverse collection of valuable fluorinated products. Notably, under these reaction conditions more activated substrates require lower amounts of the electrophilic fluorinating reagent and shorter reaction times. Mechanistic studies have provided evidence supporting an oxidative quenching pathway, in which reduction of the N–F bond of Selectfluor® initiates the photoredox cycle prior to the carboxylic acid oxidation/decarboxylation sequence.

**Acknowledgement.** Financial support was provided by the NIGMS (R01 GM093213-05) and kind gifts from Merck, Amgen, and AbbVie.

**Supporting Information Available.** Experimental procedures and spectral data are provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881.
- (2) (a) Furuya, T.; Kuttruff, C. A.; Ritter, T. *Curr. Opin. Drug Disc.* **2008**, *11*, 803. (b) Neumann, C. N.; Ritter, T. *Angew. Chem. Int. Ed.* **2015**, DOI:10.1002/anie.201410288.
- (3) (a) Hagmann, W. K. *J. Med. Chem.* **2008**, *51*, 4359. (b) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320.
- (4) (a) Furuya, T.; Kamlet, A. S.; Ritter, T. *Nature* **2011**, *473*, 470. (b) Hull, K. L.; Anani, W. Q.; Sanford, M. S. *J. Am. Chem. Soc.* **2006**, *128*, 7134. (c) Chen, K. S. L.; Wasa, M.; Wang, X.; Yu, J.-Q. *Angew. Chem. Int. Ed.* **2011**, *50*, 9081. (d) Troung, T.; Klimovica, K.; Daugulis, O. *J. Am. Chem. Soc.* **2013**, *135*, 9342. (e) Lou, S.-J.; Xu, D.-Q.; Xia, A.-B.; Wang, Y.-F.; Liu, Y.-K.; Du, X.-H.; Xu, Z.-Y. *Chem. Commun.* **2013**, *49*, 6218. (f) Watson, D. A.; Su, M.; Teverovskiy, G.; Zhang, Y.; Garcia-Fortanet, J.; Kinzel, T.; Buchwald, S. L. *Science* **2009**, *325*, 166. (g) Lee, H. G.; Milner, P. J.; Buchwald, S. L. *J. Am. Chem. Soc.* **2014**, *136*, 3792.
- (5) For selected examples see: (a) Beeson, T. D.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2005**, *127*, 8826. (b) Kalow, J. A.; Doyle, A. G. *J. Am. Chem. Soc.* **2010**, *132*, 3268. (c) Katcher, M. H.; Sha, A.; Doyle, A. G. *J. Am. Chem. Soc.* **2011**, *133*, 15902. (d) Hollingworth, C.; Hazari, A.; Hopkinson, M. N.; Tredwell, M.; Benedetto, E.; Huiban, M.; Gee, A. D.; Brown, J. M.; Gouverneur, V. *Angew. Chem. Int. Ed.* **2011**, *50*, 2613. (e) Topczewski, J. J.; Tewson, T. J.; Nguyen, H. M. *J. Am. Chem. Soc.* **2011**, *133*, 19318. (f) Barker, T. J.; Boger, D. L. *J. Am. Chem. Soc.* **2012**, *134*, 13588. (g) Sladojevich, F.; Arlow, S. I.; Tang, P.; Ritter, T. *J. Am. Chem. Soc.* **2013**, *135*, 2470.
- (6) (a) Liu, W.; Huang, X.; Cheng, M.-J.; Nielsen, R. J.; Goddard, W. A., III; Groves, J. T. *Science* **2012**, *337*, 1322. (b) Bloom, S.; Knippel, J. L.; Lectka, T. *Chem. Sci.* **2014**, *5*, 1175.
- (7) (a) Liu, W.; Groves, J. T. *Angew. Chem. Int. Ed.* **2013**, *52*, 6024. (b) Bloom, S.; Pitts, C. R.; Woltermist, R.; Griswold, A.; Holl, M. G.; Lectka, T. *Org. Lett.* **2013**, *15*, 1722. (c) Bloom, S.; Sharber, S. A.; Holl, M. G.; Knippel, J. L.; Lectka, T. *J. Org. Chem.* **2013**, *78*, 11082. (d) Amaoka, Y.; Nagatomo, M.; Inoue, M. *Org. Lett.* **2013**, *15*, 2160.
- (8) (a) Bloom, S.; Pitts, C. R.; Miller, D. C.; Haselton, N.; Holl, M. G.; Urheim, E.; Lectka, T. *Angew. Chem. Int. Ed.* **2012**, *51*, 10580. (b) Braun, M.-G.; Doyle, A. G. *J. Am. Chem. Soc.* **2013**, *135*, 12990.
- (9) (a) Nicewicz, D.; MacMillan, D. W. C. *Science* **2008**, *32*, 77. (b) Schultz, D. M.; Yoon, T. P. *Science* **2014**, *343*, 6174. (c) Yoon, T. P.; Ischay, M. A.; Du, J. *Nat. Chem.* **2010**, *2*, 527. (d) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. *Chem. Rev.* **2013**, *113*, 5322. (e) Narayanan, J. M. R.; Stephenson, C. R. J. *Chem. Soc. Rev.* **2011**, *40*, 102.

- (10) Chu, L.; Ohta, C.; Zuo, Z.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 10886.
- (11) Yin, F.; Wang, Z.; Li, Z.; Li, C. *J. Am. Chem. Soc.* **2012**, *134*, 10401.
- (12) (a) Rueda-Becerril, M.; Chatlova-Sazepin, C.; Leung, J. C. T.; Okbinoglu, T.; Kennepohl, P.; Paquin, J.-F.; Sammis, G. M. *J. Am. Chem. Soc.* **2012**, *134*, 4026. (b) Leung, J. C. T.; Chatlova-Sazepin, C.; West, J.; Rueda-Becerril, M.; Paquin, J.-F.; Sammis, G. M. *Angew. Chem. Int. Ed.* **2012**, *51*, 10804. (c) Rueda-Becerril, M.; Mahe, O.; Drouin, M.; Majewski, M. B.; West, J. G.; Wolf, M. O.; Sammis, G. M.; Paquin, J.-F. *J. Am. Chem. Soc.* **2014**, *136*, 2637.
- (13) Zuo, Z.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 5257.
- (14) Lowry, M. S.; Goldsmith, J. L.; Slinker, J. D.; Rohl, R.; Pascal, R. A.; Malliaras, G. G.; Bernhard, S. *Chem. Mater.* **2005**, *17*, 5712.
- (15) (a) Stavber, S.; Zupan, M. *Acta Chim. Slov.* **2005**, *52*, 13. (b) Girina, G. P.; Fainzil'berg, A. A.; Feoktistov, L. G. *Russ. J. Electrochem.* **2000**, *36*, 162.
- (16) Galicia, M.; Gonzalez, F. J. *J. Electrochem. Soc.* **2002**, *149*, D46.
- (17) Bockman, T. M.; Hubig, S. M.; Kochi, J. K. *J. Org. Chem.* **1997**, *62*, 2210.
- (18) Fluorides **22**, **25** and **29** were also obtained using Ir[dF(CF<sub>3</sub>)ppy]<sub>2</sub>(dtbbpy)PF<sub>6</sub> photocatalyst in 71%, 89% and 85% yield, respectively. However, we observed slightly increased efficiency with the ruthenium-based photocatalyst Ru(bpz)<sub>3</sub>(PF<sub>6</sub>)<sub>2</sub> in these three cases.
- (19) See Supporting Information for further details about fluorescence quenching studies.

Graphical Abstract

