Copper-Catalyzed Direct Alkenylation of N-Iminopyridinium Ylides**

James J. Mousseau, James A. Bull, and André B. Charette*

In memory of Keith Fagnou

The pyridine moiety is a privileged structure that is ubiquitous in nature and is often a key component of pharmaceutically active compounds.^[1] As such, there has been much interest in the synthesis of substituted pyridine derivatives. 2-Alkenyl pyridine derivatives are of biological interest (Scheme 1) and are important metal ligands. Furthermore, alkenyl pyridine derivatives can be valuable precursors to alkyl pyridines and piperidines, themselves of significant biological importance.^[2]



Scheme 1. Biologically important 2-alkenyl or 2-alkyl pyridine derivatives.

Direct arylations have become increasingly important and greener alternatives to traditional cross-coupling reactions.^[3] Recent work has shown that such reactions are applicable to activated pyridinium species.^[4] In contrast, comparatively little has been published on direct alkenylation reactions; in particular, there are few examples of direct vinylations on activated pyridines. In these cases, symmetrical alkynes with little functionality or relatively simple Heck acceptors were required as the coupling partners.^[5] Cognizant of these deficiencies, we believed that *N*-iminopyridinium ylides could provide access to functionalized 2-alkenyl pyridines by a direct alkenylation reaction with functionalized vinyl iodides.

Direct reactions often employ expensive metal catalysts (Pd or Rh). Less costly Fe and Cu catalytic systems are

[*]	J. J. Mousseau, Dr. J. A. Bull, Prof. A. B. Charette
	Department of Chemistry, Université de Montréal
	P.O. Box 6128 Stn Downtown, Montréal, Québec, H3C 3J7 (Canada)
	Fax: (+1) 514-343-5900
	E-mail: andre.charette@umontreal.ca
[***1	This work was supported by the Natural Science and Engineering

seldom reported. Indeed, although electron-deficient arenes have been reported, applications of these inexpensive metals to the direct functionalization of heterocycles more often employ electron-rich substrates.^[6] Herein, we report the first Cu-catalyzed direct alkenylation of *N*-iminopyridinium ylides with inexpensive Cu salts.

Given our success with the direct arylation of N-iminopyridinium ylides, we first applied our optimal Pd-catalyzed arylation conditions to alkenylation with (E)- β -styryl iodide (2a).^[7] These conditions gave the vinylated pyridinium 3a in 36% yield calculated by ¹H NMR spectroscopy (Table S1, entry 1; see the Supporting Information) and initial optimization did not lead to a significant increase in yield. It was reasoned that the addition of a Lewis acid could increase the reactivity of the pyridinium by coordination to the N-benzoyl moiety. Indeed, the addition of CuBr to the Pd-catalyzed reaction increased the yield of **3a** to 63% (Table S1, entry 2). In light of this result, we postulated that Cu alone might promote the transformation. Gratifyingly, CuBr displayed superior reactivity to Pd(OAc)₂ (Table S1, entry 3). Decreasing the ligand loading gave a higher yield (Table S1, entries 4 and 5), and the reaction was successful in the absence of an external ligand (Table S1, entry 6), presumably because the Lewis basic N-benzoyl moiety acts as an intramolecular stabilizing group.

Encouraged by these results, we optimized the reaction without any external ligand. The alkenylation was insensitive to the source of Cu; most Cu^{II} and Cu^I salts were compatible, as well as Cu⁰ dust, and even a penny. In fact, the reaction could be performed in a Cu vessel without any additional metal source to provide a 74% yield calculated by ¹H NMR spectroscopy. CuBr₂ was chosen because of its low cost and high stability (Table S1, entry 7). The catalyst loading could be decreased to 2.5 mol% and still provide good yields, although 10 mol% were chosen for improved results. Aromatic and ethereal solvents both gave excellent results. Chlorobenzene was chosen because of its lower volatility. The yield of **3a** increased to 85% when 1.5 equiv of the ylide **1** was used, because of statistical suppression of the formation of the 2,6-divinylated by-product (Table S1, entries 9 and 10). Owing to the low cost and high efficiency of K₂CO₃ in the reaction, we elected to continue with this base, and 2 equiv were sufficient to enable the reaction [Eq. (1); Bz = benzoyl].

© 2010 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



^[**] This work was supported by the Natural Science and Engineering Research Council of Canada (NSERC), Merck Frosst Canada Ltd., Boehringer Ingelheim (Canada), Ltd., the Canada Research Chairs Program, the Canadian Foundation for Innovation, and the Université de Montréal. J.J.M. is grateful to FQRNT for a postgraduate scholarship.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.200906020.

Communications

The scope of the Cu-catalyzed reaction was found to be quite general (Table 1).^[8] Unsubstituted arenes present on the alkene reacted well (Table 1, entries 1 and 2). The excess ylide employed in the reaction could be recovered to give 93 % yield based on recovered starting material (Table 1, entry 1). The alkenylation tolerated substitution on the phenyl functionality with very good yield (Table 1, entry 3) and also on the alkene, where the group is less removed from the reactive site (Table 1, entry 4). Bis(vinyl iodide) **2e** also reacted to give

the dipyridinium adduct in synthetically useful yields (Table 1, entry 5). The reaction was successful with both electron-rich (Table 1, entries 6–9) and electron-poor substrates (Table 1, entries 10–12). Perhaps the most striking feature of the reaction is its chemoselectivity. Halogen substituents (Cl, Br, I; Table 1, entries 13–16) were tolerated on the phenyl ring with the reaction occurring selectively on the alkenyl iodide, and no arylated product was detected. These halogenated compounds (3m-3q) are of particular

Entry	Alkenyl Iodide	Product	Yield [%] ^[b]	Entry	Alkenyl Iodide	Product	Yield [%] ^[b]
1	'~ 2a	NBZ 3a	81 (93) ^[c]	12	21		83
2	'~ 2b		81	13	2m	NBz ci	74
3	2c	NBZ 3C	78	14	Br 2n		65
4	2d		48	15	20		74
5 ^[d]	2e		63	16	2p		47
6	2f	N N NBZ OMe	87	17	2e		41
7	Bno 2g		71	18	Ph 2q	N N N Bz 3r	53
8	MeO 2h	NBz MeO	93	19	2r	NBZ 3S	52
9	2i	NBz O 3i	75	20	ормв 2s	NBZ OPMB 3t	30
10	2j	NBZ 3i	71	21	2t	NBZ 3a	71
11			89	22	CO ₂ Et		41
		JK				30	

Table 1: Scope of the alkenyl iodide in the Cu-catalyzed reaction.^[a]

[a] Reaction conditions: 1 (1.5 equiv), 2 (1.0 equiv), CuBr₂ (10 mol%), K_2CO_3 (2 equiv), PhCl (0.2 m), 125 °C, 16–24 h. Bn = benzyl, PMB = *p*-methoxybenzyl. [b] Yield of isolated product. [c] Yield based on recovered starting material. [d] 3.1 equiv of ylide 1.

interest, as they contain reactive handles and, as such, could be used as a foundation for the synthesis of more-complex molecules. Substituents on the alkene with sp³ hybridization gave the corresponding alkenylated pyridinium species in moderate yields (Table 1, entries 18 and 19). Reactants with long alkyl chains displayed poorer reactivity, as iodide **2t** gave compound **3t** in 30% yield (Table 1, entry 20). However, these molecules are still of interest and may be used as scaffolds to build compounds such as CGS23113 (Scheme 1).^[9] Z-Alkenes are reactive, yielding products with *trans* configuration, possibly as a result of thermal isomerization to the more stable alkene (Table 1, entries 21 and 22).

The scope of the pyridinium ylide was considered next (Table 2). Pyridinium species with substituents at the 2-, 3-, and 4-positions underwent the alkenylation in good yields

Table 2: Scope of the pyridinium ylide in the Cu-catalyzed reaction.^[a]



[a] Reaction conditions: 1 (1.5 equiv), 2 (1.0 equiv), $CuBr_2$ (10 mol%), K_2CO_3 (2 equiv), PhCl (0.2 M), 125 °C, 16–24 h. [b] Yield of isolated product. [c] CuBr used in place of CuBr₂.

(Table 2, entries 1–3). In the case of the 3-substituted **1b**, vinylation occurred exclusively at the 6-position. This methodology could also be applied to other *N*-imino ylides; quinolinium species **1d** (Table 2, entry 4) afforded the vinylated product in moderate yield. It is important to note that the N–N bond could be cleaved in the presence of the alkene in very good yields (77–81%) to provide the corresponding 2-alkenylated pyridines (Scheme 2).

Next, we wished to gain insight into the mechanistic pathway of the reaction. Kinetic isotope effect (KIE) studies yielded a KIE value of 1.45, indicating that the C-H bond



Scheme 2. Cleavage of the N-N bond.

Angew. Chem. Int. Ed. 2010, 49, 1115–1118

breaking event is not rate-limiting. Furthermore D exchange studies (Table 3) clearly demonstrate a directed D incorporation to the 2,6-positions of the pyridinium ring in the presence of Cu (Table 3, entries 2 and 4)^[10] compared with





[a] Incorporation determined by ${}^{1}H$ NMR spectroscopy. [b] Equal D insertion in 2,4,6-positions.

indiscriminate D incorporation when no CuBr_2 was employed (Table 3, entry 3). This result, coupled with the fact that the direct alkenylation proceeds equally well with a range of Cu salts in different oxidation states, led us to propose the following catalytic cycle (Scheme 3): We believe that Cu^{II} is



Scheme 3. Proposed catalytic cycle.

reduced to Cu^{I} by the pyridinium ylide (**A**). In cases where Cu^{0} is operative, the ylide may add into the copper to generate a Cu^{II} intermediate that yields the reactive Cu^{I} species through either reduction or disproportionation.^[11] Next, there is a carbonate–bromide exchange to generate $CuCO_{3}$, which is likely to be the reactive species in all cases, regardless of the Cu source (**B**).^[11] This species can undergo deprotonation/ metalation onto the pyridinium to generate an organocupracycle stabilized by the Lewis basic iminobenzoyl moiety (**C**). Oxidative addition (**D**) into the alkenyl iodide and subsequent reductive elimination (**E**) affords the product.

Communications

In summary, we have described the first ligand-free Cucatalyzed direct alkenylation of electron-deficient heteroarenes. The reaction is believed to pass through a Cu^I/Cu^{III} catalytic cycle to provide a range of products in good to excellent yields. The reaction is highly chemoselective towards alkenyl iodides, introducing the possibility of preparing a scaffold from which a library of biologically interesting compounds could be constructed. Work is continuing to determine the full extent of these reactions as well as a detailed mechanistic analysis of the process.

Received: October 26, 2009 Published online: December 22, 2009

Keywords: bioactivity · copper · heterocycles · homogeneous catalysis · vinylidene ligands

- a) G. Zheng, L. P. Dwoskin, A. G. Deaciuc, J. Zhu, M. D. Jones, P. A. Crooks, *Bioorg. Med. Chem.* **2005**, *13*, 3899; b) K. J. Wilson, M. B. van Niel, L. Cooper, D. Bloomfield, D. O'Connor, L. R. Fish, A. M. MacLeod, *Bioorg. Med. Chem. Lett.* **2007**, *17*, 2643; c) L.-C. Campeau, K. Fagnou, *Chem. Soc. Rev.* **2007**, *36*, 1058; d) T. Laird, *Org. Process Res. Dev.* **2006**, *10*, 851.
- [2] a) M. G. P. Buffat, *Tetrahedron* 2004, 60, 1701; b) F.-X. Felpin, J. Lebreton, *Eur. J. Org. Chem.* 2003, 3693; c) J. S. Carey, L. Laffan, C. Thompson, M. T. Williams, *Org. Biomol. Chem.* 2006, 4, 2337; d) R. W. Dugger, J. A. Ragan, D. H. B. Ripin, *Org. Process Res. Dev.* 2005, *9*, 253.
- [3] For selected reviews, see: a) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* 2007, *107*, 174; b) M. Miura, T. Satoh, *Top. Organomet. Chem.* 2005, *14*, 55; c) T. Satoh, M. Miura, *Chem. Lett.* 2007, *36*, 200; d) L.-C. Campeau, K. Fagnou, *Chem. Commun.* 2006, 1253; e) I. Y. Seregin, V. Gevorgyan, *Chem. Soc. Rev.* 2007, *36*, 1173; f) J. Ellman, *Science* 2007, *316*, 1131.

- [4] a) A. Larivée, J. J. Mousseau, A. B. Charette, J. Am. Chem. Soc. 2008, 130, 52; b) J. J. Mousseau, A. Larivée, A. B. Charette, Org. Lett. 2008, 10, 1641; c) L.-C. Campeau, S. Rousseaux, K. Fagnou, J. Am. Chem. Soc. 2005, 127, 18020; d) J.-P. Leclerc, K. Fagnou, Angew. Chem. 2006, 118, 7945; Angew. Chem. Int. Ed. 2006, 45, 7781; e) L.-C. Campeau, D. J. Schipper, K. Fagnou, J. Am. Chem. Soc. 2008, 130, 3266; f) D. J. Schipper, M. El-Salfiti, C. J. Whipp, K. Fagnou, Tetrahedron 2009, 65, 4977.
- [5] a) M. Murakami, S. Hori, J. Am. Chem. Soc. 2003, 125, 4720;
 b) K. S. Kanyiva, Y. Nakao, T. Hiyama, Angew. Chem. 2007, 119, 9028; Angew. Chem. Int. Ed. 2007, 46, 8872; c) Y. Nakao, K. S. Kanyiva, T. Hiyama, J. Am. Chem. Soc. 2008, 130, 2448; d) S. H. Cho, S. J. Hwang, S. Chang, J. Am. Chem. Soc. 2008, 130, 9254.
- [6] For Cu-catalyzed direct functionalization, see: a) H.-Q. Do, O. Daugulis, J. Am. Chem. Soc. 2007, 129, 12404; b) H.-Q. Do, O. Daugulis, J. Am. Chem. Soc. 2008, 130, 1128; c) H.-Q. Do, R. M. K. Khan, O. Daugulis, J. Am. Chem. Soc. 2008, 130, 15185; d) F. Besselièvre, S. Piguel, F. Mahuteau-Betzer, D. S. Grierson, Org. Lett. 2008, 10, 4029; e) R. J. Phipps, N. P. Grimster, M. J. Gaunt, J. Am. Chem. Soc. 2008, 130, 8172; f) G. Brasche, S. L. Buchwald, Angew. Chem. 2008, 120, 1958; Angew. Chem. Int. Ed. 2008, 47, 1932; g) D. Zhao, W. Wang, F. Yang, J. Lan, L. Yang, G. Gao, J. You, Angew. Chem. 2009, 121, 3346; Angew. Chem. Int. Ed. 2009, 48, 3296; h) R. J. Phipps, M. J. Gaunt, Science 2009, 323, 1593.
- [7] See the Supporting Information for selected optimization.
- [8] J. A. Bull, J. J. Mousseau, A. B. Charette, Org. Lett. 2008, 10, 5485.
- [9] R. H. Jackson, M. M. Morrissey, M. A. Stills, M. F. Jarvis, J. Pharmacol. Exp. Ther. 1992, 262, 80.
- [10] ¹H NMR spectroscopy demonstrated that D incorporation for Table 3, entry 2 was at the 2,6-positions, although it could not be quantified because of peak broadening caused by the presence of Cu.
- [11] For Cu⁰ to Cu¹ reactions, see: a) T. Cohen, R. A. Schambach, J. Am. Chem. Soc. **1970**, 92, 3189; b) H. Weingarten, J. Org. Chem. **1964**, 29, 3624.