This article was downloaded by: [Tufts University] On: 08 October 2014, At: 13:20 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Phase-Transfer Catalyzed Mild Synthesis of 1,5-Diacyl Thiocarbohydrazides and Their Solventless Expeditious Transformation to 1,5-Diacyl Carbohydrazides

Zheng Li^a, Yanlong Zhao^a & Jingya Yang^a ^a Gansu Key Laboratory of Polymer Materials, Northwest Normal University, Lanzhou, Gansu, People's Republic of China Published online: 01 Feb 2007.

To cite this article: Zheng Li , Yanlong Zhao & Jingya Yang (2007) Phase-Transfer Catalyzed Mild Synthesis of 1,5-Diacyl Thiocarbohydrazides and Their Solventless Expeditious Transformation to 1,5-Diacyl Carbohydrazides, Phosphorus, Sulfur, and Silicon and the Related Elements, 182:1, 79-84, DOI: <u>10.1080/10426500600867240</u>

To link to this article: http://dx.doi.org/10.1080/10426500600867240

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with

primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Phase-Transfer Catalyzed Mild Synthesis of 1,5-Diacyl Thiocarbohydrazides and Their Solventless Expeditious Transformation to 1,5-Diacyl Carbohydrazides

Zheng Li Yanlong Zhao Jingya Yang Gansu Key Laboratory of Polymer Materials, Northwest Normal University, Lanzhou, Gansu, People's Republic of China

1,5-diacyl thiocarbohydrazides were efficiently synthesized by the reactions of thiocarbohydrazide with a variety of aroyl chlorides at r.t. using PEG-400 as a phasetransfer catalyst. Meanwhile, 1,5-diacyl thiocarbohydrazides were be expeditiously transformed into corresponding 1,5-diacyl carbohydrazides with periodic acid by r.t. grinding a under solventless condition. This protocol has advantages of a mild condition, fast reaction rate, high yield, and simple work-up procedure.

 ${\bf Keywords}~$ 1,5-diacyl thiocarbohydrazide; 1,5-diacyl carbohydrazide; phase-transfer catalysis; solventless

INTRODUCTION

Thiocarbohydrazide derivatives have attracted much attention in recent years due to their applications in the synthesis of heterocyclic compounds,¹ synthesis of transition metal complexes,² and pharmacological studies.³

Meanwhile, carbohydrazide derivatives are widely used as an oxygen scavenger (metal passivator) for water treatment systems, particularly for boiler-feed systems.⁴ They also are important intermediates for metal complexes.⁵ Recently, Zhao et al.⁶ reported the self-assembly properties of substituted carbohydrazides. Wang et al.⁷ reported the

Address correspondence to Zheng Li, Gansu Key Labaratory of Polymer Materials, College of Chemistry and Chemical Engineering, Northwest Normal University, Lanzhou, Gansu, 730070 People's Republic of China. E-mail: lizheng@nwnu.edu.cn

Received March 22, 2006; accepted April 30, 2006.

The authors thank the "Chunhui Project" of Ministry of Education of China (Z2004-1-62034) and Key Laboratory of Eco-Environment-Related Polymer Materials (Northwest Normal University), Ministry of Education of China, for the financial support of this work.



SCHEME 1

application of carbohydrazides as an inhibitor. However, the general synthetic method for carbohydrazide derivatives is to use extremely toxic phosgene as a starting material, which makes the process unsafe and environmentally unfriendly.⁸

In this article, we would like to report a mild, rapid, and high yielding method to prepare both 1,5-diacyl thiocarbohydrazides and corresponding 1,5-diacyl carbohydrazides in one route by using phase-transfer catalysis and a solventless grinding method, respectively.

RESULTS AND DISCUSSION

Reactions of thiocarbohydrazide with two equivalents of aroyl chlorides at r.t. using poly(ethylene glycol-400) (PEG-400) as a liquid-liquid phase transfer catalyst and sodium hydroxide as a base afforded 1,5diacyl thiocarbohydrazides (**1a-h**) in a high yield (Scheme 1). In fact, all the reactions were very efficient. Although water was used as a solvent together with methylene chloride, no any hydrolysates were observed. The different substituents on aryl rings had no obvious effect on the reaction rate and yield. The reaction also was suitable to the furan heterocycle (compound **1h**). All the products were easily separated from the mixture by extraction.

Further, compounds **1a-h** were ground with periodic acid in an agate mortar with a pestle under solvent-free conditions to readily afford 1,5diacyl carbohydrazides (**2a-h**) in an excellent yield. It was found that the periodic acid was quite an efficient reagent for the reactions. No byproducts were observed. The 1:1 molar ratio for the two reactants and about a 3–5 min reaction time were the optimized condition for giving a high yield (Table I).

From the room-temperature ¹H NMR spectrum of compound **1a** in DMSO- d_6 , an interesting result was observed. Although there were two different kinds of NH protons in compound **1a**, four significant

Compound	R	Reaction time (min)	M.P. (°C)	Yield (%) ^a
1a	C_6H_5	30	190–192	95
1b	$4-CH_3OC_6H_4$	25	184 - 185	93
1c	$2-CH_3C_6H_4$	30	184 - 186	89
1d	$3-CH_3C_6H_4$	30	156 - 158	90
1e	$2\text{-ClC}_6\text{H}_4$	25	184 - 186	89
1f	$4-ClC_6H_4$	20	188 - 190	91
1g	$4\text{-BrC}_6\text{H}_4$	20	224 - 225	91
1h	Fur-2-yl	25	130 - 132	95
2a	C_6H_5	3	206 - 207	96
2b	$4-CH_3OC_6H_4$	4	203 - 205	94
2c	$2-CH_3C_6H_4$	4	218 - 219	90
2d	$3-CH_3C_6H_4$	4	180 - 182	91
2e	$2\text{-ClC}_6\text{H}_4$	3	219 - 220	89
2f	$4-ClC_6H_4$	3	241 - 243	88
2g	$4\text{-BrC}_6\text{H}_4$	5	246 - 248	90
2 h	Fur-2-yl	4	204 - 205	95

TABLE I The Synthesis of Compounds 1a-h and 2a-h

^{*a*}Yields refer to the isolated product.

NH single peaks at 10.59, 10.47, 10.22, and 9.80 ppm appeared. However, the high temperature ¹H NMR spectrum at 60°C for compound **1a** showed that the four peaks for NH protons were changed into two broad peaks at 9.91 and 9.06 ppm. The similar situations were also observed for compounds **1b–h**. IR spectra of compounds **1a–h** showed the characteristic absorptions at 3194–3254 cm⁻¹ for N-H, 1676–1691 cm⁻¹ for C=O, and 1250–1259 for C=S.

In addition, ¹H NMR spectra of compounds **2a–h** in DMSO- d_6 showed proton peaks at 10.10–10.23 and 8.32–8.53 ppm for NH groups. IR spectra of compounds **2a–h** showed the characteristic absorptions at 3196–3288 cm⁻¹ for N-H and 1661–1688 cm⁻¹ for C=O.

In summary, we have developed an efficient method to prepare both 1,5-diacyl thiocarbohydrazides and 1,5-diacyl carbohydrazides in one route. This protocol has advantages of a mild reaction condition, short reaction time, high yield, low cost of the catalyst, and simple handling procedure. It is a good alternative to the synthesis of carbohydrazide derivatives without using extremely toxic phosgene as a starting material.

EXPERIMENTAL

IR spectra were recorded using KBr pellets on a Digilab 300 FTIR spectrophotometer and ¹H NMR spectra on a Mercury-400BB instrument using $(CD_3)_2SO$ as a solvent and Me₄Si as an internal standard.

Elemental analyses were performed on a Vario E1 Elemental Analysis instrument. Mass spectra were recorded on a QP-1000A GC-MS using the impact mode (70 eV). Melting-points were observed in an electrothermal melting-point apparatus. Aroyl chlorides were prepared by the reactions of corresponding substituted benzoic acids with thionyl chloride. Thiocarbohydrazide was prepared according to the literature procedure.⁹

General Procedure for the Preparation of Compounds 1a-h

To a solution of sodium hydroxide (10 mmol), thiocarbohydrazide (5 mmol), and PEG-400 (0.15 mmol) in 10 mL of water, aroyl chlorides (12 mmol) in 10 mL of CH_2Cl_2 were added dropwise. Then the mixture was stirred for an appropriate time indicated in Table I. The completion of the reaction was monitored by TLC using petroleum ether, acetone, and chloroform (5:2:2) as an eluent. Then the organic layer was separated, and the aqueous layer was extracted with CH_2Cl_2 (2X5 mL). The combined organic solution was dried by anhydrous Na_2SO_4 . The solvent was evaporated off, and the residue was crystallized from EtOH-DMF- H_2O (6:3:1) to give a product. The analytic data for compounds **1a–h** follow:

1a: White crystal. ¹H NMR: (DMSO- d_6 , 400 MHz): δ 10.59 (s, 1H, NH), 10.47 (s, 1H, NH), 10.22 (s, 1H, NH), δ 9.80 (s, 1H, NH), 7.47–7.95 (m, 10H, Ar-H). IR: (KBr, ν , cm⁻¹): 3254 (N-H), 1676(C=O), 1254(C=S). MS: m/z, 314 (M⁺). Anal. calcd. for C₁₅H₁₄N₄O₂S: C, 57.31; H, 4.49; N, 17.82. Found: C, 57.19; H, 4.56; N, 17.73.

1b: White crystal. ¹H NMR: (DMSO-*d*₆, 400 MHz): δ 10.39 (s, 1H, NH), 10.28 (s, 1H, NH), 10.10 (s, 1H, NH), 9.67 (s, 1H, NH), 7.01–7.91 (m, 8H, Ar-H), 3.82 (s, 6H, CH₃). IR: (KBr, ν , cm⁻¹): 3212 (N-H), 1687 (C=O), 1254 (C=S). MS: *m/z*, 374 (M⁺). Anal. calcd. for C₁₇H₁₈N₄O₄S:C, 54.53; H, 4.85; N, 14.96. Found: C, 54.42; H, 4.93; N, 14.88.

1c: White crystal. ¹H NMR: (DMSO-*d*₆, 400 MHz): δ 10.35 (s, 1H, NH), 10.23 (s, 1H, NH), 10.05 (s, 1H, NH), 9.61 (s, 1H, NH), 6.91–7.87 (m, 8H, Ar-H), 2.40 (s, 6H, CH₃). IR: (KBr, ν , cm⁻¹): 3199 (N-H), 1681 (C=O), 1253 (C=S). MS: *m/z*, 342 (M⁺). Anal. calcd. for C₁₇H₁₈N₄O₂S:C, 59.63; H, 5.30; N, 16.36. Found: C, 59.52; H, 5.25; N, 16.47.

1d: White crystal. ¹H NMR: (DMSO- d_6 , 400 MHz): δ 10.33 (s, 1H, NH), 10.22 (s, 1H, NH), 10.03 (s, 1H, NH), 9.58 (s, 1H, NH), 6.90–7.88 (m, 8H, Ar-H), 2.39 (s, 6H, CH₃). IR: (KBr, ν, cm⁻¹): 3194 (N-H), 1680 (C=O), 1252 (C=S). MS: m/z, 342 (M⁺). Anal. calcd. for C₁₇H₁₈N₄O₂S:C, 59.63; H, 5.30; N, 16.36. Found: C, 59.71; H, 5.37; N, 16.41.

1e: White crystal. ¹H NMR: (DMSO-*d*₆, 400 MHz): δ 10.63 (s, 1H, NH), 10.51 (s, 1H, NH), 10.26 (s, 1H, NH), δ 9.82 (s, 1H, NH), 7.51–8.01

(m, 8H, Ar-H). IR: (KBr, ν , cm⁻¹): 3215 (N-H), 1688 (C=O), 1255 (C=S). MS: m/z, 382 (M⁺). Anal. calcd. for C₁₅H₁₂Cl₂N₄O₂S:C, 47.01; H, 3.16; N, 14.62. Found: C, 46.91; H, 3.22; N, 14.70.

1f: White crystal. ¹H NMR: (DMSO- d_6 , 400 MHz): δ 10.64 (s, 1H, NH), 10.50 (s, 1H, NH), 10.28 (s, 1H, NH), δ 9.83 (s, 1H, NH), 7.53–8.00 (m, 8H, Ar-H). IR: (KBr, ν , cm⁻¹): 3216 (N-H), 1688 (C=O), 1256 (C=S). MS: m/z, 382 (M⁺). Anal. calcd. for C₁₅H₁₂Cl₂N₄O₂S:C, 47.01; H, 3.16; N, 14.62. Found: C, 47.12; H, 3.08; N, 14.56.

1g: White crystal. ¹H NMR: (DMSO-*d*₆, 400 MHz): δ 10.69 (s, 1H, NH), 10.56 (s, 1H, NH), 10.31 (s, 1H, NH), δ 9.86 (s, 1H, NH), 7.58–8.08 (m, 8H, Ar-H). IR: (KBr, ν , cm⁻¹): 3217 (N-H), 1689 (C=O), 1258 (C=S). MS: *m*/*z*, 470 (M⁺). Anal. calcd. for C₁₅H₁₂Br₂N₄O₂S:C, 38.16; H, 2.56; N, 11.87. Found: C, 38.07; H, 2.66; N, 11.96.

1h: White crystal. ¹H NMR: (DMSO-*d*₆, 400 MHz): δ 10.58 (s, 1H, NH), 10.47 (s, 1H, NH), 10.23 (s, 1H, NH), 9.74 (s, 1H, NH), 7.46–7.95 (m, 6H, Fu-H). IR: (KBr, ν , cm⁻¹): 3204 (N-H), 1679 (C=O), 1250 (C=S). MS: *m*/*z*, 294 (M⁺). Anal. calcd. for C₁₁H₁₀N₄O₄S:C, 44.89; H, 3.43; N, 19.04. Found: C, 44.94; H, 3.51; N, 19.14.

General Procedure for the Preparation of Compounds 2a-h

Compounds **1a–h** (0.5 mmol) and HIO₄.2H₂O (0.5 mmol) were added to an agate mortar (6 cm in diameter). Then the mixture was ground with a pestle at r.t. for the appropriate time indicated in Table I. The completion of the reaction was monitored by TLC using petroleum ether, acetone, and chloroform (2:3:3) as an eluent. The resulting solid was washed with H₂O (3X5 mL) and recrystallized from EtOH-DMF-H₂O (6:3:1) to give the product. The analytic data for compounds **2a–h** follows:

2a: White crystal. ¹H NMR (DMSO-d₆): δ 10.17 (s, 2H, NH), 8.34 (bs, 2H, NH), 7.36–7.75 (m, 9H, Ar-H). IR (KBr, ν , cm⁻¹): 3272 (N-H), 1668 (C=O). MS: m/z, 298 (M⁺). Anal. calcd. for C₁₅H₁₄N₄O₃: C, 60.40; H, 4.73; N, 18.78. Found: C, 60.29; H, 4.88; N, 18.84.

2b: White crystal. ¹H NMR (DMSO-d₆): δ 10.19 (s, 2H, NH), 8.36 (bs, 2H, NH), 7.38–7.77 (m, 8H, Ar-H), 3.85 (s, 6H, CH₃). IR (KBr, ν , cm⁻¹): 3212 (N-H), 1664 (C=O). MS: m/z, 358 (M⁺). Anal. calcd. for C₁₇H₁₈N₄O₅: C, 56.96; H, 5.10; N, 15.63. Found: C, 57.03; H, 5.02; N, 15.51.

2c: White crystal. ¹H NMR (DMSO-d₆): δ 10.16 (s, 2H, NH), 8.33 (bs, 2H, NH), 7.34–7.74 (m, 8H, Ar-H), 2.35 (s, 6H, CH₃). IR (KBr, ν , cm⁻¹): 3196 (N-H), 1662 (C=O). MS: m/z, 326 (M⁺). Anal. calcd. for C₁₇H₁₈N₄O₃: C, 62.54; H, 5.60; N, 17.16. Found: C, 62.66; H, 5.68; N, 17.21.

2d: White crystal. ¹H NMR (DMSO-d₆): δ 10.15 (s, 2H, NH), 8.32 (bs, 2H, NH), 7.34–7.73 (m, 8H, Ar-H), 2.34 (s, 6H, CH₃). IR (KBr, ν , cm⁻¹): 3271 (N-H), 1661 (C=O). MS: m/z, 326 (M⁺). Anal. calcd. for C₁₇H₁₈N₄O₃: C, 62.54; H, 5.60; N, 17.16. Found: C, 62.47; H, 5.51; N, 17.09.

2e: White crystal. ¹H NMR (DMSO-d₆): δ 10.20 (s, 2H, NH), 8.37 (bs, 2H, NH), 7.35–7.80 (m, 8H, Ar-H). IR (KBr, ν , cm⁻¹): 3264 (N-H), 1668 (C=O). MS: m/z, 366 (M⁺). Anal. calcd. for C₁₅H₁₂Cl₂N₄O₃: C, 49.07; H, 3.29; N, 15.26. Found: C, 49.14; H, 3.35; N, 15.19.

2f: White crystal. ¹H NMR (DMSO-d₆): δ 10.21 (s, 2H, NH), 8.38 (bs, 2H, NH), 7.37–7.83 (m, 8H, Ar-H). IR (KBr, ν , cm⁻¹): 3264 (N-H), 1668 (C=O). MS: m/z, 366 (M⁺). Anal. calcd. for C₁₅H₁₂Cl₂N₄O₃: C, 49.07; H, 3.29; N, 15.26. Found: C, 48.99; H, 3.21; N, 15.32.

2g: White crystal. ¹H NMR (DMSO-d₆): δ 10.22 (s, 2H, NH), 8.39 (bs, 2H, NH), 7.36–7.81 (m, 8H, Ar-H). IR (KBr, ν , cm⁻¹): 3265 (N-H), 1670 (C=O). MS: m/z, 454 (M⁺). Anal. calcd. for C₁₅H₁₂Br₂N₄O₃: C, 39.50; H, 2.65; N, 12.28. Found: C, 39.61; H, 2.70; N, 12.30.

2h: White crystal. ¹H NMR (DMSO-d₆): δ 10.10 (s, 2H, NH), 8.53 (bs, 2H, NH), 6.65–7.88 (m, 6H, Fu-H). IR (KBr, ν , cm⁻¹): 3263 (N-H), 1669 (C=O). MS: m/z, 278 (M⁺). Anal. calcd. for C₁₁H₁₀N₄O₅: C, 47.49; H, 3.62; N, 20.14. Found: C, 47.56; H, 3.54; N, 20.20.

REFERENCES

- (a) M. S. Chande, M. A. Pankhi, and S. B. Ambhaikar, *Indian J. Chem. Sect. B*, **39**, 603 (2000); (b) A. D. Naik, S. M. Annigeri, U. B. Gangadharmath, V. K. Revankar, and V. B. Mahale, *J. Inclusion Phenom. Macrocycl. Chem.*, **43**, 291 (2002); (c) B. S. Holla, K. N. Poojary, B. S. Rao, and M. K. Shivananda, *Eur. J. Med. Chem.*, **37**, 511 (2002); (d) J. R. Choudhari and B. N. Berad, *Indian J. Heterocycl. Chem.*, **14**, 151 (2004).
- [2] (a) O. V. Mikhailov, M. A. Kazymova, T. A. Shumilova, G. A. Chmutova, and S. E. Solovieva, *Transition Met. Chem.*, **30**, 299 (2005); (b) Q. J. Zhu, R. B. Dai, Q. X. Li, J. Q. Qu, L. F. Wang, J. G. Wu, and J. R. Cui, *Chin. J. Inorg. Chem.*, **17**, 911 (2001).
- [3] G. C. Saha, K. Khayer, M. R. Islam, and M. S. K. Chowdhury, *Indian J. Chem. Sect. B*, **31**, 547 (1992).
- [4] (a) M. Slovinsky and I. L. Woodridge, US Patent 4,269,717, October 6, 1980;
 (b) R. C. Strong, T. X. Richmond, V. K. Majestic, and T.X. Stafford, US Patent 5,078,966, August 14, 1990.
- [5] D. U. Warad, C. D. Satish, V. H. Kulkarni, and C. S. Bajgur, *Indian J. Chem. Sect. A*, 39, 415 (2000).
- [6] X. Zhao, X. Z. Wang, X. K. Jiang, Y. Q. Chen, Z. T. Li, and G. J. Chen, J. Am. Chem. Soc., 125, 15128 (2003).
- [7] D. Wang, M. Pechar, W. Li, P. Kopeckova, D. Bromme, and J. Kopecek, *Biochemistry*, 41, 8849 (2002).
- [8] J. Lange and H. Paul, US Patent 4,496,761, August 17, 1982.
- [9] X. H. Sun and Y. F. Liu, Chemistry (Chinese), 62, 46 (1999).