

Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: P. J. Foth, F. Gu, T. G. Bolduc, S. S. Kanani and G. Sammis, *Chem. Sci.*, 2019, DOI: 10.1039/C9SC03570B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

New Sulfuryl Fluoride-Derived Alkylating Reagents for the 1,1-Dihydrofluoroalkylation of Thiols

Paul J. Foth^a, Frances Gu^a, Trevor G. Bolduc^a, Sahil S. Kanani^a, Glenn M. Sammis^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

ABSTRACT: Herein, we report a new method for the one-pot synthesis of 1,1-dihydrofluoroalkyl sulfides by bubbling sulfuranyl fluoride (SO_2F_2) through a solution of the corresponding alcohol and thiol. The reaction proceeds through a new class of bis(1,1-dihydrofluoroalkyl) sulfate reagents, to afford the desired 1,1-dihydrofluoroalkyl sulfides in 55-90% isolated yields. The bis(1,1-dihydrofluoroalkyl) sulfates are highly chemoselective for thiol alkylation, and are unreactive with competing, unprotected nucleophiles, including amines, alcohols, and carboxylic acids.

Sulfuryl fluoride (SO_2F_2) has been utilized since the 1960s as an industrial fumigant,¹ but it has only recently attracted significant attention as a reagent for organic synthesis.^{1b,2} Studies have demonstrated that oxygen nucleophiles, such as alcohols (**1**),^{3,4} phenol derivatives (**2**),^{1b,5} oximes (**3**),⁶ and carboxylic acids (**4**),⁷ react with sulfuranyl fluoride to form fluorosulfate derivatives (Scheme 1).² The addition of a second equivalent of the oxygen nucleophile is kinetically slow, which allows fluorosulfate **5** to undergo subsequent transformations.⁸ Fluorosulfates (**5**) have been utilized as key reactants in a diverse range of reactions, including metal-catalyzed cross couplings,^{5c,9} click reactions,^{1b,5d} deoxyfluorinations,^{5b} alkylations,^{3a,4} nitrile syntheses,⁶ and the formation of amide bonds.^{7a}

All current synthetic methods rely on a very similar protocol for the formation of the fluorosulfate intermediate. Sulfuryl


fluoride is bubbled through a solution of the requisite oxygen nucleophile and a base, which is usually *N,N*-diisopropylethylamine (DIPEA), triethylamine, or a carbonate salt.^{1b,2-7} Despite the expansion of the use of sulfuranyl fluoride, the only reactive intermediates that have been identified are fluorosulfate derivatives (**5**), and no other sulfuranyl fluoride-derived reactive intermediates have been explored.¹⁰

We previously reported that bubbling sulfuranyl fluoride through a solution of 2,2,2-trifluoroethanol (**1a**) and an amine base, such as DIPEA or triethylamine, afforded trifluoroethyl fluorosulfate (**5a**, $\text{R}'' = \text{CH}_2\text{CF}_3$) in >90% yield.^{3a,11} Following up on the synthesis and reactivity of fluorosulfate **5a** in new transformations, we serendipitously discovered that even moderately more basic reagents,¹² such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) with a pK_{aH} of 12,¹³ afforded bis(trifluoroethyl) sulfate (**6a**) as the major product, and only trace amounts of fluorosulfate **5a** were detected by ^{19}F NMR spectroscopy. Bis(trifluoroethyl) sulfate (**6a**) is an intriguing species as there are only two previous methods for its synthesis,¹⁴⁻¹⁶ and there are no studies investigating its reactivity.¹⁷


We elected to study the reactivity of this new bis(trifluoroethyl) sulfate intermediate (**6a**) for the 1,1-dihydrofluoroalkylation of thiols for two reasons: (1) the resulting fluoroalkyl sulfides are important fluorinated motifs in pharmaceuticals and agrochemicals,¹⁸⁻²⁰ and (2) the more common sulfuranyl fluoride-derived reagent, trifluoroethyl fluorosulfate **5a**, is not an effective intermediate for thiol alkylation. Previous studies by Shreeve and coworkers indicated that the reaction between methane thiol (**7**), triethylamine, and **5a** afforded the corresponding fluoroalkyl sulfide (**8**) in only 31% yield and a 2.2:1 preference for reactivity at carbon compared to sulfur (Scheme 2, A).¹⁴

To examine the viability of bis(trifluoroethyl) sulfate (**6a**) as a thiol alkylating reagent, we treated a solution of **6a** and DBU with benzyl mercaptan (**9a**), which led to a 99% ^{19}F NMR yield of 1,1-dihydrofluoroalkylated product **10a** (Scheme 2, B). The

Previous work: representative SO_2F_2 -derived fluorosulfate intermediates.



This work: new SO_2F_2 -derived bis(1,1-dihydrofluoroalkyl) sulfate intermediates.



Scheme 1 Representative examples of sulfuranyl fluoride-mediated processes that utilize fluorosulfate reactive intermediates (**5**) and a new bis(trifluoroalkyl) sulfate (**6**).




^a Department of Chemistry, University of British Columbia, 2036 Main Mall, Vancouver, British Columbia V6T 1Z1, Canada

* Corresponding authors

Email: gsammis@chem.ubc.ca

Electronic Supplementary Information (ESI) available: [Experimental procedures, methods, and optimization data; NMR, IR, and MS data including ^1H , ^{13}C , and ^{19}F NMR spectra]. See DOI: 10.1039/x0xx00000x



A. Reactivity of fluorosulfate **5a** with methane thiol.¹⁴B. Reactivity to bis(trifluoroethyl) sulfate (**6a**) with benzyl mercaptan.C. Reactivity to fluorosulfate **5a** with benzyl mercaptan.


Scheme 2 Investigations into thiol alkylation using fluorosulfate (**5a**) and bis(trifluoroethyl) sulfate (**6a**). The yields in B and C were determined by ¹⁹F NMR spectroscopy using trifluorotoluene as an internal standard.

analogous reaction with a solution of DIPEA and fluorosulfate **5a** with **9a** afforded only 52% yield of **10a** (Scheme 2, C), which is comparable to the result reported by Shreeve and coworkers.¹⁴ The addition of DBU and benzyl mercaptan to a solution of DIPEA and **5a** improved the yield; however, the reaction led to an increase in the amount of free trifluoroethanol in solution, presumably resulting from nucleophilic attack at sulfur.²¹

This initial result is noteworthy as it represents the first example of the direct conversion of an unactivated 1,1-dihydrofluoroalcohol to the corresponding fluoroalkyl sulfide in a one-pot process. Thiol 1,1-dihydrofluoroalkylation can be achieved through nucleophilic displacement of activated trifluoroalkyl moieties,^{14,22} copper-catalyzed reactions with trifluoroalkyl iodide²³ or trifluorodiazalkanes,²⁴ or reductive trifluoroalkylthiolations.²⁵ All previous work relies on activated trifluoroalkyl moieties. This is particularly problematic for select activated trifluoroethyl derivatives and longer chain 1,1-dihydrofluoroalkyl groups that are only available from the corresponding alcohols, and thus require additional synthetic steps to activate.

With an established protocol for a one-pot, sequential reaction, investigations next focused on a one-step process, where the alkylation proceeds by bubbling sulfuryl fluoride through a solution of trifluoroethanol (**1a**), benzyl mercaptan (**9a**), and DBU. At room temperature, the reaction proceeded efficiently to afford desired trifluoroethylated product **10a** in 71% yield (Table 1, entry 1). The yield increased at 40 °C (entry 2), but there was no further improvement when the reaction was run at 60 °C (entry 3). We next examined the reaction performance in different solvents (entries 4–8).²⁶ Overall, the reaction was robust in a range of solvents, providing good yield in both polar aprotic (entries 4 and 5) and nonpolar solvents (entries 6 and 7). DCM was not as effective for this transformation, with product **10a** observed in only 38% yield (entry 8).²⁷

Table 1 Optimization of the one-pot, 1,1-dihydrofluoroalkylation of benzyl mercaptan (**9a**). DOI: 10.1039/C9SC03570B




entry	solvent	temp (°C)	¹⁹ F NMR yield ^a (%)
1	DMF	25	71
2	DMF	40	86
3	DMF	60	85
4	THF	40	81
5	ACN	40	77
6	hexane	40	75
7	benzene	40	67
8	DCM	40	38

All reactions were carried out following a one-pot procedure on 0.30 mmol scale of **9a** and a 1:1 v/v trifluoroethanol:solvent ratio. ^a Yield after 20 minutes, as determined by ¹⁹F NMR spectroscopy using trifluorotoluene as an internal standard.

The one-pot reaction generally affords high yields of the desired thioalkylated product regardless of the steric bulk or the electronics of the thiol (Scheme 3). Benzyl mercaptan (**9a**) and furfuryl mercaptan (**9b**) were both efficiently trifluoroethylated to form the corresponding sulfides **10a** and **10b** in good yields. 1-Decanethiol (**9c**), 2-phenylethanethiol (**9d**), methyl thioglycolate (**9e**), and 1,9-nonanedithiol (**9f**) were effective substrates for this transformation, affording mono- and dialkylated products **10c–10f** in 58% to 73% isolated yields. The reaction was insensitive to steric bulk alpha to the thiol, with both cyclohexyl mercaptan (**9g**) and triphenylmethanethiol (**9h**) alkylated in comparable yields (**10g** and **10h**, respectively). Electron rich and electron poor thiophenol derivatives were well tolerated, regardless of the position of the substituents (**10i–q**). Importantly, longer chain 1,1-dihydrofluoroalcohols, such as 2,2,3,3,3-pentafluoropropanol (**1b**), were viable starting materials; however extended reaction times were required. The isolated yields of **11a** and **11c** were increased to 90% and 89%, respectively, by conducting the reaction in a sequential one-pot manner,²⁸ where sulfuryl fluoride was first bubbled through a solution of DBU and trifluoroethanol followed by the addition of the requisite thiol.

We next examined the functional group tolerance of this new thiol 1,1-dihydrofluoroalkylation (Scheme 4). In substrates in which there is competition between alcohol and thiol alkylation, the reaction cleanly afforded good yields of the desired thiol 1,1-dihydrofluoroalkylation products (**10r** and **11r**). Carboxylic acids were also tolerated, with good isolated yields of thiol alkylated product **10s** using either the standard one-pot or the sequential one-pot protocols. The reaction was selective for the thiol over potential competing reactivity at the nitrogen atom of aniline and pyridine derivatives (**10t**, **11t**, and **10u**). As primary nitrogen derivatives were competent nucleophiles in reactions with trifluoroalkyl fluorosulfate, we next examined the reaction of L-cysteine ethyl ester. Under our sequential one-pot conditions, reactivity was only observed at sulfur to give **10v** in 63% isolated yield.²⁹






Scheme 3 Substrate scope for the 1,1-dihydrofluoroalkylation of thiols. Reaction conditions: SO₂F₂ (2.9 equiv) was bubbled through a solution of **9** (1 equiv), DBU (5.9 equiv) in 1:1 TFE/DMF (v/v), at 40 °C for 3 minutes, and then the reaction was stirred for an additional 17 min. All reactions were run on 1 mmol scale of thiol unless otherwise indicated. Isolated yields for the one-pot reaction are reported, with ¹⁹F NMR yields (using trifluorotoluene as the internal standard) provided in parentheses. ^aThe isolated yield has been corrected to account for disulfide or solvent impurities. See the Supplementary Information for details. ^bThe reaction was conducted on 0.5 mmol scale. ^cThe product was not isolated due to volatility. ^dThe reaction was stirred for 2 hours. ^ePentafluoropropanol:DMF (1:2 v/v) was used to form the reagent, and then the thiol was added to the reaction mixture. The reaction was stirred for 30 minutes.


Finally, we investigated whether we could achieve selective 1,1-dihydrofluoroalkylation using glutathione (**9w**). Glutathione is a challenging substrate as it has two carboxylic acids, an amine, and two amides that may interfere with the desired thiol fluoroalkylation. Gratifyingly, under our sequential, one-pot reaction conditions, the thiol was selectively alkylated in 92% ¹⁹F NMR yield and 59% isolated yield. Under similar reaction conditions, trifluoroethyl triflate only afforded moderate yields of **10w**.

Intrigued by the chemoselectivity of the bis(trifluoroethyl) sulfate reagent (**6a**), we next investigated its selectivity compared to trifluoroethyl fluorosulfate (**5a**)³⁰ in a competition experiment between benzyl mercaptan (**9a**) and piperidine (**12**)³¹ (Scheme 5).³² Addition of **9a** and **12** to a preformed solution of trifluoroethyl fluorosulfate and DIPEA afforded only a slight preference for thiol alkylation (Scheme 5, A). Trifluoroethanol (**1a**) was liberated in the course of the reaction, which is likely the result of the addition of thiol to the sulfur center of the fluorosulfate reagent.^{33,34} Better selectivity for thiol versus amine alkylation could be achieved by adding DBU with **9a** and **12**;³⁵ however, there was also a concomitant increase in the amount of **1a** (Scheme 5, B). Increasing the




Scheme 4 Functional group tolerance of the 1,1-dihydrofluoroalkylation reaction. Reaction conditions: SO₂F₂ (2.9 equiv) was bubbled through a solution of **9** (1 equiv), DBU (5.9 equiv) in 1:1 TFE/DMF (v/v), at 40 °C for 3 minutes, and then the reaction was stirred for an additional 17 min. All reactions were run on a 1 mmol scale unless otherwise indicated. Isolated yields for the one-pot reaction are reported, with ¹⁹F NMR yields (using trifluorotoluene as the internal standard) provided in parentheses. ^aThe one-pot reaction was stirred for 2 hours. ^bThe sequential, one-pot reaction was stirred for 30 minutes after addition of thiol. ^cThe product was converted to an HCl salt, and the reported yield has been corrected for solvent impurities.


A. Trifluoroethyl fluorosulfate (**5a**) reactivity.




B. Trifluoroethyl fluorosulfate (**5a**) reactivity with added DBU.



C. Bis(trifluoroethyl) sulfate (**6a**) reactivity.



D. Bis(trifluoroethyl) sulfate (**6a**) reactivity using pyrrolidine as the competition substrate.



Scheme 5 Competition experiments with sulfur and nitrogen nucleophiles.

amount of fluorosulfate reagent **5a** resulted in more amine alkylated product (**13**), but did not lead to a significantly better yield of **10a**. In contrast, formation of bis(trifluoroethyl) sulfate (**6a**) followed by addition of **9a** and **12** led to 90% yield of thiol alkylated product **10a**, and only trace amounts of **13** and trifluoroethanol (Scheme 5, C). Further increasing the equivalents of the alkylating reagent led to near quantitative yield of **10a** (>97%). Even when pyrrolidine (**14**), a more nucleophilic amine,³⁶ was used in a competition experiment, trifluoroethyl sulfide **10a** was obtained almost exclusively (Figure 5, D).³⁷

Overall, we have developed a new method for the 1,1-dihydrofluoroalkylation of thiols using a previously unexplored, sulfonyl fluoride derived bis(trifluoroethyl) sulfate reagent (**6a**). This protocol enables the one-pot activation and thiolation of 1,1-dihydrofluoroalcohols to afford industrially relevant moieties in high yields, regardless of the sterics or electronics of the starting thiol. In-situ generated bis(trifluoroethyl) sulfate (**6a**) is highly selective for thiols, even in the presence of unprotected alcohols, carboxylic acids, or amines, allowing for possible late-stage functionalization. Compared to trifluoroethyl fluorosulfate, the new bis(trifluoroethyl) sulfate reagent displays superior thiol alkylation chemoselectivity over both competing amine alkylation and reactivity at the sulfate center. Efforts to further explore this new class of bis(1,1-dihydrofluoroalkyl) reagents in the context of other reactions are currently underway.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the University of British Columbia (UBC), the Natural Sciences and Engineering Research Council of Canada (NSERC), the NSERC CREATE Sustainable Synthesis Program for support for PJF, an NSERC CGSM scholarship for TGB, and a UBC Work Learn International Undergraduate Research Award to SSK.

Notes and references

- (a) E. E. Kenaga, *J. Econ. Entomol.*, 1957, **50**(1), 1–6. (b) J. Dong, L. Krasnova, M. G. Finn, K. B. Sharpless, *Angew. Chem. Int. Ed.*, 2014, **53**(36), 9430–9448. (c) Dow AgroSciences Technical Bulletin, "Sulfonyl Fluoride Gas Fumigant," April 2002.
- For a representative review, see: L. Revathi, L. Ravindar, J. Leng, K. P. Rakesh, H. -L. Qin, *Asian J. Org. Chem.*, 2018, **7**(4), 662–682.
- For select examples of 1,1-dihydrofluoroalcohol activation, see: (a) M. Epifanov, P. J. Foth, F. Gu, C. Barrillon, S. S. Kanani, C. S. Higman, J. E. Hein, G. M. Sammis, *J. Am. Chem. Soc.*, 2018, **140**(48), 16464–16468. (b) M. R. Johnson, WO Pat. 124 456, 2014.
- For a representative example, see: A. Ishii, T. Yamazaki, M. Yasumoto, (Central Glass Company, Ltd.). US Pat. 8 426 645B2, 2013.
- For representative examples, see: (a) W. C. J. Firth, *J. Polym. Sci. B*, 1972, **10**, 637 – 641. (b) S. D. Schimler, M. A. Cismesia, P. S. Hanley, R. D. J. Froese, M. J. Jansma, D. C. Bland, M. S. Sanford *J. Am. Chem. Soc.*, 2017, **139**(4), 1452–1455. (c) P. S. Hanley, M. S. Ober, A. L. Krasovskiy, G. T. Whiteker, W. J. Kruper, *ACS Catal.*, 2015, **5**(9), 5041–5046. (d) J. Dong, K. B. Sharpless, L. Kwisnek, J. S. Oakdale, V. V. Fokin, *Angew. Chem. Int. Ed.*, 2014, **53**(36), 9466–9470.
- (a) J. Gurgar, J. Baker, V. V. Fokin, *Chem. Eur. J.*, 2019, **25**(8), 1906–1909. (b) W. -Y. Fang, H. -L. Qin, *J. Org. Chem.* 2019, **84**(9), 5803–5812.
- For representative examples, see: (a) S. -M. Wang, C. Zhao, X. Zhang, H. -L. Qin, *Org. Biomol. Chem.*, 2019, **17**, 4087–4101. (b) A. Ishii, M. Yasumoto, U. Koji, JP Pat. 155 248, 2009.
- For an early report on the stability of aryl fluorosulfates, see ref 5a.
- A. Ishii, T. Ishimaru, T. Yamazaki, JP Pat. 001 653 A, 2013.
- While other minor sulfonyl fluoride-derived products have been observed (see ref 3a), they have only been detected in <10% yield.
- When non-fluorinated alcohols are used, the corresponding alkyl fluorosulfate is highly reactive and leads to numerous degradation products. The 1,1-dihydrofluoroalkyl group makes the corresponding fluorosulfate more stable and less prone to degradation. For a study on the effect of halides alpha to the electrophilic center of S_N2 reactions, see: J. Hine, W. H. Brader Jr. *J. Am. Chem. Soc.*, 1953, **75**(16), 3964 – 3966.
- Stronger bases, such as KO-tBu and KHMDS, provide comparable results as DBU. Presumably, this is because more alkoxide is formed in the reaction, thus accelerating the conversion of **5a** to **6a**. For full base optimization, see the Supplementary Information.
- D. Granitza, M. Beyermann, H. Wenschuh, H. Haber, L. A. Carpino, G. A. Truran, M. Bienert, *J. Chem. Soc., Chem. Commun.*, 1995, **21**, 2223–2224.
- Shreeve *et al.* synthesized fluorosulfate **5a** in two separate steps using SOCl₂ followed by ClF. Fluorosulfate **5a** could be converted to the corresponding bis(trifluoroethyl) sulfate (**6a**) by running the reaction under solvent-free conditions using triethylamine and warming the reaction from -196 °C to room temperature. No subsequent reactivity studies were performed on **6a**. S. A. Kinkead, R. C. Kumar, J. M. Shreeve, *J. Am. Chem. Soc.*, 1984, **106**(24), 7496–7500.
- For a synthesis of bis(trifluoroethyl) sulfate (**6a**) from trifluoroethanol and sulfonyl chloride, see: W. V. Cohen, *J. Org. Chem.*, 1961, **26**(10), 4021–4026.
- We have previously detected the formation of bis(trifluoroethyl) sulfate (**6a**) as a minor byproduct in the formation of trifluoroethyl fluorosulfate (**5a**). See ref 3a.
- While a SciFinder® search indicates that bis(trifluoroethyl) sulfate (**6a**) was reported (WO Pat. 066 559, 2000), the reagent is not mentioned in this patent.
- For selected reviews on fluoroalkyl sulfides see: (a) S. Swallow, *Prog. Med. Chem.*, 2015, **54**, 65–133. (b) F. Leroux, P. Jeschke, M. Schlosser, *Chem. Rev.*, 2005, **105**(3), 827–856.
- For selected patents containing trifluoroethyl sulfur compounds see: (a) A. Adrien EP Kohler, B. Alig, A. Becker, A. Voerste, U. Gorgens, R. Fischer, W. A. Moradi, S. Cerezo-Galvez, J. Neumann, K. Ilg, H. -G. Schwarz, T. Gomibuchi, M. Ito, D. Yamazaki, K. Shibuya, E. Shimojo, WO Pat. 092 350, 2013. (b) F. Setzu, E. -J. Umemura, K. Sasaki, K. Tadauchi, T. Okutomi, K. Ohtsuka, S. Takahata, WO Pat. 042 188, 2003. (c) F. Kaiser, S. Gross, J. Langewald, A. Narine, WO Pat. 030 262, 2013. (d) B. Alig, S. Cerezo-Galvez, R. Fischer, A. Koehler, J. J. Hahn, K. Ilg, P. Loesel, O. Malsam, D. Portz, WO Pat. 004 028, 2015.
- For the effect of fluoroalkyl moieties on the properties in a molecule, as well as the importance of fluorine in the



- pharmaceutical and agrochemical industries, see: (a) W. A. Sheppard, *J. Am. Chem. Soc.*, 1963, **85**(9), 1314–1318. (b) B. Manteau, S. Pazenok, J. -P. Vors, F. R. Leroux, *J. Fluorine Chem.*, 2010, **131**, 140–158. (c) C. Ni, M. Hu, J. Hu, *Chem. Rev.*, 2015, **115**(2), 765–825. (d) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.*, 2008, **37**, 320–330. (e) P. Jeschke, *ChemBioChem.*, 2004, **5**(5), 570–589. (f) J. Wang, M. Sánchez-Roselló, J. L. Aceña, C. del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok, H. Liu, *Chem. Rev.*, 2014, **114**(4), 2432–2506. (g) B. Manteau, S. Pazenok, J. -P. Vors, F. R. Leroux, *J. Fluorine Chem.*, 2010, **131**, 140–158. (h) K. Müller, C. Faeh, F. Diederich, *Science*, 2007, **317**(5846), 1881–1886. (i) C. Isanbor, D. O'Hagan, *J. Fluorine Chem.*, 2006, **127**(3), 303–319.
- 21 See the Supplementary Information for full reaction details.
- 22 For representative examples, see: (a) Z. -Y. Long, Q. -Y. Chen, *J. Fluorine Chem.*, 1998, **91**(1), 95–98. (b) J. Hine, R. G. Ghirardelli, *J. Org. Chem.*, 1958, **23**(10), 1550–1552. (c) T. Nakai, K. Tanaka, H. Setoi, N. Ishikawa, *Bull. Chem. Soc. Jpn.*, 1977, **50**(11), 3069–3070. (d) R. F. Langler, N. A. Morrison, *Can. J. Chem.*, 1987, **65**(10), 2385–2389. (e) A. Adin, J. J. Looker, S. Y. Farid, I. R. Gould, S. A. Godleski, J. R. Lenhard, A. A. Muentner, L. C. Vishwakarma, P. A. Zielinski, US Pat. 6 054 260, 2000. (f) T. Umemoto, Y. Gotoh, *J. Fluorine Chem.*, 1986, **31**(2), 231–236. (g) Y. Pustovit, A. Alexenko, S. Trofymchuk, O. Lukin, A. A. Tolmachev, *Synthesis*, 2010, **7**, 1159–1165.
- 23 S. Chen, M. Zhang, X. Liao, Z. Weng, *J. Org. Chem.*, 2016, **81**(17), 7993–8000.
- 24 S. Hyde, J. Veliks, B. Liegault, D. Grassi, M. Taillefer, V. Gouverneur, *Angew. Chem. Int. Ed.*, 2016, **55**(11), 3785–3789.
- 25 (a) R. Wang, L. Jiang, Y. Wenbin, *J. Org. Chem.*, 2018, **83**(15), 7789–7798. (b) L. Jian, W. Yi, Q. Liu, *Adv. Synth. Catal.*, 2016, **358**(23), 3700–3705.
- 26 The reaction also works well using 1:2 and 1:4 TFE/DMF solvent ratios, which correlates to approximately 13 and 8 equivalents of TFE, respectively. For optimization of the amount of trifluoroethanol relative to solvent, see the Supplementary Information.
- 27 The alkylation step is slower in DCM than in more polar solvents, such as DMF.
- 28 Under the standard one-pot conditions, sulfonyl fluoride can react with the thiol, which leads to disulfides, among other products. The sequential, one-pot conditions prevent this side reaction from occurring. For a discussion of this disulfide, see the Supplementary Information.
- 29 ¹⁹F NMR spectroscopic analysis of the crude reaction mixture indicated that no amine 1,1-dihydrofluoroalkylation products were observed.
- 30 Reactivity was compared to fluoroalkyl fluorosulfates, such as **5a**, because they are the only other sulfonyl fluoride-derived reagent that can be generated from any 1,1-dihydrofluoroalcohol, and used in a one-pot process.
- 31 Piperidine was selected for the competition experiment as it was an effective substrate for amine trifluoroethylation (see ref 3a).
- 32 For full reactions conditions, see the Supplementary Information.
- 33 As attack at the sulfur atom with nitrogen was not typically seen under these conditions (see ref 3a), the reagent degradation is likely to be due to thiol attack at sulfur.
- 34 The amount of trifluoroethanol is due to nucleophile-mediated degradation of the alkylating reagent. Therefore, the percentage of trifluoroethanol was calculated relative to the total amount of benzyl mercaptan.
- 35 Similar moderate selectivity for *S*- versus *N*-trifluoroalkylation was observed with trifluoroethyl triflate. See the Supplementary Information for details.
- 36 T. Kanzian, T. A. Nigst, A. Maier, S. Pichl, H. Mayr, *Eur. J. Org. Chem.*, 2009, **36**, 6379–6385. DOI: 10.1039/C9SC03570B
- 37 The competition experiment was run using an equimolar amount of benzyl mercaptan (**9a**) and pyrrolidine, and 1.0 equiv. of sulfate **6a**. No amine alkylation product was detected by ¹⁹F NMR spectroscopy using trifluorotoluene as an internal standard. See the Supplementary Information for full reaction details.