Accepted Manuscript

Title: Direct benzylic functionalization of pyridines: Palladium-catalyzed mono- α -arylation of α -(2-pyridinyl)acetates with heteroaryl halides



Authors: Chaochao Jin, Kun Xu, Xiao Fan, Changyao Liu, Jiajing Tan

PII:	S1001-8417(19)30357-2
DOI:	https://doi.org/10.1016/j.cclet.2019.06.028
Reference:	CCLET 5059
To appear in:	Chinese Chemical Letters
Received date:	9 April 2019
Revised date:	10 June 2019
Accepted date:	14 June 2019

Please cite this article as: Jin C, Xu K, Fan X, Liu C, Tan J, Direct benzylic functionalization of pyridines: Palladium-catalyzed mono- α -arylation of α -(2-pyridinyl)acetates with heteroaryl halides, *Chinese Chemical Letters* (2019), https://doi.org/10.1016/j.cclet.2019.06.028

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Communication

Direct benzylic functionalization of pyridines: Palladium-catalyzed mono- α -arylation of α -(2-pyridinyl)acetates with heteroaryl halides

Chaochao Jin^a, Kun Xu^b, Xiao Fan^a, Changyao Liu^c, Jiajing Tan^{a,*} tanjj@mail.buct.edu.cn

^aDepartment of Organic Chemistry, College of Chemsitry, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing 100029, China

^bCollege of Chemistry and Pharmaceutical Engineering, Nanyang Normal University, Nanyang 473061, China ^cSchool of Food and Chemical Engineering, Beijing Technology and Business University, Beijing 100048, China

School of 1 oou and chemical Engineering, Deging Teenhology and Dusine

*Corresponding author.

Graphical Abstract



Herein, we report a Pd-catalyzed mono- α -arylation reaction to synthesize di-heteroaryl acetates in good yields and selectivities. The method is applicable to challenging heteroaryl substrate combinations, and exhibits great functional group tolerance.

ARTICLE INFO Article history: Received 9 April 2019 Received in revised form 10 June 2019 Accepted 11 June 2019 Available online

ABSTRACT

Keywords: a-Arylation Palladium Heterocycles

Herein, we report a Pd-catalyzed mono-a-arylation reaction for pyridine benzylic functionalization. This approach serves as an efficient alternative to synthesize di-heteroaryl acetates in good yields and selectivities. Moreover, the method is applicable to heteroaryl substrate combinations, and exhibits great functional group tolerance. A streamlined protocol also enables the rapid synthesis of diheteroaryl ketones. The synthetic value was also demonstrated by scale-up experiments

Pyridines Cross-coupling

Heterocycles are of great significance to the pharmaceutical industry due to their ability to interact with biological systems through polar interactions [1]. In particular, the unique electronic and conformational properties of pyridine rings renders them one simple yet privileged class of structure motifs in medicinal chemistry (Fig. 1) [2]. The incorporation of pyridine rings often had profound impacts on the biological properties of natural products. For instance, azatyrosine, with replacement of phenyl ring with pyridine, displays potent antibiotic and antitumor properties, when compared to corrsponding tyrosine [3].

The major progress of synthetic approaches to pyridine-containing molecules relied on the manipulation of pre-functionalized structural motifs [4]. Among those progress, the benzylic functionalization of pyridines is of great importance, as it provides access to useful building blocks such as α -pyridyl carbonyl, germinal dipyridyl moiety [5]. A common procedure for the benzylic functionalization of pyridines involved the deprotonation of picoline precursors with organometallic reagents, followed by the treatment with electrophiles (Scheme 1) [6]. Recently, alternative approaches have also been reported. For instance, Londregan and co-workers reported the PyBroP-mediated the addition of silyl ketene acetals to azine-*N*-oxides [7]. Nevertheless, these protocols showed broad synthetic appeal and were useful in certain contexts, but the reaction reagent nature still limited their utilities to a certain degree [6,8].

Pyridine-containing bio-active molecules



Fig. 1. Pyridine motifs in medicinal chemistry. Classic pyridine benzylic functionalization



Scheme 1. Pyridine benzylic functionalization.

Given our group's continuous efforts in heterocyclic chemistry and transition metal catalysis [9], our attention was therefore drawn to employ the Pd-catalyzed α -arylation reaction to achieve the pyridyl-benzylic functionalization. Since the seminal work from Buchwald, Hartwig and Miura groups, the Pd-catalyzed α -arylation of carbonyl moiety has emerged as an effective Csp^2-Csp^3 bond formation strategy [10]. A wide array of protocols has been found to be effective to connect highly functionalized substrates. Moreover, these progresses have transformed the way in which synthetic chemists designed and implemented synthesis on both bench-top and industrial scales [11]. Notwithstanding such progress, the substrate scope of both coupling partners remains challenging with respect to the pyridine functionality. In principle, any Pd-catalyzed α -arylation reactions that is tolerable to both the Lewis basic functionality as well as the inherent electronic deficient property of pyridine rings, would be highly appreciated, and offer alternative efficient solution to pyridine benzylic functionalization [9a,12]. Another critical task is to identify the suitable reaction conditions to avoid the inhibition caused by the enolization of 2-(pyridin-2-yl)acetate and resulting coordination with palladium center as the ligand. Despite these challenges, we would like to report a highly efficient pyridyl-benzylic functionalization protocol *via* the palladium-catalyzed mono- α -arylation reaction of α -(2-azaheteroaryl) acetates with heteroaryl halides. Due to the immense usefulness of pyridine motifs, this protocol in principle should be of great interests to the community.

Table 1

Optimization of the reaction conditions.^a

N la	Br CO ₂ Et + 2a	Pd (5 mol%) Ligand (5 mol%) Base (3 equiv.) Solvent, 100 °C, N		Jaa
Entry	Pd catalyst	Ligand	Solvent	Yield (%) ^b
1	$Pd(OAc)_2$	L_1	Dioxane	23
2	$Pd(OAc)_2$	L_2	Dioxane	29
3	$Pd(OAc)_2$	L_3	Dioxane	20
4	$Pd(OAc)_2$	L_4	Dioxane	85
5	$Pd(OAc)_2$	L_5	Dioxane	69
6	$Pd(OAc)_2$	L_6	Dioxane	96
7	Pd ₂ (dba) ₃	L_6	Dioxane	94
8	[PdCl(allyl)] ₂	L_6	Dioxane	91
9	PdCl ₂	L_6	Dioxane	90
10	$Pd(OAc)_2$	L_6	Glyme	13
11	$Pd(OAc)_2$	L_6	DCE	67
12	$Pd(OAc)_2$	L_6	Toluene	79
13°	$Pd(OAc)_2$	L_6	Dioxane	50
14 ^d	Pd(OAc) ₂	L_6	Dioxane	82

^a 1a (0.3 mmol), 2a (0.4 mmol), Pd catalyst (0.015 mmol), ligand (0.015 mmol), and Cs₂CO₃ (0.9 mmol) in dioxane (1 mL), 100 °C, N₂, 10 h.

^b Isolated yields.

° Na2CO3 used instead.

^d K₃PO₄ used instead.

We initiated the optimization study with commercially available ethyl 2-(pyridin-2-yl)acetate (**1a**) and 6-bromoquinoline (**2a**) (Table 1). Among the phosphine ligands examined, the highest yield of ethyl 2-(pyridin-2-yl)-2-(quinolin-6-yl)acetate (**3aa**) was obtained in the presence of X-Phos (**L**₆) (Fig. 2) [13]. Next, a series of palladium sources were examined, all of which led to slightly lower yields, when compared to palladium acetate. Attention was then switched to determine the optimal solvent for the reaction. The control experiments revealed that dioxane was the ideal choice (entries 8-12). Keeping all other variables constant, the coupling reactions with either Na₂CO₃ or K₃PO₄ were then performed, and both failed to offer better results (entry 6 *vs.* entry 13-14).





Scheme 2. Substrate scope. Reaction conditions: 1a (0.3 mmol), 2 (0.4 mmol), Pd(OAc)₂ (0.015 mmol), X-Phos (0.015 mmol) and Cs₂CO₃ (0.9 mmol) in dioxane (1 mL), 100 °C, N₂, 10 h. Isolated yields.

With the optimal conditions in hands, the transformation was demonstrated on a range of heteroaryl halides. As shown in Scheme 2, investigations of the heteroaryl bromide scope showed that quinoline-based heteroaryl bromides were viable substrates to give desired products in excellent yields (**3aa-3ae**, 75%-96% yields). Moreover, pyridine-containing substrates proved to be well-tolerant to give the desired adducts **3af-3am** in 83%-97% yields. It is worth mentioning that these substrate classes are discovered to be non-sensitive to the electronic effect of substitution groups on the pyridine rings. Besides, thiophene derived substrates, which are also widely present in medicinally relevant compounds, underwent the reaction smoothly to deliver coupling products (**3an, 3bd, 3be**, 63%-83% yields). As expected, when simple bromobenzene was employed, the desired arylation product **3ao** was obtained in 95% yield.

Next, we examined the generality of our coupling protocol with respect to the heteroaryl substituted acetates (Scheme 3). The methyl and *t*-Bu 2-(pyridin-2-yl)acetate are viable substrates, providing quick access to desired products (**3ba-3cg**) in good yields (63%-97% yields). We were also glad to discovery that the protocol can be extended to quinolidyl acetates, which performed well under the standard conditions to give **3dh** in 88% yield. More importantly, the mono-arylation product could be utilized to further react with bromocyclopropane to construct the all carbon quaternary center on **5**, which is presumably involve β -elimination before coupling [14,15]. Unfortunately, when we employed **1e**, the palladium complex **6** was obtained instead of the corresponding coupling products (Scheme 4). Single crystals suitable for X-ray diffraction analysis were grown from a mixture of dichloromethane and petroleum ether. The central palladium atom displayed a square-planar arrangement with the nitrogen atoms of the pyridine ring and oxygen of the enol moiety situated mutually in the *trans* position.



Scheme 3. Substrate scope. Reaction conditions: 1 (0.3 mmol), 2 (0.4 mmol), Pd(OAc)₂ (0.015 mmol), X-Phos (0.015 mmol) and Cs₂CO₃ (0.9 mmol) in dioxane (1 mL), 100 °C, N₂, 10 h. Isolated yields.



Scheme 4. Ketone substrate and resulting palladium complex. Crystal data: $C_{26}H_{20}N_2O_2Pd$, M = 498.84, triclinic, P $\overline{1}$, a = 9.8373(3), b = 12.0510(4), c = 17.9512(6) Å, a = 105.345(3), $\beta = 95.871(3)$, $\gamma = 93.103(3)$, V = 2034.16(11) Å³, Z = 4, Mo Ka radiation ($\lambda = 0.71073$). For details, see Supporting information.

Alteration of the halide fragments was evaluated next, using heteroaryl chlorides, triflates and tosylates as the coupling partners. Under optimized conditions, the arylated products **3aa** were still obtained in good chemical yields, respectively. Additionally, our protocol was effective to incorporate benzothiazole units into arylation products using 5-chloro-2-methylbenzothiazole (Scheme 5). Noteworthy, all the above-mentioned reactions showed no evidence of over-arylation by-products, which is a major side pathway for the arylation reactions of α -carbonyl compounds [16].



Scheme 5. Substrate scope.

This arylation protocol is amenable to gram-scale preparation of **3aa**. We reacted 6 mmol of **1a** with a slight excess amount (1.3 equiv., 7.8 mmol) of **2a**. Under the same reaction conditions, **3aa** was produced in 91% yield (Scheme 6). Next, the hydrolysis and subsequent decarboxylation went without issue to provide the diheteroaryl ketone product **10** in good efficiency. This one-pot, two-step protocol was further demonstrated by the quick synthesis of **11** and **12**, which often displayed photochemical reactivities [17].



Scheme 6. Synthetic applications.

In conclusion, we have developed an efficient Pd-catalyzed mono- α -arylation reaction of α -(2-azaheteroaryl) ester with heteroaryl halides. The broad heterocyclic substrate scope enabled the rapid synthesis of a wide array of functionalized α, α' -di-heteroaryl acetate products with consistently good yields. Remarkably, our protocol proved to be not only efficient in scale-up reactions, but further

functionalization was also readily possible, setting the base for rapidly accessing diheteroaryl ketone derivatives. In comparison to classical approaches for pyridine benzylic functionalization, we anticipate that our catalytic protocol will provide an efficient alternative for divergent parallel synthesis efforts.

Acknowledgments

Tan is grateful to the support by the National Natural Science Foundation of China (No. 21702013), Beijing Natural Science Foundation (No. 2184115) and the Fundamental Research Funds from the Central Universities (No. XK1802-6, buctrc201721) in Beijing University of Chemical Technology.

References

- [1] (a) R.E. Dolle, B.L. Bourdonnec, K. Worm, et al., Comb. Chem. 12 (2010) 765-806;
 - (b) A. Facchetti, Chem. Mater. 23 (2011) 733-758;
 - (c) L.M. Blair, J. Sperry, J. Nat. Prod. 76 (2013) 794-812;
 - (d) E. Vitaku, D.T. Smith, J.T. Njardarson, J. Med. Chem. 57 (2014) 10257-10274;
 - (e) U.H.F. Bunz, Acc. Chem. Res. 48 (2015) 1676-1686;
 - (f) R.K. Alan, A.R. Christopher, F.V.S. Eric, et al., Comprehensive Heterocyclic Chemistry III, Elsevier, Oxford, 2008;
 - (g) K.C. Majumdar, S.K. Chattopadhyay, Heterocycles in Natural Product Synthesis, Wiley-VCH, Weinheim, 2011.
- [2] (a) J. Alvarez-Builla, J.J. Vaquero, J. Barluenga, Modern Heterocyclic Chemistry, Wiley-VCH, Weinheim, 2011;
 - (b) M. Baumann, I.R. Baxendale, Beilstein, J. Org. Chem. (9) 2013 2265-2319;
 - (c) M.D. Hill, Chem. -Eur. J. 16 (2010) 12052-12062;
 - (d) J.S. Carey, D. Laffan, C. Thomson, et al., Org. Biomol. Chem. 4 (2006) 2337-2347;
 - (e) M. Schlosser, F. Mongin, Chem. Soc. Rev. 36 (2007) 1161-1172;
 - (f) L. Ackermann, H.K. Potukuchi, A.R. Kapdi, et al., Chem. -Eur. J. 16 (2010) 3300-3303;
 - (g) Y. S. Kumar, F. -R. N. Khan, Chin. Chem. Lett. 28 (2017) 1607-1612;
 - (h) A. P. Krinochkin, D. S. Kopchuk, N. V. Chepchugov, et al., Chin. Chem. Lett. 28 (2017) 1099-1103.
- [3] R.A. Aycock, D.B. Vogt, N. T. Jui, Chem. Sci. 8 (2017) 7998-8003.
- [4] (a) A. Kotschy, G. Timári, Heterocycles from Transition Metal Catalysis, Springer, Dordrecht, 2005;
 (b) S. Schröter, C. Stock, T. Bach, Tetrahedron, 61 (2005) 2245-2267;
 (c) M. Beller, C. Baler, Transition Metals for Organic Surface Pailling, Physical Prior Chemicals, 2nd education of the Chemical Pailling, Physical Physical Pailling, Physical Phys
- (c) M. Beller, C. Bolm, Transition Metals for Organic Synthesis: Building Blocks and Fine Chemicals, 2nd ed. Wiley-VCH, Weinheim, 2004. [5] (a) H.-Y. Lin, B. B. Snider, J. Org. Chem. 77 (2012) 4832-4836;
- (b) J.A. Lowe, D.L. Hageman, S.E. Drozda, et al., J. Med. Chem. 37 (1994) 3789-3811;
- (c) J.J. Mousseau, A. Larivée, A.B. Charette, Org. Lett. 10 (2008) 1641-1643;
- (d) S. Duez, A.K. Steib, S.M. Manolikakes, et al., Angew. Chem. Int. Ed. 50 (2011) 7686-7690;
- (e) K.P. Bogeso, A.V. Christensen, J. Hyttel, et al., J. Med. Chem. 28 (1985) 1817-1828;
- [6] (a) R.B. Woodward, E.C. Kornfeld, Org. Synth. 29 (1949) 44-44;
 (b) W.G. Kofron L.M. Baclawski, Org. Synth. 52 (1972) 75-75.
- (c) R. Zhu, G. Cheng, C. Jia, et al., J. Org. Chem. 81 (2016) 7539-7544.
- [7] (a) A.T. Londregan, S. Jennings, L. Wei, Org. Lett. 12 (2010) 5254-5257;
- (b) A.T. Londregan, S. Jennings, L. Wei, Org. Lett. 13 (2011) 1840-1843.
- [8] (a) P.S. Fier, J. Am. Chem. Soc. 139 (2017) 9499-9502;
- (b) D.D. Zhai, X.Y. Zhang, Y.F. Liu, et al., Angew. Chem. Int. Ed. 57 (2018) 1650-1653. and references therein. [9] (a) J.J. Tan, Y. Chen, H. Li, et al., J. Org. Chem. 79 (2014) 8871-8876;
- (b) H.Q. Wang, W.T. Xu, Z.Q. Wang, et al., J. Org. Chem. 80 (2015) 2431-2435; (c) L. Liu, C. Tan, R. Fan, et al., Org. Biomol. Chem. 17 (2019) 252-256.
- [10] (a) S. Lee, N.A. Beare, J.F. Hartwig, J. Am. Chem. Soc. 123 (2001) 8410-8411;
- (b) M. Jorgensen, S. Lee, X. Liu, et al., J. Am. Chem. Soc. 124 (2002) 12557-12565;
- (c) T. Hama, X. Liu, D.A. Culkin, et al., J. Am. Chem. Soc. 125 (2003) 11176-11177;
- (d) X. Liu, J.F. Hartwig, J. Am. Chem. Soc. 126 (2004) 5182-5191;
- (e) T. Hama, J.F. Hartwig, Org. Lett. 10 (2008) 1545-1548;
- (f) M.R. Biscoe, S.L. Buchwald, Org. Lett. 11 (2009) 1773-1775;
- (g) T. Hama, D.A. Culkin, J.F. Hartwig, J. Am. Chem. Soc. 128 (2006) 4976-4985;
- (h) B. Zheng, T. Jia, P.J. Walsh, Org. Lett. 15 (2013) 4190-4193;
- (i) R. Martin, S.L. Buchwald, Angew. Chem. Int. Ed. 46 (2007) 7236-7239;
- (j) G.D. Vo, J.F. Hartwig, Angew. Chem. Int. Ed. 47 (2008) 2127-2130;
- (k) R. Martin, S.L. Buchwald, Org. Lett. 10 (2008) 4561-4564;
- (1) D.A. Culkin, J.F. Hartwig, Acc. Chem. Res. 36 (2003) 234-245;
- (m) F. Bellina, R. Rossi, Chem. Rev. 110 (2010) 3850-3850;
- (n) C.C.C. Johansson, T.J. Colacot, Angew. Chem. Int. Ed. 49 (2010) 676-707;
- (o) Z. Liu, M. Li, B. Wang, et al., Org. Chem. Front. 5 (2018) 1870-1876;
- (p) G. Gao, Y. Fu, M. Li, et al., Adv. Synth. Catal. 359 (2017) 2890-2894;
- (q) K. Ablajan, G.B Panetti, X. Yang, et al., Adv. Synth. Catal. 359 (2017) 1927-1932;
- (r) G. Saini, P. Kumar, G.S. Kumar, et al., Org. Lett. 20 (2018) 441-444;
- (s) I. Astarloa, R. SanMartin, M.T. Herrero, et al., Adv. Synth. Catal. 360 (2018) 1711-1718;
- (t) D.J. Leonard, J.W. Ward, J. Clayden, Nature 562 (2018) 105-109.
- [11] (a) Á. Molnár, Palladium-Catalyzed Coupling Reactions Practical Aspects and Future Developments, Wiley-WCH, Weinheim, 2013;
 (b) M.L. Crawley, B.M. Trost, Applications of Transition Metal Catalysis in Drug Discovery and Development, Wiley, New York, 2012;
 (c) C. Torborg, M. Beller, Adv. Synth. Catal. 351 (2009) 3027-3043.
- [12] (a) M.A. Oberli, S.L. Buchwald, Org. Lett. 14 (2012) 4606-4609;
 (b) P.E. Maligres, J. Li, S.W. Krska, et al., Angew. Chem. Int. Ed. 51 (2012) 9071-9074;
 (c) J.C. Tellis, D.N. Primer, G.A. Molander, Science 345 (2014) 433-436.

- [13] (a) X. Huang, K.W. Anderson, D. Zim, J. Am. Chem. Soc. 125 (2003) 6653-6655;
 (b) N.C. Bruno, M.T. Tudge, S.L. Buchwald, Chem. Sci. 4 (2013) 916-920;
 (c) P. Novak, R. Martin, Curr. Org. Chem. 15 (2011) 3233-3262.
- [14] (a) C.J. Douglas, L.E. Overman, Proc. Natl. Acad. Sci. U. S. A. 101 (2004) 5363-5367;
 (b) J. Christoffers, A. Baro, Adv. Synth. Catal. 347 (2005) 1473-1482;
 (c) K.W. Quasdorf, L.E. Overman, Nature 516 (2014) 181-191.
- [15] (a) J. Gu, X. Wang, W. Xue, Org. Chem. Front. 2 (2015) 1411-1421;
- (b) X. Wang, S.Z. Stankovich, R.A. Widenhoefer, Organometallics 21 (2002) 901-905.
- [16] (a) W.A. Moradi, S.L. Buchwald, J. Am. Chem. Soc. 123 (2001) 7996-8002;
 (b) T. Hama, J.F. Hartwig, Org. Lett. 10 (2008) 1549-1552;
 (c) F. Churruca, R. SanMartin, R. Tellitu, Tetrahedron Lett. 44 (2003) 5925-5929;
 (d) K.B. Urkalan, M.S. Sigman, Angew. Chem. Int. Ed. 48 (2009) 3146-3149;
 (e) F. Churruca, R. SanMartin, M. Carril, Tetrahedron 60 (2004) 2393-2408.
- [17] G. Favaro, F. Ortica, A. Romani, J. Photochem. Photobiol. C 16 (2013) 22-45.