

Rapid Access to 3-Aryltetralin Skeleton via C(sp³)-H Bond Functionalization: Investigation on the Substituent Effect of Aromatic Ring Adjacent to C-H Bond in Hydride Shift/Cyclization Sequence

Keiji Mori, Shosaku Sueoka, and Takahiko Akiyama*

Department of Chemistry, Faculty of Science, Gakushuin University, Mejiro, Toshima-ku, Tokyo 171-8558

(Received October 22, 2011; CL-111039; E-mail: takahiko.akiyama@gakushuin.ac.jp)

The concise construction of 3-aryltetralin skeleton via hydride shift mediated C-H bond functionalization was achieved. In this process, the benzylic [1,5]-H shift occurred smoothly to furnish tetralin derivatives in good to excellent chemical yields. The electronic and steric properties of the aromatic ring adjacent to the C-H bond influenced significantly the reactivity of this transformation.

The direct and selective replacement of carbon-hydrogen bonds with new bonds (C-H functionalization) represents an important and longstanding goal in synthetic organic chemistry.¹ Considering the high abundance of C-H bonds, the precise one-step substitution of C-H bonds with C-C and/or C-Y bonds (Y = O, N, etc.) without disruption of the surrounding molecular structure and the prefunctionalization to C-X bonds (X = halogen, OSO₂CF₃, etc.) offer a promising approach for the synthesis of various complex molecules.

Recently, the C(sp³)-H bond functionalization via hydride shift/cyclization, namely, "internal redox process," has attracted much attention for its unique features (Scheme 1).² Its key feature is the [1,5]-hydride shift of the C(sp³)-H bond α to the heteroatom. Subsequent 6-endo cyclization affords heterocycle **2**. Although C-H functionalization is, in general, promoted by a transition-metal catalyst, this type of C-H functionalization process typically proceeds under thermal conditions or, in some cases, under the Brønsted or Lewis acid catalysis.³⁻⁶


Whereas a range of related reactions with heteroatom-containing substrates have been reported, the corresponding C-version had been overlooked until quite recently. This is because the key [1,5]-H shift required the electronic assistance of an adjacent heteroatom for the stabilization of the carbocation generated by the hydride shift. Recent work on the internal redox process has led to the realization of the benzylic, nonadjacent heteroatom hydride shift/cyclization sequence.⁷ The groups of Chatani^{7a} and He^{7b} independently reported that the benzylic C-H bond without an adjacent heteroatom could also participate in this type of catalytic cycloisomerization reaction. Quite

recently, our group has disclosed^{8a} that the employment of benzylidene barbiturate as the electrophilic portion enabled the benzylic [1,5]-hydride shift of the C-H bond at the α -position of the phenethyl moiety, affording 3-phenyltetralin in excellent chemical yield.⁸ Although there are several elegant pathways leading to benzylic C-H bond functionalization, details of the substituent effect of the aromatic ring adjacent to the C-H bond remain to be investigated.


Herein we report our study of the benzylic hydride shift/transfer system, focusing on the substituent effect of phenethyl aromatic ring (Scheme 2). We have found that the electronic nature of the aromatic group of the phenethyl moiety influenced significantly the reactivity, and the desired compounds were obtained in excellent chemical yields with low catalyst loading (as low as 0.5 mol %).

According to our previous report,^{8a} various benzylidene barbiturates **5** were synthesized from salicylic acid derivatives. The preparation of *p*-tolyl analog **5b** is illustrated in Scheme 3 as a representative example. The Sonogashira coupling of triflate **3**, which is readily synthesized from commercially available salicylic acid, with *p*-tolyl acetylene afforded adduct **4** in 90% yield. Hydrogenation, reduction of the ester group, oxidation of the resulting alcohol, and condensation with 1,3-dimethylbarbituric acid gave **5b** in 76% yield from **4** (Scheme 3).


With the desired substrates in hand,⁹ the substituent effect of the aromatic ring was surveyed under optimal conditions (cat. Sc(OTf)₃, ClCH₂CH₂Cl, reflux).^{8a,10} Phenyl-substituted substrate **5a** underwent the internal redox process in 24 h by means of 5 mol % Sc(OTf)₃ (Table 1, Entry 1). Interestingly, the



Scheme 2. Rapid access to tetralin skeleton.



Scheme 1. C(sp³)-H functionalization via internal redox process.



Scheme 3. Preparation of benzylidene barbiturate.

Table 1. Investigation of the effect of aromatic ring on phenethyl moiety^a

Reaction scheme showing the conversion of compound **5** to compound **6**. Compound **5** is a benzylidene barbiturate derivative with an Ar group. The reaction conditions are: cat. Sc(OTf)₃, ClCH₂CH₂Cl, reflux. Compound **6** is a 3-aryltetralin derivative.



Entry	Ar	Catalyst loading /mol %	Time /h	Yield /%	
1		6a	5	24	94
2		6b	5	2	98
3		6b	1	10	95
4		6c	1	8	96
5		6c	0.5	12	90
6		6d	30	24	88
7		6d	5	24	25
8		6e	5	24	79
9		6e	3	24	47
10		6f	5	24	96
11		6f	1	24	6
12		6g	30	24	75
13		6g	5	24	trace

^aUnless otherwise noted, all reactions were performed with 0.2 mmol benzylidene barbiturate and a catalytic amount of Sc(OTf)₃ in ClCH₂CH₂Cl (2.0 mL) at refluxing temperature.

electronic nature of the aromatic ring changed the reactivity significantly: the reaction of **5b** with *p*-tolyl group (electron-rich group) was completed within 2 h to give **6b** in excellent chemical yield (Entry 2 vs. Entry 1). In addition, the catalyst loading could be reduced to 1 mol % without sacrificing both chemical yield and reaction rate (95%, 10 h, Entry 3). Notably, even 0.5 mol % catalyst sufficed to afford **5c** with a *p*-methoxyphenyl group in excellent chemical yield (Entry 5). On the other hand, an electron-withdrawing substituent lowered the reactivity considerably: 30 mol % catalyst was required to achieve a satisfactory chemical yield for **6d** (Ar = *p*-chlorophenyl, 88%, 24 h, Entry 6).

Further investigation revealed that the steric factor also strongly affected the reactivity (Entries 8–13). To obtain **6f** (Ar = *o*-tolyl) in excellent chemical yield, a longer reaction time and an increased catalyst loading (24 h with 5 mol % catalyst) were required compared with *p*-tolyl **5b** (Entry 10 vs. Entry 3). Interestingly, **5g** with a mesityl group, which has high electron density compared with **5b**, showed considerably low reactivity: the chemical yield was at most 75% even with 30 mol % catalyst loading (Entry 12).

We assumed that the hyperconjugative interaction of the σ^* orbital of the C–H bond with neighboring substituents is

**Figure 1.** Rationalization of the difference of hydride transfer.**Figure 2.** Further examination of substrate scope.

responsible for the promotion of the desired [1,5]-hydride shift (Figure 1). The electronic assistance from the n -orbital of heteroatom to the σ^* C–H orbital is the most important element in the case of heteroatom analog **1** (X = NR and O).¹¹ On the other hand, the orbital overlap between the π -orbital of the aromatic ring and the σ^* C–H orbital is the critical factor in **5a**, in good agreement with the result that the reactivity was changed by the electronic nature of the aromatic ring of the phenethyl moiety (Table 1).

Taking this into consideration, the low reactivity of **5g** could be well rationalized. For the efficient promotion of the desired [1,5]-hydride shift, the aromatic ring should be oriented vertical to the σ^* C–H orbital. The steric repulsion between the two *ortho*-methyl groups in the mesityl group and the methylene group in **5g** rendered desired conformer **C** unfavorable, lowering the reactivity.

Further investigation of the substrate scope is shown in Figure 2. Anthracene derivative **6h** was obtained in good yield (85%). It is notable that excellent chemical yield with low catalyst loading could be achieved in the case of **6i** (96% with 0.5 mol %). This was mainly due to the cooperative electronic assistance from the π orbital of the benzene ring and the σ^* C–H orbital of the adjacent methyl group (hyperconjugation). Diastereoselective reaction was also attainable, giving desired tetracycle **6j** in good yield with fairly high selectivity (81%, *cis/trans* = 6.4/1).^{12,13} This stereoselective transformation underscores the synthetic potential of the present methodology.

In summary, an Sc(OTf)₃-catalyzed internal redox process from simple phenethyl derivatives was achieved. We have found that the reactivity was strongly affected by the electronic nature of the aromatic group adjacent to the C–H bond, and the desired 3-aryltetralins were obtained in excellent chemical yields with low catalyst loadings (as low as 0.5 mol %). It is worthy of note

that the desired tetralines were obtained in good to excellent chemical yields even with low-reactive substrates (Ar = *p*-chlorophenyl or mesityl). The results described here offer important information for rationalizing the reactivity of the benzylic hydride shift/cyclization sequence.

References and Notes

- For recent reviews on C–H activation, see: a) F. Kakiuchi, N. Chatani, *Adv. Synth. Catal.* **2003**, *345*, 1077. b) H. M. L. Davies, *Angew. Chem., Int. Ed.* **2006**, *45*, 6422. c) K. Godula, D. Sames, *Science* **2006**, *312*, 67. d) R. G. Bergman, *Nature* **2007**, *446*, 391. e) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, *107*, 174. f) H. M. L. Davies, J. R. Manning, *Nature* **2008**, *451*, 417. g) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem., Int. Ed.* **2009**, *48*, 5094. h) R. Jazzar, J. Hitce, A. Renaudat, J. Sofack-Kreutzer, O. Baudoin, *Chem.—Eur. J.* **2010**, *16*, 2654. i) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147. j) H. M. L. Davies, J. Du Bois, J.-Q. Yu, *Chem. Soc. Rev.* **2011**, *40*, 1855.
- For selected recent references, see: a) S. J. Pastine, K. M. McQuaid, D. Sames, *J. Am. Chem. Soc.* **2005**, *127*, 12180. b) S. J. Pastine, D. Sames, *Org. Lett.* **2005**, *7*, 5429. c) M. Tobisu, N. Chatani, *Angew. Chem., Int. Ed.* **2006**, *45*, 1683. d) J. Barluenga, M. Fañanás-Mastral, F. Aznar, C. Valdés, *Angew. Chem., Int. Ed.* **2008**, *47*, 6594. e) K. M. McQuaid, D. Sames, *J. Am. Chem. Soc.* **2009**, *131*, 402. f) D. Shikanai, H. Murase, T. Hata, H. Urabe, *J. Am. Chem. Soc.* **2009**, *131*, 3166. g) P. A. Vadola, D. Sames, *J. Am. Chem. Soc.* **2009**, *131*, 16525. h) I. D. Jurberg, Y. Odabachian, F. Gagosz, *J. Am. Chem. Soc.* **2010**, *132*, 3543. i) G. Zhou, J. Zhang, *Chem. Commun.* **2010**, *46*, 6593. j) M. C. Haibach, I. Deb, C. Kanta De, D. Seidel, *J. Am. Chem. Soc.* **2011**, *133*, 2100. k) M. Alajarin, M. Marin-Luna, A. Vidal, *Adv. Synth. Catal.* **2011**, *353*, 557. l) J. Barluenga, M. Fañanás-Mastral, A. Fernández, F. Aznar, *Eur. J. Org. Chem.* **2011**, *47*, 1961.
- These types of reactions have been classified under the term “*tert*-amino effect.” For reviews, see: a) O. Meth-Cohn, H. Suschitzky, *Adv. Heterocycl. Chem.* **1972**, *14*, 211. b) W. Verboom, D. N. Reinhoudt, *Recl. Trav. Chim. Pays-Bas* **1990**, *109*, 311. c) O. Meth-Cohn, *Adv. Heterocycl. Chem.* **1996**, *65*, 1. d) J. M. Quintela, *Recent Research Developments in Organic Chemistry*, **2003**, Vol. 7, p. 259. e) P. Mátyus, O. Éliás, P. Tapolsányi, Á. Polonka-Bálint, B. Halász-Dajka, *Synthesis* **2006**, 2625.
- a) D. N. Reinhoudt, G. W. Visser, W. Verboom, P. H. Benders, M. L. M. Pennings, *J. Am. Chem. Soc.* **1983**, *105*, 4775. b) W. Verboom, D. N. Reinhoudt, R. Visser, S. Harkema, *J. Org. Chem.* **1984**, *49*, 269. c) W. H. N. Nijhuis, W. Verboom, D. N. Reinhoudt, S. Harkema, *J. Am. Chem. Soc.* **1987**, *109*, 3136. d) W. H. N. Nijhuis, W. Verboom, D. N. Reinhoudt, *Synthesis* **1987**, 641. e) L. C. Groenen, W. Verboom, W. H. N. Nijhuis, D. N. Reinhoudt, G. J. Van Hummel, D. Feil, *Tetrahedron* **1988**, *44*, 4637. f) W. H. N. Nijhuis, W. Verboom, A. Abu El-Fadl, S. Harkema, D. N. Reinhoudt, *J. Org. Chem.* **1989**, *54*, 199. g) W. H. N. Nijhuis, W. Verboom, A. Abu El-Fadl, G. J. Van Hummel, D. N. Reinhoudt, *J. Org. Chem.* **1989**, *54*, 209. h) B. De Boeck, S. Jiang, Z. Janousek, H. G. Viehe, *Tetrahedron* **1994**, *50*, 7075. i) B. De Boeck, Z. Janousek, H. G. Viehe, *Tetrahedron* **1995**, *51*, 13239. j) V. Ojea, I. Muínelo, J. M. Quintela, *Tetrahedron* **1998**, *54*, 927. k) N. Kaval, B. Halász-Dajka, G. Vo-Thanh, W. Dehaen, J. Van der Eycken, P. Mátyus, A. Loupy, E. Van der Eycken, *Tetrahedron* **2005**, *61*, 9052. l) C. Zhang, C. Kanta De, R. Mal, D. Seidel, *J. Am. Chem. Soc.* **2008**, *130*, 416. m) X. Che, L. Zheng, Q. Dang, X. Bai, *Synlett* **2008**, 2373. n) Á. Polonka-Bálint, C. Saraceno, K. Ludányi, A. Bényei, P. Mátyus, *Synlett* **2008**, 2846. o) S. Murarka, C. Zhang, M. D. Konieczynska, D. Seidel, *Org. Lett.* **2009**, *11*, 129. p) J. C. Ruble, A. R. Hurd, T. A. Johnson, D. A. Sherry, M. R. Barbachyn, P. L. Toogood, G. L. Bundy, D. R. Graber, G. M. Kamilar, *J. Am. Chem. Soc.* **2009**, *131*, 3991. q) K. M. McQuaid, J. Z. Long, D. Sames, *Org. Lett.* **2009**, *11*, 2972. r) Á. A. Földi, K. Ludányi, A. C. Bényei, P. Mátyus, *Synlett* **2010**, 2109. s) P. Dunkel, G. Túrós, A. Bényei, K. Ludányi, P. Mátyus, *Tetrahedron* **2010**, *66*, 2331.
- Examples of asymmetric version of internal redox reactions: a) S. Murarka, I. Deb, C. Zhang, D. Seidel, *J. Am. Chem. Soc.* **2009**, *131*, 13226. b) Y. K. Kang, S. M. Kim, D. Y. Kim, *J. Am. Chem. Soc.* **2010**, *132*, 11847. c) W. Cao, X. Liu, W. Wang, L. Lin, X. Feng, *Org. Lett.* **2011**, *13*, 600. d) G. Zhou, F. Liu, J. Zhang, *Chem.—Eur. J.* **2011**, *17*, 3101.
- For other types of internal redox reactions, see: a) N. K. Pahadi, M. Paley, R. Jana, S. R. Waetzig, J. A. Tunge, *J. Am. Chem. Soc.* **2009**, *131*, 16626. b) C. Zhang, D. Seidel, *J. Am. Chem. Soc.* **2010**, *132*, 1798. c) I. Deb, D. Seidel, *Tetrahedron Lett.* **2010**, *51*, 2945. d) C. Zhang, D. Das, D. Seidel, *Chem. Sci.* **2011**, *2*, 233.
- a) M. Tobisu, H. Nakai, N. Chatani, *J. Org. Chem.* **2009**, *74*, 5471. b) S. Yang, Z. Li, X. Jian, C. He, *Angew. Chem., Int. Ed.* **2009**, *48*, 3999. Fillion's group found that the *p*-methoxyphenyl group is important for the benzylic hydride shift/ring closure for the formation of tetralin derivatives. See: c) S. J. Mahoney, D. T. Moon, J. Hollinger, E. Fillion, *Tetrahedron Lett.* **2009**, *50*, 4706. For hydride shifts from di- and/or triarylmethane, see: d) G. B. Bajracharya, N. K. Pahadi, I. D. Gridnev, Y. Yamamoto, *J. Org. Chem.* **2006**, *71*, 6204. e) M. Alajarin, B. Bonillo, M.-M. Ortin, P. Sanchez-Andrada, A. Vidal, R.-A. Orenes, *Org. Biomol. Chem.* **2010**, *8*, 4690.
- a) K. Mori, S. Sueoka, T. Akiyama, *J. Am. Chem. Soc.* **2011**, *133*, 2424. For another internal redox process developed by our group: see, b) K. Mori, Y. Ohshima, K. Ehara, T. Akiyama, *Chem. Lett.* **2009**, *38*, 524. c) K. Mori, T. Kawasaki, S. Sueoka, T. Akiyama, *Org. Lett.* **2010**, *12*, 1732. For an asymmetric version of internal redox reaction catalyzed by chiral phosphoric acid: see, d) K. Mori, K. Ehara, K. Kurihara, T. Akiyama, *J. Am. Chem. Soc.* **2011**, *133*, 6166.
- Supporting Information is available electronically on the CSJ-Journal Web site: <http://www.csj.jp/journals/chem-lett/>.
- Only the combination of barbituric acid as the electrophilic portion and Sc(OTf)₃ as the catalyst afforded the desired products. For example, treatment of dimethyl benzylidenemalonate with 30 mol % various acids (TfOH, SnCl₄, TiCl₄, BF₃·OEt₂, Yb(OTf)₃, and Sc(OTf)₃) in refluxing ClCH₂CH₂Cl led to only the recovery of the starting material. When benzylidenemeldrumate was used as the substrate, the decomposition of the starting material was observed in most cases.
- The hyperconjugative interaction between the lone pair electrons of heteroatom and σ^{*}C–H orbital weakened and polarized the adjacent C–H bond: P. Brunet, J. D. Wuest, *J. Org. Chem.* **1996**, *61*, 2020, and references cited therein.
- The diastereomer ratio was determined by comparing the integration value of each C-9 hydrogen in ¹H NMR (CDCl₃). The relative stereochemistry of the major isomer was unambiguously established by single-crystal X-ray analysis. CCDC-844878 contains the supplementary crystallographic data of *cis*-**6j**. These data can be obtained free of charge from The Cambridge Crystallographic Data Center via <http://www.ccdc.cam.ac.uk/data-request>.
- Due to the fixed configuration of the C-10 carbon, the nucleophilic attack mainly occurred from the same face of the aromatic ring in the cationic intermediate to afford *cis*-**6j**.