Nickel-Catalyzed Coupling of Arynes, Alkenes, and Boronic Acids: Dual Role of the Boronic Acid**

Thiruvellore Thatai Jayanth and Chien-Hong Cheng*

Nickel-catalyzed domino reactions that involve the coupling of two π components with main-group organometallic reagents or metal hydrides have emerged as highly useful methods for assembling complex molecular structures from simple starting materials.^[1] This class of reactions has evolved significantly in scope as well as substrate choice. Most commonly employed π systems include alkynes, alkenes, dienes, allenes, aldehydes, imines, and allylic halides. The organometallic reagents employed are usually based on zinc, aluminum, zirconium, tin, boron, and silanes.^[2] The utility of this concept has been demonstrated well in organic synthesis.^[3] A similar reaction sequence employing arynes as a π component would be highly interesting and useful, considering that arynes are generally very reactive and generated in situ and that the reaction would lead to the formation of two different C-C bonds in ortho positions of a benzene ring.^[4] A mild method for the preparation of arynes in situ at moderate temperatures from commercially available o-silyl aryl triflates $\mathbf{1}^{[5]}$ has rekindled the interest, especially in the past few years, in employing arynes as substrates in organic synthesis.^[6] As part of our ongoing efforts in employing arynes as useful substrates in organic synthesis,^[7] we report an unprecedented nickel-catalyzed three-component coupling of arynes, enones, and organoboronic acids. Thus, when a solution of benzyne precursor 1a (2-(trimethylsilyl)phenyl triflate), ethyl vinyl ketone (2a), and trans-2-phenylvinylboronic acid (3a) in the presence of $[Ni(cod)_2]$ (cod = 1,5cyclooctadiene; 10 mol%), PPh₃ (20 mol%), and CsF in CH₃CN was stirred for 8 h at 40 °C, coupling product 4a was obtained in 92% yield (Table 1, entry 1).^[8] This process is the first example involving arynes and is also one of the first to utilize organoboronic acids in nickel-catalyzed multicomponent coupling reactions.^[9]

A variety of arynes are compatible with this threecomponent coupling reaction (Table 1). The aryne precursor **1b** with two methyl groups on the phenyl ring furnished product **4b** in 93% yield (Table 1, entry 2) and indanyl derivative **1c** afforded **4c** in 84% yield (Table 1, entry 3). Aryne precursors that bear other electron-donating substitu-

| [*] T. T. Jayanth, Prof. Dr. CH. Cheng |
|----------------------------------------|
| Department of Chemistry |
| National Tsing Hua University |
| Hsinchu 30013 (Taiwan) |
| Fax: (+886) 3-572-4698 |
| E-mail: chcheng@mx.nthu.edu.tw |
| |

^[**] We thank the National Science Council of the Republic of China (NSC 95-2119M-007-005) for support of this research.

Supporting information for this article is available on the WWW under http://www.angewandte.org or from the author.

ents such as 3,4-dimethoxy and 1,3-benzodioxole derivatives **1d** and **1e**, respectively, gave products in excellent yields (Table 1, entries 4 and 5). Furthermore, the difluoro substituted benzyne precursor **1f** gave the corresponding three-





[a] Reaction conditions: aryne precursor 1 (0.30 mmol), enone 2 (0.45 mmol), boronic acid 3 (0.30 mmol), $[Ni(cod)_2]$ (0.030 mmol), PPh₃ (0.060 mmol), and CsF (0.90 mmol) in CH₃CN (1 mL) at 40 °C for 8 h. [b] Yields of isolated products. [c] The structure of the major isomer is represented. [d] Mixture of regioisomers (1:1).

© 2007 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



Communications

component product **4f** (Table 1, entry 6). Very interestingly, the aryne precursor **1g** with a methoxy group at the *ortho* position afforded regioisomers in 85:15 ratio in which the predominant product was that with the vinyl group from the boronic acid attached *ortho* to the methoxy group (Table 1, entry 7). In the case of benzyne precursor **1h**, the reaction gave a 1:1 mixture of regioisomers (Table 1, entry 8), as expected for the reaction of the unsymmetrical 4-methylben-zyne substrate.

This three-component assembly works well with both acyclic and cyclic enones (Table 2). Propyl vinyl ketone (**2b**) worked smoothly under the reaction conditions (Table 2, entry 1). Cyclic enones (five-, six-, and seven-membered rings) reacted equally efficiently to yield the expected products in very good yields (Table 2, entries 2–4). Acrylates

were also found to be compatible in the present reaction (Table 2, entries 5 and 6). When acrylonitrile (**2h**) was employed the three-component coupling product **4o** was obtained in 71 % yield (Table 2, entry 7). When boronic acid **3a** was replaced by its pinacol ester the reaction did not proceed to give the expected product, and boroxine ((4-MeOC₆H₅)₃B₃O₃) gave trace amounts of product. The reaction also failed with other soft organometallic reagents based on tin (PhC=CSnBu₃) and silicon (PhSiMe₃). The boronic acid acting as a proton source is the likely reason for this observation (see below). The boronic acid was found to be crucial for the success of this reaction. Alkenyl boronic acids worked particularly well. Thus hexenylboronic acid afforded the three-component product in excellent yield (Table 2, entry 8). Of the other organoboronic acids tested, *p*-, *o*-, and

Table 2: Three-component coupling of benzyne with alkenes 2 and organoboronic acids 3.^[a]



[a] Reaction conditions: aryne precursor 1 (0.30 mmol), enone 2 (0.45 mmol), boronic acid 3 (0.30 mmol), $[Ni(cod)_2]$ (0.030 mmol), PPh₃ (0.060 mmol), and CsF (0.90 mmol) in CH₃CN (1 mL) at 40 °C for 8 h. [b] Yields of isolated products. [c] P(p-C₆H₄F)₃ was employed instead of PPh₃.

5922 www.angewandte.org

© 2007 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

m-methoxybenzeneboronic acids (3c-e) gave the corresponding products in 67–84 % yield (Table 2, entries 9–11). Benzeneboronic acid (3f) provided product 4t in 60 % yield (Table 2, entry 12). In the case of boronic acids 3e and 3f, phosphine ligand P(*p*-C₆H₄F)₃ was employed, as the reaction did not proceed when PPh₃ was used as the ligand.

A plausible mechanistic rationale for the present catalytic reaction is shown in Scheme 1. The catalysis is likely initiated by the reaction of Ni^0 with the enone and the aryne to form a nickelacycle 5. Protonation of the α carbon atom of the



Scheme 1. Proposed mechanism for the three-component coupling reaction.

ketone moiety by the boronic acid and transmetalation of the organic group onto the nickel center leads to **6**. Subsequent reductive elimination gives the desired coupling product and regenerates the Ni⁰ catalyst. To verify this pathway, a deuterated boronic acid, *p*-OMeC₆H₄B(OD)₂, was carefully prepared and employed. As expected, deuterium was incorporated at the α carbon atoms of the ketone product (Scheme 2).^[10] Protonation of **5** by the organoboronic acid likely enhances electron density at the boron center and thus the nucleophilicity of the carbon nucleophile to facilitate the transmetalation and catalysis. Thus, the organoboronic acid plays a dual role in the present catalytic process: it acts both as a proton source and as a carbon nucleophile.^[11] An alternative pathway for the formation of **6** from **5** via a boron enolate **7** and subsequent protonation cannot be completely ruled out.



Scheme 2. Deuterium-labeling experiment.

Angew. Chem. Int. Ed. 2007, 46, 5921-5924

The regioselectivity observed in the reaction of 1g (Table 1, entry 7) can be rationalized by the formation of *ortho*-metalated complex 5g on the basis of the coordination of the methoxy group to nickel.^[12]

A seven-membered ring nickel *O*enolate generated by the oxidative cyclization of an enal and an alkyne with Ni^0 species containing a *Z* carbon– carbon double bond is documented as the intermediate for protonation during the coupling reaction.^[13] However, in



the present case, a seven-membered ring nickel enolate intermediate from 5 can be ruled out, considering that the seven-membered ring intermediate generated by a cyclic enone such as 2-cyclopenten-1-one and an aryne with Ni⁰ would contain a strained *E* double bond in the seven-membered ring.^[14]

In summary, we have demonstrated a nickel-catalyzed domino reaction of arynes, enones, and boronic acids. This reaction represents the first example of the use of arynes in nickel-catalyzed coupling reactions. The organoboronic acid acts both as a proton source and as a carbon nucleophile.

Experimental Section

General Procedure: A round-bottomed side-arm flask containing $[Ni(cod)_2]$ (0.030 mmol), PPh₃ (0.060 mmol), CsF (0.90 mmol), and boronic acid **3** (0.30 mmol) was evacuated and purged with nitrogen gas. CH₃CN (1.0 mL), alkene **2** (0.450 mmol), and aryne precursor **1** (0.30 mmol) were then added to the flask through syringes. The reaction mixture was allowed to stir at 40 °C for 8 h. At the end of the reaction, the reaction mixture was diluted with CH₂Cl₂ and filtered through celite and silica gel. The filtrate was concentrated, and the residue was purified through a silica-gel column with ethyl acetate/ hexane as eluent to give pure product **4**.

Received: March 10, 2007 Published online: June 20, 2007

Keywords: arynes · boronic acids · enones · multicomponent reactions · nickel

- For reviews, see: a) J. Montgomery, Angew. Chem. 2004, 116, 3980; Angew. Chem. Int. Ed. 2004, 43, 3890; b) S. Ikeda, Acc. Chem. Res. 2000, 33, 511; c) J. Montgomery, Acc. Chem. Res. 2000, 33, 467; d) J. Montgomery, K. K. D. Amarasinghe, S. K. Chowdhury, E. Oblinger, J. Seo, A. V. Savchenko, Pure Appl. Chem. 2002, 74, 129; e) K. M. Miller, C. Molinaro, T. F. Jamison, Tetrahedron: Asymmetry 2003, 14, 3619.
- [2] For very recent examples, see: a) C.-Y. Ho, T. F. Jamison, Angew. Chem. 2007, 119, 796; Angew. Chem. Int. Ed. 2007, 46, 782; b) S.-S. Ng, C.-Y. Ho, T. F. Jamison, J. Am. Chem. Soc. 2006, 128, 11513; c) M. Song, J. Montgomery, Tetrahedron 2005, 61, 11440; d) S.-S. Ng, T. F. Jamison, Tetrahedron 2005, 61, 11405; e) M. Takimoto, Y. Kajima, Y. Sato, M. Mori, J. Org. Chem. 2005, 70, 8605; f) S.-S. Ng, T. F. Jamison, J. Am. Chem. Soc. 2005, 127, 7320; g) H. P. Hratchian, S. K. Chowdhury, V. M. Gutierrez-Garcia, K. K. D. Amarasinghe, M. J. Heeg, H. B. Schlegel, J. Montgomery, Organometallics 2004, 23, 4636; h) M. Kimura, A. Ezoe, M. Mori, Y. Tamaru, J. Am. Chem. Soc. 2005, 127, 201; i) E. Shirakawa, Y. Yamamoto, Y. Nakao, S. Oda, T. Tsuchimoto, T. Hiyama, Angew. Chem. 2004, 116, 3530; Angew. Chem. Int. Ed.

© 2007 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Communications

2004, *43*, 3448; j) G. M. Mahandru, A. R. L. Skauge, S. K. Chowdhury, K. K. D. Amarasinghe, M. J. Heeg, J. Montgomery, *J. Am. Chem. Soc.* **2003**, *125*, 13481; k) C. Molinaro, T. F. Jamison, *J. Am. Chem. Soc.* **2003**, *125*, 8076; for earlier work, see reference [1].

- [3] For examples of the application of this concept in total synthesis, see: a) E. A. Colby, K. C. O'Brien, T. F. Jamison, J. Am. Chem. Soc. 2005, 127, 4297; b) J. Chan, T. F. Jamison, J. Am. Chem. Soc. 2004, 126, 10682; c) Y. Ni, K. K. D. Amarasinghe, B. Ksebati, J. Montgomery, Org. Lett. 2003, 5, 3771; d) K. K. D. Amarasinghe, J. Montgomery, J. Am. Chem. Soc. 2002, 124, 9366; e) E. A. Colby, K. C. O'Brien, T. F. Jamison, J. Am. Chem. Soc. 2004, 126, 998; f) J. Chan, T. F. Jamison, J. Am. Chem. Soc. 2004, 126, 11514.
- [4] a) U. K. Tambar, B. M. Stoltz, J. Am. Chem. Soc. 2005, 127, 5340;
 b) M. Jeganmohan, C.-H. Cheng, Org. Lett. 2004, 6, 2821; c) M. Jeganmohan, C.-H. Cheng, Synthesis 2005, 1693; d) T. T. Jayanth, M. Jeganmohan, C.-H. Cheng, Org. Lett. 2005, 7, 2921; e) J. L. Henderson, A. S. Edwards, M. F. Greaney, J. Am. Chem. Soc. 2006, 128, 7426.
- [5] Y. Himeshima, T. Sonoda, H. Kobayashi, Chem. Lett. 1983, 1211.
- [6] For reviews, see: a) E. Guitián, D. Perez, D. Peña, Top. Organomet. Chem. 2005, 14, 109; b) D. Peña, D. Perez, E. Guitián, Angew. Chem. 2006, 118, 3659; Angew. Chem. Int. Ed. 2006, 45, 3579; c) E. Guitian, D. Perez, D. Peña, Top. Organomet. Chem. 2005, 14, 109; for recent application of this aryne precursors 1 in total synthesis, see: d) U. K. Tambar, D. C. Ebner, B. M. Stoltz, J. Am. Chem. Soc. 2006, 128, 11752; e) Y. Sato, T. Tamura, M. Mori, Angew. Chem. 2004, 116, 2490; Angew. Chem. Int. Ed. 2004, 43, 2436.
- [7] a) T. T. Jayanth, C.-H. Cheng, *Chem. Commun.* 2006, 894;
 b) T. T. Jayanth, M. Jeganmohan, M.-J. Cheng, S.-Y. Chu, C.-H. Cheng, *J. Am. Chem. Soc.* 2006, 128, 2232; c) S. Bhuvaneswari, M. Jeganmohan, C.-H. Cheng, *Org. Lett.* 2006, *8*, 5581; d) T. T.

Jayanth, M. Jeganmohan, C.-H. Cheng, *J. Org. Chem.* **2004**, *69*, 8445; e) J.-C. Hsieh, D. K. Rayabarapu, C.-H. Cheng, *Chem. Commun.* **2004**, 532; f) J.-C. Hsieh, D. K. Rayabarapu, C.-H. Cheng, *Chem. Commun.* **2004**, 2459.

- [8] For a recent report on the metal-catalyzed cyclotrimerization of alkenes and arynes, see: I. Quintana, A. J. Boersma, D. Peña, D. Perez, E. Guitián, Org. Lett. 2006, 8, 3347.
- [9] For an earlier report of boronic acids in nickel-catalyzed multicomponent reactions, see: S. J. Patel, T. F. Jamison, *Angew. Chem.* 2003, 115, 1402; *Angew. Chem. Int. Ed.* 2003, 42, 1364.
- [10] Even though, mechanistically, deuterium-labeling is expected at only one of the α carbons atoms, we obtained, approximately, an equal mixture of [D]4u and [D]4u'. This could be due to the scrambling of the deuterium between the α carbon atoms of the ketone catalyzed by CsF. Organoboronic acids acting as a proton source is proposed in literature; see: C. H. Oh, H. H. Jung, K. S. Kim, N. Kim, Angew. Chem. 2003, 115, 829; Angew. Chem. Int. Ed. 2003, 42, 805.
- [11] No deuteration was observed, either, when the reaction (Table 1, entry 1) was performed in CD_3CN or when quenched with D_2O at the end of the reaction.
- [12] For a similar methoxy-directing effect in catalytic reactions, see: a) N. Chatani, A. Kamitani, M. Oshita, Y. Fukumoto, S. Murai, J. Am. Chem. Soc. 2001, 123, 12686; b) M. Sonoda, F. Kakiuchi, N. Chatani, S. Murai, Bull. Chem. Soc. Jpn. 1997, 70, 3117; c) R. Takeuchi, H. Yasue, J. Org. Chem. 1993, 58, 5386.
- [13] a) S. K. Chowdhury, K. K. D. Amarasinghe, M. J. Heeg, J. Montgomery, J. Am. Chem. Soc. 2000, 122, 6775; b) G. M. Mahandru, A. R. L. Skauge, S. K. Chowdhury, K. K. D. Amarasinghe, M. J. Heeg, J. Montgomery, J. Am. Chem. Soc. 2003, 125, 13481.
- [14] S. Ikeda, R. Sanuki, H. Miyachi, H. Miyashita, M. Taniguchi, K. Odashima, J. Am. Chem. Soc. 2004, 126, 10331.