

View Article Online View Journal

# **RSC Advances**

This article can be cited before page numbers have been issued, to do this please use: A. Hajra, R. Hashim, M. Rahman, I. Ling and N. Abdullah, *RSC Adv.*, 2014, DOI: 10.1039/C4RA16374E.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This Accepted Manuscript will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/advances

Published on 22 December 2014. Downloaded by McMaster University on 22/12/2014 20:38:14.

### **Graphical Abstract**

# Organocatalysis by *p*-sulfonic acid calix[4]arene: a convenient and efficient route to 2,3-dihydroquinazolin-4(1*H*)-ones in water

Matiur Rahman, Irene Ling, Norbani Abdullah, Rauzah Hashim\*and Alakananda Hajra\*



## **ARTICLE TYPE**

# Organocatalysis by *p*-sulfonic acid calix[4]arene: a convenient and efficient route to 2,3-dihydroquinazolin-4(1*H*)-ones in water

Matiur Rahman,<sup>a</sup> Irene Ling,<sup>a</sup> Norbani Abdullah,<sup>a</sup> Rauzah Hashim<sup>a</sup>\*and Alakananda Hajra<sup>b</sup>\*

Received (in XXX, XXX) Xth XXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX 5 DOI: 10.1039/b000000x

An efficient and eco-friendly method is reported for the synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones from the direct cyclocondensation of anthranilamide with aldehydes using *p*-sulfonic acid calix[4]arene (*p*-SAC) as a recyclable <sup>10</sup> organocatalyst in excellent yields in water at room temperature. The catalyst was reusable without significant loss of catalytic efficiency. Operational simplicity, the compatibility with various functional groups, non-chromatographic purification technique, high yields and mild <sup>15</sup> reaction conditions are the notable advantages of this procedure. Large scale reaction demonstrated the practical applicability of this methodology.

#### Introduction

Published on 22 December 2014. Downloaded by McMaster University on 22/12/2014 20:38:14.

Catalysis has played a crucial role in the success of the chemistry <sup>20</sup> in the twentieth century.<sup>1</sup> Nowadays organocatalysis is one of the hot research topics in advanced organic chemistry. In the past decade, organocatalysis has opened a new window for carrying out organic transformations and has become a powerful tool in the synthesis of biologically active and structurally complex <sup>25</sup> compounds.<sup>2</sup> Compared with transition-metal catalysts, the cost and toxicity of organocatalysts are low, thus making organocatalysts beneficial for the production of pharmaceutical intermediates.<sup>2,3</sup> Moreover, organocatalysts are tolerant of water and air, are usually easy to use and finally avoidance of expensive <sup>30</sup> metal reagents or catalysts.<sup>4</sup>

The development of eco-friendly methodologies that are environmentally benign and waste-free, as well as being able to produce the desired products in high purity, have received much attention in recent years because of the increasing tendency of the

- <sup>35</sup> chemical industry towards greener processes.<sup>5</sup> The major drive towards this idea is the replacement of volatile organic solvents by green solvents,<sup>6</sup> as organic solvents are the major contributors to environmental pollution. In this view, water is the most ideal solvent<sup>7</sup> and the use of water in organic reactions as solvent has
- <sup>40</sup> received much attention.<sup>8</sup> Water is a safe, harmless and environmentally benign solvent in comparison with a large number of harmful organic solvents; the unique physicochemical properties of water can even accelerate some reactions.<sup>9</sup> However, the poor solubility of most organic compounds in water
- <sup>45</sup> often makes an unfavorable impact on water mediated organic synthesis. Therefore, the development of water-tolerant catalysts that allow organic reactions to be carried out in aqueous media is

a challenging task in the field of green catalysis. Aqueous biphasic reaction systems using water-soluble catalysts have the <sup>50</sup> practical advantages that the catalysts can be reused after simple decantation or extraction of the water-insoluble products and the catalysts show high catalytic activity in water.<sup>10</sup> The thought of environmental factor (E-factor) and atom economy have gradually become incorporated into conventional organic <sup>55</sup> synthesis in both industry and academia. Solvents are the main reason for an insufficient E-factor, especially in synthesis of fine chemicals and pharmaceutical industries.<sup>11</sup>

The calixarenes are a class of cyclooligomers formed via condensation between para substituted phenols and 60 formaldehyde.<sup>12</sup> Calixarenes are good receptors for the complexation with various kinds of guests such as anions, cations and neutral organic/inorganic molecules.<sup>13</sup> Calixarenes have been widely used in sensors, enzyme-mimics, ion carriers, solid-phase support materials, ion selective electrodes, drug-delivery agents 65 etc.<sup>14</sup> The discovery of water-soluble calixarenes as catalyst in organic reaction has attracted the attention of researchers with the objective of faster reaction and use of lower amount of catalyst.<sup>15</sup> In this sense, water-soluble calix[n]arenes could be used as surfactant-type Brønsted acid catalyst to facilitate the reactions 70 through the formation of a host-guest complex, with a nucleophile component in the organic-aqueous interfacial layer.10 So the use of calix[n]arenes as catalyst in water is in demand from the aspect of green chemistry. However the use of calixarenes as catalyst in water is very rare.<sup>15a</sup>

<sup>75</sup> The *N*-containing heterocyclic compounds play a crucial role in the field of drug scaffolds, synthetic organic chemistry, medicinal chemistry as well as material sciences.<sup>16</sup> 2,3-Dihydroquinazolinone derivatives act as important intermediates in the synthesis of drug molecules, and natural products.<sup>17</sup> These <sup>80</sup> also exhibit various activities such as antitumor, anti-cancer, antibiotic, antidefibrillatory, antipyretic, analgesic, antihypertonic, diuretic, antihistamine, antidepressant, and vasodilating agents<sup>18</sup> (Fig 1).



Fig. 1 Biologically important quinazolinone derivatives.

85

Published on 22 December 2014. Downloaded by McMaster University on 22/12/2014 20:38:14.

In view of their significant uses, green methodologies for the synthesis of 2,3-dihydroquinazolin-4(1H)-ones should be benefited in the synthetic and medicinal chemistry. Although numerous protocols have been developed for the synthesis of 2,3-<sup>5</sup> dihydroquinazolinones,<sup>19</sup> regardless their efficiency and reliability, most of these methods suffer from one or more of these disadvantages, such as the use of hazardous organic solvents, requires higher temperature, low yields, strongly acidic conditions, expensive moisture sensitive catalysts, and tedious 10 work-up procedure. Therefore, it is important to develop economically and environmentally more viable procedure for the synthesis of 2,3-dihydroquinazolin-4(1H)-ones. Very recently, we have developed an environmentally benign greener strategy using recyclable nano indium oxide as catalyst for the one-pot synthesis 15 of 2,3-dihydroquinazolin-4(1H)-ones by a three-component condensation of isatoic anhydride with primary amines or ammonium salts and aromatic aldehydes in ethanol.<sup>190</sup> As a part of our ongoing research for the synthesis of heterocycles through

a greener way,  $^{19o,20}$  herein we report a simple and practical <sup>20</sup> metal-free method for the synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones by the cyclocondensation of anthranilamide with aldehydes using *p*-sulfonic acid calix[4]arene (*p*-SAC) as a recyclable organocatalyst in water at room temperature (Scheme 1).



Scheme 1 Synthesis of 2,3-dihydroquinazolin-4(1H)-ones.

We started our study with preparing *p*-sulfonic acid calix[*n*]arenes in our laboratory according to the literature <sup>30</sup> procedure.<sup>21</sup> Next the catalytic activity of these organocatalyst were investigated for the synthesis of 2,3-dihydroquinazolin-4(1H)-ones. In order to find out the optimum reaction conditions, we have selected the reaction of anthranilamide **1** and 4methylbenzaldehyde **2a** as the model reaction (Table 1). To our <sup>35</sup> delight, the desired product **3a** was obtained in 92% yield in presence of 1 mol% *p*-sulfonic acid calix[4]arene (*p*-SAC) in water at room temperature for 20 min (Table 1, entry 1). No further improvement was noticed by increasing the reaction time (Table 1, entry 2). Increasing the amount of catalyst (5 mol%) did <sup>40</sup> not improve the yield noticeably (Table 1, entry 3,) whereas decreasing the amount of catalyst (0.5 mol%) decreased the yield

- (Table 1, entry 4). To clarify the role of the catalyst (p-SAC) in this process, the same reaction was carried out without p-SAC (Table 1, entry 5). However the reaction did not proceed at all
- <sup>45</sup> without the catalyst after 20 min of reaction. Water appeared to be the best choice as solvent among the common solvents like MeOH, MeCN, toluene, DMF, 1,2-DCE (Table 1, entries 6-10).

The reaction did not proceed well in absence of any solvent (Table 1, entry 11). The chain length of calixarenes also played a <sup>50</sup> significant role. By increasing the chain length of 4 to 6 (*n* = 1 to 3) the yield of the desired product decreased noticeably (Table 1, entry 12). To explore the role of *p*-sulfonic acid calix[4]arene (*p*-SAC) we used *p*-hydroxy benzenesulfonic acid (*p*-HSA) as a catalyst. It was observed that *p*-HSA was less efficient than *p*-<sup>55</sup> SAC (Table 1, entry 13). This indicates that the sulfonyl and phenolic groups in calixarene moiety are not solely responsible for fast and efficient reaction. On the other hand, *p*-TSA was not also effective for this conversion (Table 1, entry 14). Phenol was less efficient than *p*-SAC (Table 1, entry 15). Other Brønsted acid <sup>60</sup> catalyst like *p*-dodecylbenzenesulfonic acid (DBSA, Kobayashi's catalyst<sup>2m</sup>) was also tested, but no improvement of the yield was

observed (Table 1, entries 16). Thus, optimal reaction conditions were obtained using anthranilamide (1, 1 mmol), 4methylbenzaldehyde (2a, 1 mmol) in presence of 1 mol% of p-<sup>65</sup> sulfonic acid calix[4]arene (p-SAC) in water (1 mL) at room temperature for 20 min (Table 1, entry 1).

Table 1 Optimization of the reaction conditions<sup>a</sup>

$\bigcup_{NH_2}^{O} + \bigcup_{CH_3}^{CHO} \xrightarrow{Catalyst}_{Solvent, rt} \qquad \bigcup_{H_2}^{O} + \bigcup_{CH_3}^{O} \xrightarrow{Catalyst}_{CH_3}$							
Entry	Catalyst (mol%)	Solvent (1 mL)	Time (min)	Yields $(\%)^b$			
1	<i>p</i> -SAC (1)	$H_2O$	20	92			
2	<i>p</i> -SAC (1)	$H_2O$	60	92			
3	<i>p</i> -SAC (5)	$H_2O$	20	93			
4	<i>p</i> -SAC (0.5)	H <sub>2</sub> O	20	84			
5	-	$H_2O$	20	n. r. <sup>c</sup>			
6	<i>p</i> -SAC (1)	MeOH	20	80			
7	<i>p</i> -SAC (1)	MeCN	20	72			
8	<i>p</i> -SAC (1)	Toluene	20	38			
9	<i>p</i> -SAC (1)	DMF	20	75			
10	<i>p</i> -SAC (1)	DCE	20	32			
11	<i>p</i> -SAC (1)	-	20	20			
12	<i>p</i> -Sulfonic acid calix[6]arene $[n = 3]$ (1)	H <sub>2</sub> O	20	67			
13	<i>p</i> -HSA (1)	H <sub>2</sub> O	20	60			
14	<i>p</i> -TSA (1)	H <sub>2</sub> O	20	57			
15	Phenol (1)	$H_2O$	20	18			
16	DBSA (5)	H <sub>2</sub> O	20	58			

	<sup>a</sup> Reaction conditions: 1 mmol of 1 and 1 mmol of 2a in the presence of
70	catalyst and solvent (1 mL) at room temperature. <sup>b</sup> Isolated yields. <sup>c</sup> No
	reaction. $p$ -SAC = $p$ -Sulfonic acid calix[4]arene. $p$ -HSA = $p$ -
	Hydroxybenzenesulfonic acid. $p$ -TSA = $p$ -Toluenesulfonic acid. TBAB =
	Tetrabutyl ammonium bromide. $DBSA = p$ -Dodecylbenzenesulfonic acid.

Next, we studied the scope and limitation of this reaction by employing a wide range of aldehydes (2) under optimized reaction conditions. The results are summarized in Table 2. To our delight, the corresponding dihydroquinazolin-4(1*H*)-ones (3) were obtained with good to excellent yields in all cases. Aromatic aldehydes with electron-donating as well as electron-withdrawing substituents reacted very well, affording good to excellent yields

50

of 2,3-dihydroquinazolinones. The chloro- and bromosubstituted benzaldehydes afforded the corresponding products 3d and 3e in 87% and 86% yields respectively. Substituent at the ortho position of the phenyl ring also afforded the corresponding 5 products with excellent yields (3j and 3k). Heteroaryl aldehyde like 2-thiophenecarboxaldehyde was also compatible under this reaction conditions to give the desired product (3p) without polymerization. Gratifyingly, the current methodology is also applicable for the aliphatic aldehydes. Cyclohexane 10 carboxaldehyde gave the desired product (3q) with high yield. Butvraldehvde also afforded the corresponding 2,3dihydroquinazolinone (3r) with good yield however longer reaction time is required. Moreover, ketone like acetone also afforded the desired product (3s) with moderate yield. The all 15 synthesized compounds have been characterized by spectral and analytical data. Results in Table 2 clearly show that the present protocol is indeed superior to several reported methods in terms of product yields, reaction time, avoiding the use of volatile organic solvents, and reaction temperature. Moreover, we have 20 developed a greener reaction conditions bearing lower Efactor<sup>11,22</sup> of 0.17 and 0.15 in the cases of synthesizing **3a** and **3b** 

factor<sup>11,22</sup> of 0.17 and 0.15 in the cases of synthesizing 3a and respectively (see Supporting Information).

 Table 2 Substrates scopes of the reaction<sup>a</sup>

Published on 22 December 2014. Downloaded by McMaster University on 22/12/2014 20:38:14.

$1 \qquad 2 \qquad $							
Entry	$\mathbf{R}^1$	R <sup>2</sup>	Products	Time (min)	Yields $(\%)^b$		
1	4-Me-C <sub>6</sub> H <sub>4</sub>	Н	3a	20	92, 86 <sup>c</sup>		
2	Ph	Н	3b	18	94		
3	4-MeO-C <sub>6</sub> H <sub>4</sub>	Н	3c	20	90		
4	4-Cl-C <sub>6</sub> H <sub>4</sub>	Н	3d	22	87		
5	4-Br-C <sub>6</sub> H <sub>4</sub>	Н	3e	22	86		
6	$4-NO_2-C_6H_4$	Н	3f	25	82		
7	4-MeS-C <sub>6</sub> H <sub>4</sub>	Н	3g	20	88		
8	4-CN-C <sub>6</sub> H <sub>4</sub>	Н	3h	22	86		
9	$4-OH-C_6H_4$	Н	3i	24	82		
10	$2-Cl-C_6H_4$	Н	3ј	20	90		
11	$2-OH-C_6H_4$	Н	3k	22	88		
12	4-OH-3,5- (OMe) <sub>2</sub> -C <sub>6</sub> H <sub>2</sub>	Н	31	24	84		
13	5-Br-2-OH-C <sub>6</sub> H <sub>3</sub>	Н	3m	22	85		
14	$3-NO_2-C_6H_4$	Н	3n	24	88		
15	1-Pyrenyl	Н	30	26	82		
16	2-Thiophenyl	Н	3р	22	86		
17	Cyclohexyl	Н	3q	30	82		
18	$C_3H_7$	Н	3r	40	68		
19	CH <sub>3</sub>	CH <sub>3</sub>	3s	90	64		

- <sup>25</sup> <sup>a</sup> Reaction conditions: 1 (1 mmol), 2 (1 mmol) in water (1 mL) at room temperature in presence of 1 mol% of *p*-SAC. <sup>b</sup> Isolated yields. <sup>c</sup> 1 (20 mmol), 2a (20 mmol) in water (20 mL) at room temperature in presence of 1 mol% of *p*-SAC.
- <sup>30</sup> The dihydroquinazolinones were generally precipitated from the reaction mixtures and the solid precipitation was filtered off and recrystallized from hot ethanol to obtain pure products. This methodology is also applicable on a gram-scale synthesis. We

have successfully synthesized the dihydroquinazolinone **3a** in 35 86% yield by the reaction of anthranilamide (**1**, 20 mmol) and 4methylbenzaldehyde (**2a**, 20 mmol) (Table 2, entry 1).

Finally the single crystal X-ray diffraction analysis of **3n** was carried out for further confirmation of the structure of product (Fig. 2).<sup>23</sup> In crystalline state the compound shows the 2,3-<sup>40</sup> dihydroquinazoline moiety is planar. The pyrimidine ring is in a flattened half-chair conformation. The nitro-substituted benzene ring forms dihedral angle of 85.91°, with the benzene ring of the dihydroquinazoline group. In the crystal, molecules are arranged in crisscross manner along *a*-axis, with N—O...H hydrogen <sup>45</sup> bonds linking molecules at 2.451 to 2.708 Å in the extended network. Short contacts of hydrogen bonding interaction related to carbonyl oxygen atoms with neighboring hydrogen atoms is observed with O…H distances ranging from 2.080 to 2.636 Å.



Fig. 2 The single crystal XRD structure of compound 3n.

Next, we turned our attention towards the recovery and reusability of the catalyst. For this purpose, we have chosen the reaction of anthranilamide (1) with 4-methylbenzaldehyde (2a) in presence of 1 mol% *p*-sulfonic acid calix[4]arene (*p*-SAC) in <sup>55</sup> water at room temperature as the model reaction. After completion of the reaction distilled water was added to the reaction mixture. The reaction mixture was then filtered off and the catalyst was recovered by evaporating the water. Pure product was obtained by recrystallizing the residue from hot ethanol. The <sup>60</sup> recovered catalyst was reused for a subsequent fresh batch of the reaction. The catalytic activity was checked upto fourth cycle and it showed similar (Table 3).

**Table 3** Recycling of the *p*-sulfonic acid calix[4]arene (*p*-SAC) for  ${}^{65}$  synthesizing  $3a^a$ 

Entry	Run	Yield $(\%)^b$
1	1st	92
2	2nd	92
3	3rd	90
4	4th	89
5	5th	87

<sup>*a*</sup> Carried out with 1 mmol of **1** and 1 mmol of **2a** in presence of catalyst in water (1 mL) at room temperature. <sup>*b*</sup> Isolated yields.

Based o the literature reports,  $^{19g}$  a plausible mechanistic 70 pathway to 2,3-dihydroquinazolin-4(1*H*)-ones (**3**) is illustrated in 65

75

95

100

**RSC Advances Accepted Manuscrip** 

Scheme 2. In the initial step, condensation of anthranilamide (1) with the aldehyde (2) gives imine A, which upon intramolecular cyclization afforded the final product 3. The catalyst might activate the aldehyde through the co-ordination with the oxygen s of the corresponding carbonyl group which facilitates the subsequent nucleophilic attack by the nitrogen atom on the carbonyl carbon.



Scheme 2 Plausible mechanistic pathway

#### Conclusions

10

In summary, we have demonstrated a remarkable catalytic activity of *p*-sulfonic acid calix[4]arene (*p*-SAC) as an organocatalyst for the synthesis of 2,3-dihydroquinazolin-4(1*H*)-<sup>15</sup> ones by the condensation between various aldehydes and anthranilamide in water at room temperature. This protocol is also applicable on a gram-scale synthesis. Clean reaction, ease of product isolation, short reaction time, mild reaction conditions (room temperature), low catalyst loading, low E-factor, and use <sup>20</sup> of water as solvent are the notable advantages of the present methodology and these features make this procedure to be a green synthetic protocol. We believe that our novel procedure will open up a new practical and convenient route for the synthesis of biologically important 2,3-dihydroquinazolin-4(1*H*)-ones.

#### 25 Experimental section

**General:** Melting points were determined on a glass disk with an electric hot plate and are uncorrected. <sup>1</sup>H NMR (400 MHz) and <sup>13</sup>C NMR (100 MHz) spectra were run in in CDCl<sub>3</sub> & DMSO-*d*<sub>6</sub> solutions (Bruker Avance 400). Chemical shift were recorded as

 $_{30}$   $\delta$  values in parts per million (ppm), and the signals were reported as s (singlet), d (doublet), t (triplet), m (multiplet) and coupling constants *J* were given in Hz. IR spectra were taken in a Perkin Elmer FTIR-Spectrum 400. Commercially available substrates were freshly distilled before the reaction. Solvents, reagents and  $_{35}$  chemicals were purchased from Sigma Aldrich and Merck.

#### General procedure for the synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones

*p*-Sulfonic acid calix[4]arene (1 mol%) was added to a solution of anthranilamide (1 mmol) and aldehyde/ketone (1 mmol) in water

<sup>40</sup> (1 mL) and the mixture was stirred at room temperature for a specified period of time. After completion of the reaction, cold distilled water (5 mL) was added to the reaction mixture. Then the product was filtered off and the catalyst was recovered by

evaporating the water. The recovered catalyst was reused for a <sup>45</sup> subsequent fresh batch of the reaction after reactivation. The crude product was recrystallized from hot ethanol to afford the pure product.

#### Acknowledgements

We thank the University of Malaya and the Malaysian Ministry 50 of Higher Education High Impact Research Grant UM.C/625/1/HIR/MOHE/05 for financial support.

#### Notes and references

<sup>a</sup> Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia. E-mail: rauzah@um.edu.my

55 <sup>b</sup> Department of Chemistry, Visva-Bharati (A Central University), Santiniketan 731235, India. E-mail: alakananda.hajra@visvabharati.ac.in

<sup>†</sup>*Electronic Supplementary Information (ESI) available:* [<sup>1</sup>*H*, <sup>13</sup>*C NMR data and Spectra, crystallographic data, E-factor calculations]. See* <sup>60</sup> DOI: 10.1039/b000000x/</sup>

- 1 (a) R. A. Sheldon, *Chemtech*, 1994, **25**, 38-47; (b) R. A. Sheldon and H. van Bekkum, Eds., *Fine Chemicals Through Heterogeneous Catalysis*, Wiley-VCH, Weinheim, Germany, 2001.
- 2 (a) P. I. Dalko and L. Moisan, Angew. Chem., Int. Ed., 2001, 40, 3726-3748; (b) B. List, J. Am. Chem. Soc., 2000, 122, 9336-9337; (c) J. Seayad and B. List, Org. Biomol. Chem., 2005, 3, 719-724; (d) J. H. Kim, I. Coric, S. Vellalath and B. List, Angew. Chem., Int. Ed., 2013, 52, 4474-4477; (e) B. List, Synlett, 2001, 1675-1686; (f) D. W. C. MacMillan, Nature, 2008, 455, 304-308; (g) D. Cantillo, B. Gutmann and C. O. Kappe, J. Am. Chem. Soc., 2011, 133, 4465-4475; (h) A. C. Cole, J. L. Jensen, I. Ntai, K. L. T. Tran, K. J. Weaver, D. C. Forbes and J. H. Davis, Jr, J. Am. Chem. Soc., 2002, 124, 5961-5963; (i) L. Myles, N. Gathergood and S. J. Connon, Chem. Commun., 2013, 49, 5316-5318; (j) S. Bertelsen and K. A. Jørgensen, Chem. Soc. Rev., 2009, 38, 2178-2189; (k) M. Gruttadauria, F. Giacalone and R. Noto, Chem. Soc. Rev., 2008, 37, 1666-1688; (1) L. W. Xu, L. Li and Z. H. Shi, Adv. Synth. Catal., 2010, 352, 243-279; (m) K. Manabe, Y. Mori and S. Kobayashi, Tetrahedron, 2001, 57, 2537-2544.
  - (a) P. I. Dalko and L. Moisan, Angew. Chem. Int. Ed., 2004,
    43, 5138-5175; (b) B. List, Chem. Commun., 2006, 819-824;
    (c) A. Dondoni and A. Massi, Angew. Chem., Int. Ed., 2008,
    47, 4638-4660; (d) E. N. Jacobsen and D. W. C. MacMillan,
    Proc. Natl. Acad. Sci. USA, 2010, 107, 20618-20619; (e) J. G.
    Hernandez and E. Juaristi, Chem. Commun., 2012, 48, 5396-5409.
  - 4 (a) Organic Catalysis; Wiley-VCH: Weinheim, Germany, 2004; pp. 1007-1249; (b) Organocatalysis; American Chemical Society: Washington, DC, USA, 2007; pp. 5413-5883.
  - 5 (a) P. T. Anastas and J. C. Warner, Green Chemistry: Theory and Practice, Oxford University Press, Oxford, 1998; (b) S. Shimizu, S. Shirakawa Y. Sasaki and C. Hirai, Angew. Chem. Int. Ed., 2000, 39, 1256-1259; (c) F. Zhang and H. Li, Chem. Sci., 2014, 5, 3695-3707; (d) R. A. Sheldon, I. Arends and U. Hanefeld, Green Chemistry and Catalysis, Wiley-VCH, Weinheim, 2007; (e) R. T. Baker and W. Tumas, Science, 1999, 284, 1477-1479; (f) C.-J. Li, Chem. Rev., 2005, 105, 3095-3166; (g) U. K. Lindstrom, Chem. Rev., 2002, 102, 2751-2772; (h) A. Lubineau, J. Auge' and Y. Queneau, Synthesis, 1994, 741-760; (i) G. Fioroni, F. Fringuelli, F. Pizzo and L. Vaccaro, Green Chem., 2003, 5, 425-428; (j) B. C. Ranu and K. Chattopadhyay, in Eco-friendly Processes and Procedures for the Synthesis of Fine Chemicals, ed. R. Ballini, Royal Society of Chemistry, Cambridge, UK, 2009, 186-215.

Page 6 of 6 View Article Onlin DOI: 10.1039/C4RA16374E

- (a) C. Capello, U. Fischer and K. Hungerbuhler, Green Chem., 6 2007, 9, 927-934; (b) D. Kumar, K. Seth, D. N. Kommi, S. Bhagat and A. K. Chakraborti, RSC Adv., 2013, 3, 15157-15168; (c) R. B. Nasir-Baig and R. S. Varma, Chem. Soc. Rev., 2012, 41, 1559-1584.
- 7 (a) K. Shanab, C. Neudorfer, E. Schirmer and H. Spreitzer, Curr. Org. Chem. 2013, 17, 1179-1187; (b) K. Alfonsi, J. Colberg, P. J. Dunn, T. Fevig, S. Jennings, T. A. Johnson, H. P. Kleine, C. Knight, M. A. Perry and M. Stefaniak, Green Chem., 2008, 10, 3-10; (c) H. C. Hailes, Org. Process Res. Dev., 2007, 11, 114-120.

10

15

20

25

30

35

40

45

60

Published on 22 December 2014. Downloaded by McMaster University on 22/12/2014 20:38:14.

- (a) R. N. Butler and A. G. Coyne, Chem. Rev., 2010, 110, 8 6302-6337; (b) S. Minakata and M. Komatsu, Chem. Rev., 2009, 109, 711-724; (c) A. Chanda and V. V. Fokin, Chem.
- Rev., 2009, 109, 725-748; (d) M. Chen, F. Zhang and H. Li, Curr. Org. Chem. 2013, 17, 1051-1057; (e) J.-N. Tan, H. Li and Y. Gu, Green Chem. 2010, 12, 1772-1782; (f) Y. Gu, R. De Sousa, G. Frapper, C. Bachmann, J. Barrault and F. Jérôme, Green Chem. 2009, 11, 1968-1972.
- 9 (a) L. Liu and D. Wang, "On Water" for Green Chemistry in Handbook of Green Chemistry 5: Reaction in Water., ed. P. T. Anastas and C.-J. Li, p. 207-228. Wiley-VCH, Weinbeim, 2010; (b) J. Mlynarski and J. Paradowska, Chem. Soc. Rev., 2008, 37, 1502-1511; (c) F. Fringuelli, O. Piermatti, F. Pizzo
- and L. Vaccaro, Eur. J. Org. Chem., 2001, 3, 439-455; (d) A. Manna and A. Kumar, J. Phys. Chem. A., 2013, 117, 2446-2454; (e) A. K. Chakraborti, S. Rudrawar, K. B. Jadhav, G. Kaur and S. V. Chankeshwara, Green Chem., 2007, 9, 1335-1340; (f) G. L. Khatik, R. Kumar and A. K. Chakraborti, Org. Lett., 2006, 8, 2433-2436.
- 10 S. Shimizu, N. Shimada and Y. Sasaki, Green Chem., 2006, 8, 608-614.
  - R. A. Sheldon, Chem. Soc. Rev., 2012, 41, 1437-1451. 11
- 12 (a) S. Sayin and M. Yilmaz, RSC Adv., 2014, 4, 2219-2225; (b) A. Casnati, Chem. Commun., 2013, 49, 6827-6830; (c) D. M. Homden and C. Redshaw, Chem. Rev., 2008, 108, 5086-5130.
- (a) F. Cardona, G. Isoldi, F. Sansone, A. Casnati and A. Goti, 13 J. Org. Chem., 2012, 77, 6980-6988; (b) S. Sayin, E. Yilmaz and M. Yilmaz, Org. Biomol. Chem., 2011, 9, 4021-4024; (c) A. A. Bhatti, M. A. Kamboh, I. B. Solangi and S. Memon, J. Appl. Polym. Sci., 2013, 130, 776-785.
- 14 (a) S. J. Dalgarno, K. M. Claudio-Bosque, J. E. Warren, T. E. Glass and J. L. Atwood, Chem. Commun., 2008, 1410-1412;
- (b) B. M. Rambo, S. K. Kim, J. S. Kim, C. W. Bielawski and J. L. Sessler, Chem. Sci., 2010, 1, 716-722; (c) J. L. Atwood, S. J. Dalgarno, M. J. Hardie and C. L. Raston, Chem. Commun., 2005, 337-339; (d) C. D. Gutsche and K. C. Nam, J. Am. Chem. Soc., 1988, 110, 6153-6162;
- 15 (a) L. Liu, Y. L. Wang, Y. C. Han and Y. J. Chen, Green 50 Chem., 2008, 10, 635-640; (b) D. L. da Silva, S. A. Fernandes, A. A. Sabino and A. de Fátima, Tetrahedron Lett., 2011, 52, 6328-6330; (c) S. A. Fernandes, R. Natalino, P. A. R. Gazolla, M. J. da Silva and G. N. Jham., Tetrahedron Lett., 55 2012. 53. 1630–1633.
  - (a) A. M. Farghaly, R. Soliman, M. A. Khalil, A.A. Bekhit, A. 16 el-Din and A. Bekhit, Boll. Chem. Farm., 2002, 141, 372-378; (b) H. J. Yale and M. Kalkstein, J. Med. Chem., 1967, 10, 334-336; (c) M. Hour, L. Huang, S. Kuo, Y. Xia, K. Bastow,
  - Y. Nakanishi, E. Hamel and K. Lee, J. Med. Chem., 2000, 43, 4479-4487; (d) M. Rahman, A. K. Bagdi, S. Mishra and A. Hajra, Chem. Commun., 2014, 50, 2951-2953.
    - 17 R. P. Maskey, M. Shaaban, I. Grun-Wollnu and H. Laatsch, J. Nat. Prod., 2004, 67, 1131-1134.
- (a) Y. S. Sadanadam, R. M. Reddy and A. Bhaskar, Eur. J. 18 65 Med. Chem., 1987, 22, 169-173; (b) J. I. Levin, P. S. Chan, T. Bailey, A. S. Katocs and A. M. Venkatesan, Bioorg. Med. Chem. Lett., 1994, 4, 1141-1146; (c) G. L. Neil, L. H. Li, H. H. Buskirk and T. E. Moxley, Cancer Chemother. Rep., 1972, 56, 163-173. 70

- 19 (a) P. VNS Murthy, D. Rambabu, G. R. Krishna, C. M. Reddy, K. R. S. Prasad, M. V. B. Rao and M. Pal, Tetrahedron Lett., 2012, 53, 863-867; (b) N. B. Darvatkar, S. V. Bhilare, A. R. Deorukhkar, D. G. Raut and M. M. Salunkhe, Green Chem. Lett. Rev., 2010, 3, 301-306; (c) S. Rostamizadeh, A. M. Amani, G. H. Mahdavinia, H. Sepehrian and S. Ebrahimi, Synthesis, 2010, 1356-1360; (d) X. Cheng, S. Vellalath, R. Goddard and B. List, J. Am. Chem. Soc., 2008, 130, 15786-15787; (e) Z.-H. Zhang, H.-Y. Lu, S.-H. Yang and J.-W. Gao, J. Comb. Chem., 2010, 12, 643-646; (f) P. Salehi, M. Dabiri, M.A. Zolfigol and M. Baghbanzadeh, Svnlett, 2005, 1155-1157; (g) F. Li, Y. Feng, Q. Meng, W. Li, Z. Li, Q. Wang and F. Tao, Arkivoc, 2007, (i), 40-50; (h) M. Desroses, M. Scobie and T. Helleday, New J. Chem., 2013, 37, 3595-3597; (i) A. Davoodnia, S. Allameh, A. R. Fakhari and N. Tavakoli-Hoseini, Chin. Chem. Lett., 2010, 21, 550-553; (j) M. Sharma, S. Pandey, K. Chauhan, D. Sharma, B. Kumar and P. M. S. Chauhan, J. Org. Chem., 2012, 77, 929-937; (k) M. Min Wang, T. T. Zhang and Z. G. Song, Chin. Chem. Lett., 2011, 22, 427-430; (l) J. Safari and S. Gandomi-Ravandi, J. Mol. Catal. A: Chem., 2014, 390, 1-6; (m) K. Ramesh, K. Karnakar, G. Satish, K. H. V. Reddy and Y. V. D. Nageswar, Tetrahedron Lett., 2012, 53, 6095-6099; (n) D. Huang, X. Li, F. Xu, L. Li, and X. Lin, ACS Catal., 2013, 3, 2244-2247; (o) S. Santra, M. Rahman, A. Roy, A. Majee and A. Hajra, Catal. Commun., 2014, 49, 52-57; (p) Y. Zheng, M. Bian, X.-Q. Deng, S.-B. Wang and Z.-S. Quan, Arch. Pharm. Chem. Life Sci., 2013, 346, 119-126; (q) M. Abdollahi-Alibeik and E. Shabani, J. Iranian Chem. Soc., 2014, 11, 351-359; (r) S. Vasudhevan and R. J. Karunakaran, Int. J. ChemTech Res., 2013, 5, 2844-2853; (s) J. Wang, Y. Zong, R. Fu, Y. Niu, G. Yue, Z. Quan, X. Wang and Y. Pan, Ultrason. Sonochem., 2014, 21, 29-34; (t) V. B. Labade, P. V. Shinde and M. S. Shingare, Tetrahedron Lett., 2013, 54, 5778-5780; (u) G. A. N. K. Durgareddy, R. Ravikumar, S. Ravi and S. R. Adapa, J. Chem. Sci., 2013, 125, 175-182; (v) X. Liu, D.-H. Hu and H. Shen, Asian J. Chem., 2012, 24, 1365-1367.
  - 20 (a) S. Santra, A. K. Bagdi, A. Majee and A. Hajra, RSC Adv., 2013, 3, 24931-24935; (b) S. Santra, A. Majee and A. Hajra, Tetrahedron Lett., 2012, 53, 1974-1977; (c) M. Rahman, D. Kundu, A. Hajra and A. Majee, Tetrahedron Lett., 2010, 51, 2896-2899.
- (a) S. Shinkai, Pure Appl. Chem., 1986, 58, 1523-1528; (b) S. 21 Shinkai, K. Araki, T. Tsubaki, T. Arimura and O. Manabe, J. Chem. Soc., Perkin Trans. 1, 1987, 11, 2297-2299; (c) M. Makha and C. L. Raston, Tetrahedron Lett., 2001, 42, 6215-6217; (d) M. Makha and C. L. Raston, Chem. Commun., 2001, 2470-2471.
- A. Kamal, V. Srinivasulu, B. N. Seshadri, N. Markandeya, A. 22 Alarifi and N. Shankaraiah, Green Chem., 2012, 14, 2513-2522
- CCDC deposition code of compound 3n: 1026285. 23

## 125

105

110

115

120