

Tetrahedron Letters 39 (1998) 991-992

TETRAHEDRON LETTERS

Oxidation of 3-Sulfanyl-alcohols with Sodium Metaperiodate: New Synthesis of Sultines

S. Yolka, R. Fellous*, L. Lizzani-Cuvelier, M. Loiseau

Laboratoire de Chimie des Arômes, Université de Nice-Sophia Antipolis, Faculté des Sciences, Parc Valrose, 06108 Nice Cedex 2, France.

Received 22 October 1997; accepted 1 December 1997

Abstract: A simple and highly efficient method for preparation of sultines by oxidation of 3-sulfanyl-alcohols with sodium metaperiodate is described. © 1998 Elsevier Science Ltd. All rights reserved.

There are no reports available about sultines as natural products, except from Rochard's¹ work which showed the presence of the 3-propyl-1,2-oxathiolane-2-oxide in the yellow passion fruit. Nevertheless, many synthetic routes have been investigated from hydroxysulfoxydes upon reaction with $SO_2Cl_2^2$, sulfones³ and chloration of sulfanyl-alcohols⁴. In this latter case, Givens et al.⁴⁶ suggested that the sultines were formed via a disulfide intermediate. Their electrophilic cleavage was facilitated by neighboring nucleophilic species to give alkylsulfonates⁵⁶ or intramolecular cyclisations⁵⁶.

We report herein a new synthetic route to sultines with sodium metaperiodate oxidation of 3-sulfanylalcohols. This synthesis first forms a disulfide compound which was isolated. The sultines can be obtained easily in a short time (4 hours) with good yields using excess of oxidizing agent (Table 1).

3-suifanyi-alcohois" (10 mmoi)	NaIO4 (mmol)	Reaction Time	Ratio of disulfides (%)	Ratie of sultines ^b (%)	Structure of sultines
CH3-CH-CH2-CH2-OH SH	20	4h	0*	80*	
	10	30 min 24h	50▲ 31▲	39▲ 54▲	1
	1.4	30 min 24h	89 [▲] 84 [▲]	0≜ 3≜	
CH3-CH-CH2-CH2-OH SH	20	4 <u>h</u>	0*	98*	٥
	10	30 min 24h	36▲ 22▲	64▲ 76▲	2 5 0 Pr
СH3 СH3 SH CH3	20	4h	0*	75*	
	10	30 min 24h	32▲ 32▲	57▲ 57▲	3

Table 1

*The 3-suifanyl-alcohols were obtained as described in note 6. Mixture of 40/60 diastereomers cis/trans. *Isolated yields. *Ratio established by GC.

Sultines 2 and 3 have never been described. Attempts to synthetise 3 according to the method of King et al.⁴⁰, by chloration of the 4-methyl-4-sulfanylpentan-2-ol in aqueous solution, were unsuccessful. The Table 1 shows that we are able to obtain either disulfide, sultine or a mixture of both depending of the amount of oxidizing agent.

Together with the formation of disulfide and sultine, the presence of iodine was observed. This suggests a change in the oxidation number of iodine from +7 to 0. The presence of iodine is confirmed by sodium thiosulfate titration.

We have established that 1/7 mol of NaIO₄ is necessary to oxidize one mole of 3-sulfanyl-alcohol in disulfide according to the equation (1):

$$14 \text{ RSH} + 2 \text{ IO}_4^- + 2 \text{ H}_3\text{O}^+ \rightarrow 7 \text{ RSSR} + \text{I}_2 + 10 \text{ H}_2\text{O}$$
(1)

Iodine is also involved in the formation of sultines by reaction with the disulfide but this reaction is very slow: only 3% were formed in 24 hours (Table 1). Further addition of 6/7 mol of NaIO₄ gave the sultines according to the equation (2):

$$7 \text{ RSSR} + 6 \text{ IO}_4^- + 6 \text{ H}_3\text{O}^+ \rightarrow 14 \text{ sultimes} + 3 \text{ I}_2 + 16 \text{ H}_2\text{O}$$
(2)

Conclusion

The oxidation of 3-sulfanyl-alcohols with NaIO₄ represents a new suitable route to sultines. This method proceeds under very simple reaction conditions and can be performed with sodium metaperiodate, a cheap and safe oxidizing agent. A detailed mechanistic study of this oxidation reaction is in progress.

General procedure for oxidation reactions

A solution of NaIO₄ (20 mmol) in water (40 ml) was added to a solution of 3-sulfanyl-alcohol (10 mmol) in acetonitrile (20 ml) at room temperature. The mixture was stirred for 4 hours and then extracted with Et_2O (3 x 20 ml). The combined organic solutions were washed with a solution of sodium thiosulphate (0.1M) and dried over Na₂SO₄. The sultines were obtained as a yellow oil with a purity up to 95%, after evaporation of the solvent under reduced pressure. No further purification is required. The sultines were identified by FTIR, GC/MS and multinuclear NMR spectroscopy⁷.

Acknowledgements: We wish to thank Dr. Stephane Daniele for private communication.

References and Notes

- 1. Rochard, S. Doctorat ès Sciences Chimie, Université de Nice Sophia-Antipolis, 1994.
- Sharma, N. K.; De Reinach-Hirtzbach, J. F.; Durst, T. Can. J. Chem. 1976, 54, 3012-25. Connolly, T. J.; Durst, T. Tetrahedron Lett. 1997, 38, 1337-40.
- Durst, T.; Charlton, J. L.; Mount, D. B. Can. J. Chem. 1986, 64, 246-49. Harpp, D. N.; Gleason, J. G.; Ash, D. K. J. Org. Chem. 1971, 36, 322-26. Dodson, R. M.; Hammen, P. D.; Davis, R. A. J. Org. Chem. 1971, 36, 2693-98.
- (a) Givens, E. N.; Hamilton, L. A. J. Org. Chem. 1967, 32, 2857-60. (b) King, J. F.; Rathore, R. Tetrahedron Lett. 1989, 30, 2763-66. (c) King, J. F.; Skonieczny, S.; Khemani, K. C.; Stothers, J. B. J. Am. Chem. Soc. 1983, 105, 6514-15.
- 5. (a) Oae, S.; Takata, T. Tetrahedron Lett. 1980, 21, 3213-16. (b) Doi, J. T.; Luehr, G. W.; Musker, W. K. J. Org. Chem. 1985, 50, 5716-19.
- The 3-sulfanyl-alcohols were prepared via the reaction of thioacetic acid with crotonaldehyde, trans-2-hexenal and mesityl oxide. Reduction with LiAlH4 gave 3-sulfanylbutan-1-ol, 3-sulfanylhexan-1-ol and 4-methyl-4sulfanylpentan-2-ol in 65% yield after purification.
- Spectroscopic data for the selected compounds are given below. ¹H and ¹³C(¹H) NMR were run on Brucker AC-200 spectrometer with CDCl₃ as solvent. COSY 2D NMR experiments were used for the assignments of signals.

3-methyl-1,2-oxathiolane-2-oxide: IR: 1121 cm⁻¹. <u>Cis isomer</u>: ¹H NMR: δ (ppm) 4.88-4.73 (m, 1H), 4.46-4.25 (m, 1H), 3.19-2.96 (m, 1H), 2.39-2.19 (m, 2H), 1.41 (d, 3H, J=6.5 Hz); ¹³C[¹H] NMR: δ 75.3 (C-5), 62.7 (C-3), 29.7 (C-4), 11.1 (CH₃); MS (70eV): 39 (34), 41 (84), 42 (18), 43 (15), 48 (13), 50 (26), 55 (M⁺, 100), 56 (23), 78 (59), 90 (7), 120 (36), 121 (1), 122 (1). <u>Trans isomer</u>: ¹H NMR: δ (ppm) 4.88-4.73 (m, 2H), 4.58-4.25 (m, 1H), 3.46-3.27 (m, 1H), 2.86-2.62 (m, 1H), 1.99-1.83 (m, 1H), 1.23 (d, 3H, J=7.5 Hz); ¹³C[¹H] NMR: δ 74.7 (C-5), 66.6 (C-3), 30.1 (C4), 13.1 (CH₃); MS (70eV): 39 (32), 41 (76), 42 (16), 43 (15), 48 (12), 50 (23), 55 (M⁺, 100), 56 (22), 78 (54), 90 (6), 120 (37), 121(<1), 122 (<1).

3, 3-dimethyl, 5-methyl-1, 2-oxathiolane-2-oxide: IR: 1117 cm⁻¹ <u>Cis isomer</u>: ¹H-NMR: δ (ppm) 4.90-4.70 (m, 1H), 2.42-1.95 (m, 2H), 1.54 (d, 3H, J=6.2 Hz), 1.39 (s, 3H), 1.25 (s, 3H); ¹³C(¹H) NMR: δ 86.7 (C-5), 43.3 (C-4), 23.5 (C-5), 21.4 (C-3), 20.5 (C-3); MS (70eV): 41 (70), 42 (7), 43 (71), 55 (89), 56 (28), 69 (55), 83 (M⁺, 100), 92 (43), 104 (14), 133 (8), 148 (3). <u>Trans isomer</u>: ¹H-NMR: δ (ppm) 5.31-5.09 (m, 1H), 2.42-1.95 (m, 2H), 1.44 (d, 3H, J=6.4 Hz), 1.36 (s, 3H), 1.35 (s, 3H); ¹³C(¹H) NMR: δ 84.1 (C-5), 44.3 (C-4), 24.3 (C-5), 21.2 (C-3), 20.1 (C-3) MS (70eV): 41 (70), 42 (7), 43 (91), 55 (M⁺, 100), 56 (32), 69 (48), 83 (99), 92 (60), 104 (17), 133 (<1), 148 (2).