Inorganica Chimica Acta 385 (2012) 135-139

Contents lists available at SciVerse ScienceDirect

Inorganica Chimica Acta

journal homepage: www.elsevier.com/locate/ica

Activation reactions of 1,1-dialkoxoalkanes and unsaturated O-donors by titanium tetrafluoride

Fabio Marchetti, Guido Pampaloni*, Lorenzo Biancalana

Università di Pisa, Dipartimento di Chimica e Chimica Industriale, Via Risorgimento 35, I-56126 Pisa, Italy

ARTICLE INFO

Article history: Received 8 November 2011 Received in revised form 20 January 2012 Accepted 23 January 2012 Available online 6 February 2012

Keywords: Titanium Fluoride Activation O-Donor Coordination

1. Introduction

The reactivity of titanium tetrafluoride has been much less developed with respect to that of the heavier congeners, especially the tetrachloride TiCl₄ [1]. One of the reasons is probably related to the particular inertness of TiF₄, due to the polymeric structure in the solid state [2] and the strength of the Ti–F bond (D_0 = 584 kJ/mol, to be compared with Ti–Cl, D_0 = 429.3 kJ/mol, Ti–Br, D_0 = 366.9 kJ/mol, and Ti–I, D_0 = 296.2 kJ/mol) [3].

Titanium tetrafluoride and *O*-donors L generally form 1:2 adducts of formula TiF₄L₂, L = thf [4], Ph₃PO [5], disso [6], dimethylformamide [6c] dimethylacetamide [6c,7], tetramethylurea [6c], substituted pyridine- and quinoline-*N*-oxide [8]. Complex TiF 4(dme) is obtained by straightforward 1:1 reaction of TiF₄ with 1,2-dimethoxyethane (dme) [9]. Polymeric fluorine-bridged complexes may be obtained with weak Lewis bases such as pyridine [6c] and benzonitrile [10].

Compounds of formula TiF_4L_2 show generally the *cis*-configuration although *cis*-*trans* equilibria in solution have been observed when strongly donating, sterically hindered ligands are involved [8a,11].

The strong Ti–F bond is not cleaved by alcohols, which react with TiF₄ affording TiF₄(ROH)₂ adducts, R = Me [12], Et [6c,12], Prⁱ [12]. It is remarkable that TiF₄ forms a stable coordination adduct upon reaction with water [13], in contrast with the behavior

ABSTRACT

The reactivity of TiF₄ with a variety of non cyclic 1,1-dialkoxoalkanes $[CH_2(OR)_2, R = Me, Et, Me_2C(OMe)_2, MeCH(OEt)_2, ClCH_2CH(OEt)_2, CH(OMe)_3, PhC=CCH(OEt)_2], 1,3-dioxolane, N_2CHCO_2Et and 1,2-epoxybutane has been investigated. Activation, including fragmentation and/or rearrangement of the organic moiety, has been observed at room temperature in some cases; it generally occurs unselectively via C-O bond fission and the formation of new C-O, C-H and C-C bonds. Small differences in the structure of the organic substrate may determine significant differences in the reactivity with TiF₄.$

© 2012 Elsevier B.V. All rights reserved.

Inorganica Chimica Acta

normally exhibited by other early transition metal halides in high oxidation state, which rapidly hydrolyze to give oxide derivatives [14]. Moreover HF is not produced in the reaction of TiF₄ with salenH₂ in boiling thf, Scheme 1, which affords the 1:1 adduct TiF₄(salenH₂): cleavage of the Ti–F bond and formation of TiF₂(salen) is achieved in the presence of the SiMe₃ fragment [15].

Recently we reported on the reactions of MX_5 (M = Nb, Ta; X = F, Cl, Br, I), MoCl₅ and WCl₆ with O-donor species, showing that the formation of the strong M–O bond favors the C–O cleavage within the organic unit, possibly resulting in fragmentation and rearrangement reactions even at room temperature [16]. In particular the reactions involving diethers showed the formation of new C–C and C–O bonds, and higher reactivity was observed with 1,1-dial-koxyalkanes rather than 1,2-dialkoxyalkanes [17,18]. These reactions are strongly influenced by the nature of the halide, the fluorides MF₅ (M = Nb, Ta) usually activating the diether at high temperatures only [16a].

At variance to niobium and tantalum derivatives, the group 4 tetrahalides MX_4 (M = Ti, Zr, X = Cl, Br, I) adducts with 1,2-dialkoxoalkanes are less prone to undergo activation of the coordinated organic fragments [9]. On the other hand, it was reported that TiCl₄ may show the "deprotecting" behavior typical of Lewis acids with respect to 1,1-dialkoxylkanes (acetal and ketals), thus determining the generation of the relevant carbonylic compounds [19]. Conversely, it was found that TiCl₄ was able to activate substituted dioxolanes to give chlorinated organic compounds [20].

In this context, we decided to explore the chemistry of the relatively inert TiF_4 with 1,1-dialkoxoalkanes or unsaturated oxygen compounds. The results of this investigation are reported in the present paper.



^{*} Corresponding author. Tel.: +39 050 2219 219; fax: +39 050 2219 246.

E-mail addresses: fabmar@dcci.unipi.it (F. Marchetti), pampa@dcci.unipi.it (G. Pampaloni).

^{0020-1693/\$ -} see front matter \odot 2012 Elsevier B.V. All rights reserved. doi:10.1016/j.ica.2012.01.051



Scheme 1. The reactions of salenH₂ with TiF₄.

2. Results and discussion

Titanium tetrafluoride reacted slowly with $CH_2(OR)_2$, R = Me, Et, $Me_2C(OMe)_2$, $MeCH(OEt)_2$, $CICH_2CH(OEt)_2$, either at room temperature or at ca. 80 °C, affording mixtures of products which prevented the identification of any coordination compound. However, it was possible to identify the products of the activation of the organic substrate after exhaustive hydrolysis of the reaction mixture, Table 1, runs 1–5.

The data of Table 1 suggest that the reactions are not selective, similarly to what observed in the analogous reactivity of Group 5 metal pentahalides [17]. The reaction of TiF₄ with $CH_2(OMe)_2$ (Table 1, run 1) produced methyl formate and dimethylether. These compounds were identified together with a significative amount of unreacted $CH_2(OMe)_2$. In addition MeOH was found in the reaction mixture after hydrolysis. An increase of the reaction temperature and/or the $CH_2(OMe)_2/Ti$ molar ratio caused a substantial increase of dimethylether (Table 2, run 2). The presence of HCO₂Me and Me₂O suggests that the activation reaction proceeds with cleavage of C–O and C–H bonds followed by formation of new C–O and C–H bonds, see Scheme 2a and b. The presence of methanol might be the consequence of the formation of the [Ti–OMe] unit, see Scheme 2c.

According to the $Me_2O/MeOH$ molar ratio detected (16:1, Table 1, run 1), process **b** should occur more easily than process **c** (Scheme 2).

The reaction of TiF_4 with $CH_2(OEt)_2$ proceeds similarly to what discussed for $CH_2(OMe)_2$. Thus HCO_2Et and EtOH (ratio 1:3) have been identified after hydrolysis of the reaction mixture (Table 1,

run 2). It is interesting to note that the analogous reaction involving MF_5 (M = Nb, Ta) gave the mixed ether OMeEt, as a consequence of the concerted activation of C–O and C–H bonds [17]. In the case of TiF₄ and CH₂(OEt)₂, ethers do not form thus suggesting that pathways **b** and **c** (Scheme 2) are almost non viable (see Scheme 3) and that a small variation in the structure of the substrate can determine strong difference in the nature of the reaction products.

The hydrolysis of the mixture obtained by reaction of TiF_4 with $MeCH(OEt)_2$ gave only EtOH and Et_2O as identifiable materials (Table 1, run 3). Although a similar process to that reported in Scheme 2 may be invoked, it has not been possible to identify the other products necessary for the full justification of the fragmentation scheme.

A substantially different reactivity was observed by formal substitution of a methyl hydrogen with chlorine. In fact the reaction between TiF₄ and ClCH₂CH(OEt)₂ gave ClCH₂CHO (Table 1, run 4): in this case, TiF₄ seems to behave as a traditional Lewis acid, "deprotecting" the carbonyl functionality (see Section 1) [19,21]. Hydrolysis of the reaction mixture did not change the relative composition of the organic compounds. A similar result was obtained on increasing the reaction temperature or the substrate/Ti molar ratio (Table 2, run 6).

The reaction of $Me_2C(OMe)_2$ with TiF₄ was fast even at room temperature and gave, after hydrolysis, MeOH plus minor amounts of MeCO₂Me, Me₂O and Me₂CO (Table 1, run 5). A possible scheme of formation is given (Scheme 4). It is interesting to note that the same reaction performed with NbF₅ afforded mesityloxide [17], i.e. the product of acetone dehydration, thus confirming the higher affinity for water of NbF₅ with respect to TiF₄ [1,14].

Trimethoxymethane reacted with TiF_4 at room temperature affording Me₂O and HCO₂Me in comparable amounts (molar ratio ca. 14:10; Table 1, run 6). Analogous activation process was observed previously in the case of NbF₅ [17]. The presence of dimeth-ylether suggests that the coupling between different fragments, including the formation of a new C–O bond, is operative in the system, see Scheme 5. Methanol was observed in the reaction mixture after hydrolysis.

An interesting case of C–C bond formation was observed when a dichloromethane suspension of TiF₄ was treated with 1,3-dioxolane. After heating and hydrolysis steps, HCO₂Me, 1,4-dioxane, MeO(CH₂)₂OMe, and Me₂O were identified in 8:20:10:10 molar ratios (Table 1, run 7). With a conversion of ca. 20%, 1,4-dioxane represents the product formed in higher amount in this reaction. A tentative suggestion concerning the formation of 1,4-dioxane from 1,3-dioxolane is sketched in Scheme 6 and implies the formation of a new C–C bond with intermolecular transfer of a [CH₂] fragment.

Table 1

Summary of the organic compounds obtained from TiF₄ and 1,1-dialkoxoalkanes, CH(OMe)₃, PhC=CCH(OEt)₂, 1,2-epoxybutane, N₂CHCO₂Et, after hydrolysis of the reaction mixtures.

Run	Reagent (L)	Products (molar ratios) ^d
1	$CH_2(OMe)_2^a$	$HCO_2Me + CH_2(OMe)_2 + MeOH + Me_2O (10:5:1:16)$
2	$CH_2(OEt)_2^a$	$CH_2Cl_2 + HCO_2Et + CH_2(OEt)_2 + EtOH (10:1:5:3)$
3	MeCH(OEt) ₂ ^a	$CH_2Cl_2 + MeCH(OEt)_2 + EtOH + Et_2O(4:1:4:1)$
4	CICH ₂ CH(OEt) ₂ ^{b,c}	CH ₂ Cl ₂ + ClCH ₂ CHO + ClCH ₂ CH(OEt) ₂ + ClCH ₂ CO ₂ Et (10:14:1:0.5)
5	$Me_2C(OMe)_2^a$	CH ₂ Cl ₂ + MeCO ₂ Me + MeOH + Me ₂ O + Me ₂ CO (10:0.4:4:0.3:0.4)
6	$HC(OMe)_3^a$	$CH_2Cl_2 + HCO_2Me + MeOH + Me_2O$ (10:7:1:5)
7	1,3-dioxolane ^a	CH ₂ Cl ₂ + HCO ₂ Me + 1,3-dioxolane + 1,4-dioxane + MeO(CH ₂) ₂ OMe + Me ₂ O (100:8:14:20:10:10)
8	$PhC \equiv CCH(OEt)_2^a$	$CH_2Cl_2 + PhCC \equiv CH(OEt)_2 + EtOH (5:3:3)$
9	1,2-epoxybutane ^b	$CH_2Cl_2 + CH_2(OH)CH(OH)Et (10:8)$
10	N ₂ CHCO ₂ Et ^a	$CH_2Cl_2 + CH_2FCO_2Et$ (3:1)

^a At 298 K.

^b After heating.

^c L/Ti molar ratio = 2.

^d CH₂Cl₂ used as internal standard (CH₂Cl₂/Ti molar ratio = 1).

F.	Marchetti	et	al./	'Inorganica	Chimica	Acta	385	(2012)	135–	13	9
----	-----------	----	------	-------------	---------	------	-----	--------	------	----	---

Table 2
Experimental details of the reactions between TiF4 and O-donors.

Run	TiF ₄ (mmol)	L	L/M molar ratio	Reaction time (h)	NMR spectra
1	0.44	CH ₂ (OMe) ₂	1	48	¹ H NMR: HCO ₂ Me [8.11 (s, CH), 3.79 ppm (s, Me)]; CH ₂ (OMe) ₂ [4.61 (s, CH ₂), 3.39 ppm (s, Me)]; 3.48 ppm (br); Me ₂ O [3.39 ppm (s, Me)]. HCO ₂ Me:CH ₂ (OMe) ₂ :Me ₂ O = 10:6:3 After hydrolysis (12 h): ¹ H NMR: HCO ₂ Me [8.08 (s, CH), 3.76 ppm (s, Me)]; CH ₂ (OMe) ₂ [4.58 (s, CH ₂), 3.37 ppm (s, Me)]; MeOH [3.48 ppm (s, Me)]; Me ₂ O [3.33 ppm (s, Me)]. HCO ₂ Me:CH ₂ (OMe) ₂ :MeOH:Me ₂ O = 10:51:16. ¹³ C NMR: CH ₂ (OMe) ₂ [97.4 (CH ₂), 55.0 ppm (CH ₃)]; HCO ₂ Me [161.9 ppm] (CH): 50.8 ppm (CH): 50.8 ppm (CH): 50.8 ppm (SH); MeO = 10:51:16. ¹³ C NMR: CH ₂ (OMe) ₂ [97.4 (CH ₂), 55.0 ppm (CH ₃)]; HCO ₂ Me [161.9 ppm] (CH): 50.8 ppm (CH): 50.8
2	0.45	CH ₂ (OMe) ₂	2	48	After heating, no hydrolysis: ¹ H NMR: HCO ₂ Me [8.05 (s, CH), 3.73 ppm (s, Me)]; CH ₂ (OMe) ₂ [4.56 (s, CH ₂), 3.34 ppm (s, Me)]: Me ₂ O [3.36 ppm (s, Me)]. HCO ₂ Me (H ₂ OMe) ₂ : Me ₂ O = 4:1:7
3	0.69	CH ₂ (OEt) ₂	1	96	¹ H NMR: CH ₂ Cl ₂ [5.31 ppm]; HCO ₂ Et [8.06 (s, CH), 4.24 (q, CH ₂), 1.31 ppm (t, CH ₃)]; CH ₂ (OEt) ₂ [4.69 (s, CH ₂ (OEt) ₂), 3.61 ppm (q, CH ₂ (OCH ₂ CH ₃) ₂) 1.22 ppm (t, CH ₃)]; 3.38 ppm (s); 1.38 ppm (t). CH ₂ Cl ₂ :HCO ₂ Et:CH ₂ (OEt) ₂ = 10:1:5 After hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.30 ppm]; HCO ₂ Et [8.05 (s, CH), 4.23 (m, CH ₂), 1.31 ppm (t, CH ₃)]; CH ₂ (OEt) ₂ [4.67 (s, CH ₂ (OEt) ₂), 3.61 (m, CH ₂ (OCH ₂ CH ₃) ₂), 1.22 ppm (m, CH ₃)]; EtOH [3.70 (quint, CH ₂), 2.02 (m, CH ₄)]; 3.37 ppm CH ₂ Cl ₂ :HCO ₂ Et/CH ₄ (OET) ₂ :FtOH = 10:1:5:3
4	0.71	MeCH(OEt) ₂	1	96	After hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.29 ppm]; MeCH(OEt) ₂ [4.67 ppm (q, CH), 3.57 ppm (m, CH ₂), 1.29 ppm (d, CH ₃ CH), 1.21 (t, OCH ₂ CH ₃)]; EtOH [3.66 (q, CH ₂), 2.66 (s, OH), 1.20 ppm (t, Me)]; Et ₂ O [3.47 (q, CH ₂), 1.18 ppm (t, Me)]. CH ₂ Cl ₂ :MeCH(OEt) ₂ :EtOH:Et ₂ O = 4:1:4:1. ¹³ C NMR: CH ₂ Cl ₂ [53.4 ppm]; Et ₂ O [67.5 (CH ₂), 15 2 ppm (Me)]: EtOH [58.0 (CH ₂), 18 1 ppm (Me)]
5	0.48	CICH ₂ CH(OEt) ₂	1	96	At 298 K: ¹ H NMR: CICH ₂ CHO [9.62 (s, CH), 4.07 ppm (s, CH ₂)]; CICH ₂ CH(OEt) ₂ [4.64 (t, CH), 3.61 ppm (m, OCH ₂), 3.52 (d, CICH ₂ CH (OE), 1.24 ppm (t, Me]. CICH ₂ CHO:CICH ₂ CH(OEt) ₂ = 1:5. ¹³ C NMR: CICH ₂ CH(OEt) ₂ [101.6 (CH), 62.5 ppm (OCH ₂), 43.8 (CICH ₂), 15.1 ppm (CH ₃)] After heating (18 h): ¹ H NMR: CICH ₂ CHO [9.62 (s, CH), 4.07 ppm (s, CH ₂)]; CICH ₂ CH(OEt) ₂ [4.64 ppm (t, CH), 3.51 ppm (d, CICH ₂), 3.66 (m, OCH ₂), 12.4 (t, Me]). CICH ₂ CHO:CICH ₂ CH(OEt) ₂ = 1:5
6	0.70	CICH ₂ CH(OEt) ₂	2	96	¹ H NMR: CH_2Cl_2 [5.29 ppm]; $ClCH_2CH0$ [9.41 ppm (s, CH), 4.01 ppm (s, br, CH ₂)]; $ClCH_2CH(OEt)_2$ [4.60 (t, CH), 3.48 (d, ClCH ₂), 3.63 (m, OCH ₂), 1.21 ppm (m, Me)]. CH_2Cl_2 : $ClCH_2CH0:ClCH_2CH(OEt)_2 = 10:2:10.$ ¹³ C NMR: CH_2Cl_2 [5.3.4 ppm]; $ClCH_2CH(OEt)_2$ [101.6 (CH), 62.5 (OCH ₂), 43.8 (ClCH ₂), 15.1 ppm (Me)] After heating (18 h): ¹ H NMR δ (ppm): CH_2Cl_2 [5.29 ppm]; $ClCH_2CH0$ [9.40 ppm (s, CH), 4.04 ppm (s, CH ₂)]; 5.90 ppm; $ClCH_2CH(OEt)_2$ [4.60 (m, CH), 3.62 (m, OCH ₂), 3.47 (m, ClCH ₂), 1.20 (m, Me)]. CH ₂ Cl ₂ : $ClCH_2CH0:ClCH_2CH0:ClCH_2CH(OEt)_2$ = 10:1:15 After heating (18 h) and hydrolysis: ¹ H NMR: CH_2Cl_2 [5.27 ppm]; $ClCH_2CH0$ [9.39 (s, CH), 4.02 ppm (s, CH ₂)]; $ClH_2CH0(OEt)_2$ [4.58 (t, CH), 3.61 (m, OCH ₂), 3.46 (d, ClCH ₂), 1.18 ppm (t, Me)]; $ClCH_2CO_2Et$ [4.20 (q, OCH ₂), 4.02 (s, ClCH ₂), 1.29 ppm (br, Me)].
7	0.75	Me ₂ C(OMe) ₂	1	96	CH ₂ Cl ₂ :ClCH ₂ CHO:ClCH ₂ CH(OEt) ₂ :ClCH ₂ CO ₂ Et = 10:14:1:0.5 ¹ H NMR: CH ₂ Cl ₂ [5.31 ppm]; 3.72 (br) ppm. After hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.31 ppm]; MeCO ₂ Me [3.67 (s, CO ₂ Me), 2.06 ppm (s, MeCO ₂)]; MeOH [3.48 (s, Me), 1.85 ppm (br, OH)]; Me ₂ O [3.32 ppm (s, Me)]; Me ₂ CO [2.17 ppm (s, Me)]. CH ₂ Cl ₂ : MeCO ₂ Me: MeOH: Me ₂ O: Me ₂ CO = 10:0.4:4:0.3:0.4. ¹³ C NMR: CH ₂ Cl ₂ [5.3 4 ppm]; MeOH [3.66 ppm]
8	0.65	HC(OMe) ₃	1	96	^{CH} ₂ Cl ₂ [3.5, 4 ppm], McO1 [363 ppm]; ^H NMR: CH ₂ Cl ₂ [5.30 ppm]; HCO ₂ Me [8.06 (s, CH), 3.74 ppm (s, Me)]; Me ₂ O [3.31 (s, Me)]. CH ₂ Cl ₂ :HCO ₂ Me:Me ₂ O = 10:7:5. ¹³ C NMR: CH ₂ Cl ₂ [53.4 ppm]; HCO ₂ Me [161.3 (CH), 50.8 ppm (Me)]; Me ₂ O [60.5 ppm (Me)] After hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.30 ppm]; HCO ₂ Me [8.05 (s, CH), 3.74 (s, Me)]; MeOH [3.45 (s, Me)]; Me ₂ O [3.30 (s, Me)]. CH ₂ Cl ₂ :HCO ₂ Me: MeOH:Me ₂ O = 10:7:1:5. ¹³ C NMR: CH ₂ Cl ₂ [53.4 ppm]; HCO ₂ Me
9	0.84	1,3-dioxolane	1	96	[161.9 (CH), 50.8 (Me)]; Me ₂ O [60.3 ppm]; MeOH [50.8 ppm] ¹ H NMR δ (ppm): CH ₂ Cl ₂ [5.31 ppm]; HCO ₂ Me [8.09 (s, CH), 3.74 ppm (s, Me)]; 1,3-dioxolane [4.92 (s, OCH ₂ O), 3.89 ppm (s, OCH ₂ CH ₂)]. CH ₂ Cl ₂ :HCO ₂ Me 1,3-dioxolane = 5:1:2 After heating (12 h): ¹ H NMR: CH ₂ Cl ₂ [5.31 ppm]; HCO ₂ Me [8.08 (s, CH), 3.76 ppm (s, Me)]; 1,3-dioxolane [4.91 (s, OCH ₂ O), 3.89 ppm (s, OCH ₂ CH ₂)]. CH ₂ Cl ₂ :HCO ₂ Me:1,3-dioxolane = 100:13:8 After heating (18 h) and hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.30 ppm]; HCO ₂ Me [8.07 (s, CH), 3.74 ppm (s, Me)]; 1,3-dioxolane [4.89 ppm (s, OCH ₂ O), 3.87 ppm (s, OCH ₂ CH ₂]; 1,4-dioxane [3.69 (s, CH ₂)]; MeO(CH ₂) ₂ OMe [3.53 (s, CH ₂), 3.37 ppm (s, CH ₃)]; Me ₂ O [3.30 ppm (s, Me)]. CH ₂ Cl ₂ :HCO ₂ Me:1,3- dioxolane(1.4 dioxange(MOC(L)) OM(2) OM(2) OH(2)010010
10	0.43	PhC≡CCH(OEt) ₂	1	72	dioxolatie: 1,4-choxane: MeO(CH ₂) ₂ O(Me:Me ₂ O = 100:8: 14:20:10:10 After hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.30 ppm]; 9.44 ppm; PhCC=CH(OEt) ₂ [7.50, 7.34 (m, Ph), 5.51 (s, CH), 3.85, 3.71 (m, CH ₂), 1.30 ppm (t, Me)]; EtOH [3.68 (q, CH ₂), 1.25 ppm (t, CH ₃)]. CH ₂ Cl ₂ :PhCC=CH(OEt) ₂ :EtOH = 5:3:3. ¹³ C NMR: CH ₂ Cl ₂ [53.4 ppm]; PhC=CCH(OEt) ₂ [133.3, 131.9, 128.8, 128.2 (CH), 121.9 (m, Ph), 91.8, 85.2 (C=C), 60.9 (CH ₂), 15.1 ppm (CH ₃)]; EtOH [58.3 (CH ₂),
11	0.75	1,2- epoxybutane	1	96	^{10.5} ppmi (Cr ₃₃). ¹ H NMR: CH ₂ Cl ₂ [5.32 ppm]; 3.52 (br), 1.57 (br), 0.96 ppm (m) After hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.31 ppm]; CH ₂ (OH)CH(OH)Et [3.53 (m), 1.47 (m), 0.97 ppm (m)]. CH ₂ Cl ₂ :CH ₂ (OH)CH(OH)Et = 1:1 After heating (5 h) and hydrolysis: ¹ H NMR: CH ₂ Cl ₂ [5.31 ppm]; CH ₂ (OH)CH(OH)Et [3.52 (br), 1.57 (br), 0.96 ppm (m)]. CH ₂ Cl ₂ :CH ₂ (OH)CH(OH)CH(OH)Et = 10:8
12	0.73	N ₂ CHCO ₂ Et	1	96	¹ H NMR: CH_2CI_2 [5.33 ppm]; CH_2FCO_2Et [4.85 (d, ¹ J_{HF} = 47 Hz, CH ₂ F), 4.30 (q, CH ₂ O), 1.33 ppm (t, CH ₃)]. CH ₂ CI ₂ : CH ₂ FCO ₂ Et = 2:1. ¹³ C NMR: CH ₂ CI ₂ [53.4 ppm]; CH ₂ FCO ₂ Et [168.1 (d, CO ₂), 77.8 (d, CH ₂ F), 61.5 (CH ₂ O), 14.2 ppm (CH ₃)] After hydrolysis: ¹ H NMR: CH ₂ CI ₂ [5.33 ppm]; CH ₂ FCO ₂ Et [4.85 (d, ¹ J_{HF} = 47 Hz, CH ₂ F), 4.30 (q, CH ₂ O), 1.33 ppm (t, CH ₃)]. CH ₂ CI ₂ : CH ₂ FCO ₂ Et = 3:1

In agreement with the relatively weak Ti–Cl bond (see Section 1), the reaction between $TiCl_4$ and substituted dioxolanes proceeds with ring opening and chlorination of the organic fragment [20].

1,3-Dioxolane and $CH_2(OEt)_2$ were reacted with the acetonitrile adduct $TiF_4(CH_3CN)_2$, which was considered more reactive than TiF_4 in view of its mononuclear structure [10]. We observed that



Scheme 2. Activation of CH₂(OMe)₂ by TiF₄.



Scheme 3. Activation of CH₂(OEt)₂ by TiF₄.



Scheme 4. Activation of CMe₂(OMe)₂ by TiF₄.

a low amount of acetonitrile (8% and 20%, respectively) was released after 20 days at room temperature or some hours at ca. 70 °C. No activation of the organic molecule was observed under the same experimental conditions used for the reactions with TiF₄.

In order to investigate in more detail the possibility of fluorinetransfer reactions, we allowed TiF₄ to react with compounds containing unsaturated functionalities such as propargylic alcohol



Scheme 5. Activation of CMe(OMe)₃ by TiF₄.

and the diethylacetal of phenylpropynal, $PhC \equiv CCH(OEt)_2$ (Table 1, run 8). No activation was observed in the first case while small amounts of EtOH were formed under drastic conditions in the latter reaction.

Even the three-membered ring of 1,2-epoxybutane was not activated by means of TiF_4 , and the probable formation of a coordination adduct could not be demonstrated (Table 1, run 9). The treatment of the reaction mixture with water caused the presumable metal-assisted quantitative conversion to 1,2-diol [22]. It has to be remarked that epoxides easily undergo ring-opening reactions in the presence of metal halides of Group 5, affording either halo-alkoxo or ketone complexes depending on the halide [23].

Titanium tetrafluoride reacted with ethyldiazoacetate at room temperature affording $CH_2FCOOEt$ as prevalent product, although in low yield (ca. 35%) (Table 1, run 10, Scheme 7).





Scheme 6. Formation of 1,4-dioxane from 1,3-dioxolane and TiF₄.

Scheme 7. Fluorine transfer to ethyldiazoacetate mediated by TiF₄.

The formation of ethylfluoroacetate is the result of the formal addition of HF to the fragment obtained by dinitrogen loss. It is noteworthy that, under comparable experimental conditions, the reaction of NbF₅ with N₂CHCO₂Et generated the stable coordination adduct NbF₅[$O=C(OEt)CHN_2$] [24].

3. Conclusions

Suspensions of TiF₄ in chlorinated solvents promote room temperature fragmentation and rearrangement of *O*-containing species, mainly 1,1-dialkoxyalkanes. The reactions are generally slow due to the polynuclear structure of TiF₄. The organic compounds identified after hydrolysis of the reaction mixtures suggest that both cleavage and formation of C–O, C–H and C–C bonds may occur during the reactions. Interestingly, small structural differences within the organic compound may determine strong differences in the nature of the products. A summary of the reaction products is reported in Table 1.

The possible formation of Ti-OR species, evidenced on the basis of the nature of the products, may represent an exception to the general observation that the strong Ti–F bond is not cleaved by oxygen containing species, including protic reagents like water and alcohols.

The presence of one unsaturated group adjacent to the *O*-functionality does not substantially influence the nature of the products. Fluorine transfer has been clearly observed only in the case of the reaction of TiF₄ with ethyldiazoacetate, yielding CH₂FCO₂Et.

4. Experimental

All manipulations of air and/or moisture sensitive compounds were performed under an atmosphere of pre-purified argon using standard Schlenk techniques. The reaction vessels were oven dried at 150 °C prior to use, evacuated (10^{-2} mm Hg) and then filled with argon. TiF₄ was purchased from Sigma Aldrich, stored in sealed tubes under argon as received, and used without further purification. TiF₄(CH₃CN)₂ was prepared according to the literature [10].

¹H NMR spectra were recorded on Bruker Avance DRX400 (BBFO probe) on CDCl₃ solutions at 298 K. The chemical shifts were referenced to the non-deuterated aliquot of the solvent.

GC/MS analyses were performed on a HP6890 instrument, interfaced with a MSD-HP5973 detector and equipped with a Phenonex Zebron column.

4.1. Reactions between TiF_4 and $TiF_4(CH_3CN)_2$ with O-donors: general procedure

Only the detailed procedure is described for TiF₄, that used with the acetonitrile adduct being substantially the same. TiF₄, CDCl₃ (0.5 mL), CH₂Cl₂ (CH₂Cl₂/Ti molar ratio = 1) and the organic reagent were introduced into a NMR tube. The tube was sealed, stirred to obtain a homogeneous suspension and stored at room temperature. After variable periods of time (see Table 2), NMR spectra were recorded. Then the mixture was heated at temperature as high as 80 °C. Thus the tube was cooled to ca. -30 °C and opened: the mixture was treated with water (ca. 1 mL). After 1 h stirring at room temperature, the resulting suspension was filtered and the solution was examined by NMR. Table 2 reports the experimental details of the reactions.

In the case of the reactions of $TiF_4(CH_3CN)_2$ with $CH_2(OEt)_2$ and 1,3-dioxolane, respectively 20% and 8% of acetonitrile was released into the solution after 20 days stirring at room temperature. In both cases, the organic reactant was the only product which could be detected after prolonged heating of the reaction mixture followed by hydrolysis.

Acknowledgments

The authors wish to thank the Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR, Roma), for financial support.

References

- (a) R.J.H. Clark, The Chemistry of Titanium and Vanadium, Elsevier, Amsterdam, 1968;
- (b) R. Colton, J.H. Canterford, Halides of the Transition Elements: Halides of the First Row Transition Metals, Wiley Interscience, London, 1969;
- (c) R.J. Clark, in: J.C. Bailar Jr., H.J. Emeleus, R.S. Nyholm, A.F. Trotman-Dickenson (Eds.), Comprehensive Inorganic Chemistry, Pergamon, Oxford, 1973, p. 355.
- [2] H. Bialowons, M. Müller, B.G. Müller, Z. Anorg. Allg. Chem. 621 (1995) 1227.
- [3] J.E. Huheey, E.A. Keiter, R.L. Keiter, Inorganic Chemistry: Principles of Structure and Reactivity, fourth ed., Harper and Collins, New York, 1993.
- [4] M. Jura, W. Levason, E. Petts, G. Reid, M. Webster, W. Zhang, Dalton Trans. 39 (2010) 10264.
- [5] E.G. Il'in, G.B. Nikiforov, G.G. Aleksandrov, Dokl. Chem. 396 (2004) 111.
- [6] (a) G.B. Nikiforov, S.G. Sakharov, E.G. Il'in, Yu.A. Buslaev, Russ. J. Inorg. Chem. 46 (2001) 1045;
 - (b) E.G. Il'in, G.B. Nikiforov, G.G. Aleksandrov, V.S. Sergienko, Yu.A. Buslaev, Dokl. Akad. Nauk 367 (1999) 772;
- (c) E.L. Muetterties, J. Am. Chem. Soc. 84 (1960) 1082.
- [7] D.S. Dyer, R.O. Ragsdale, Inorg. Chem. 6 (1967) 8.
- [8] (a) F.E. Dickson, E.W. Gowling, F.F. Bentley, Inorg. Chem. 6 (1967) 1099;
 (b) D.S. Dyer, R.O. Ragsdale, Inorg. Chem. 6 (1967) 1116;
 (c) C.E. Michelson, D.S. Dyer, R.O. Ragsdale, J. Chem. Soc. A (1970) 2296.
- [9] F. Marchetti, G. Pampaloni, Inorg. Chim. Acta 363 (2010) 3670.
- [10] G.B. Nikiforov, C. Knapp, J. Passmore, A. Decken, J. Fluorine Chem. 127 (2006) 1398.
- [11] E. Turin, R.M. Nielson, A.E. Merbach, Inorg. Chim. Acta 134 (1987) 67.
- [12] R.O. Ragsdale, B.B. Stewart, Inorg. Chem. 2 (1963) 1002.
- [13] (a) T.S. Cameron, A. Decken, E.G. Ilyin, G.B. Nikiforov, J. Passmore, Eur. J. Inorg. Chem. (2004) 3865;
 (b) N.N. Greenwood, A. Earnshaw, second ed., Elsevier, Amsterdam, 1997;
- (c) Yu.A. Buslaev, D.S. Dyer, R.O. Ragsdale, Inorg. Chem. 6 (1967) 2208. [14] (a) F. Fairbrother, The Halides of Niobium and Tantalum, in: V. Gutmann (Ed.),
- Halogen Chemistry, Academic Press, London, vol. 3, 2012, p. 123;
 (b) A.D. Beveridge, H.C. Clark, Pentahalides of Transition Metals, in: V. Gutmann (Ed.), Halogen Chemistry, Academic Press, London, vol. 3, 1967, p. 179.
- [15] S.J. Coles, M.B. Hursthouse, D.G. Kelly, A.J. Toner, N.M. Walker, J. Chem. Soc., Dalton Trans. (1988) 3489.
- [16] (a) F. Marchetti, G. Pampaloni, Chem. Commun. (2012) 635;
- (b) S. Dolci, F. Marchetti, G. Pampaloni, S. Zacchini, Dalton Trans. (2010) 5367; (c) S. Dolci, F. Marchetti, G. Pampaloni, S. Zacchini, Inorg. Chem. 50 (2011) 3846.
- [17] F. Marchetti, G. Pampaloni, S. Zacchini, Dalton Trans. (2009) 8096.
- 18] F. Marchetti, G. Pampaloni, S. Zacchini, Dalton Trans. (2008) 7026.
- [19] G. Balme, J. Gorè, J. Org. Chem. 48 (1983) 3336.
- [20] P. Mastagli, M. de Nanteuil, C. R. Acad. Sci. Paris 268 (1969) 1970.
- [21] (a) T.W. Greene, P.G.M. Wuts, Protecting Groups in Organic Synthesis, third ed., J. Wiley, New York, 1999 (Chapter 4, p. 293);
 (b) R.C. Larock, Comprehensive Organic Transformations, second ed., Wiley, VCH, New York, 1999;
 (c) M.B. Smith, J. March, March's Advanced Organic Chemistry, fifth ed., J. Wiley, New York, 2001.
- [22] (a) A. Gansäuer, A. Barchuk, F. Keller, M. Schmitt, S. Grimme, M. Gerenkamp, C. Mück-Lichtenfeld, K. Daasbjerg, H. Svith, J. Am. Chem. Soc. 129 (2007) 1359;
 (b) A.W. Eppley, N.I. Totah, Tetrahedron 53 (1997) 16545;
 (c) K. Fujiwara, T. Tokiwano, A. Murai, Tetrahedron Lett. 36 (1995) 8063;
 (d) M. Emziane, P. Lhoste, D. Sinou, J. Mol. Catal. 49 (1988) L23;
 (e) A.E. Vougioukas, H.B. Kagan, Tetrahedron Lett. 28 (1987) 6065.
- [23] F. Marchetti, G. Pampaloni, S. Zacchini, Polyhedron 28 (2009) 1235.
- [24] F. Marchetti, G. Pampaloni, S. Zacchini, Dalton Trans. (2008) 6759.