Domino reactions initiated by intramolecular hydride transfers from tri(di)arylmethane fragments to ketenimine and carbodiimide functions†‡

Mateo Alajarin,*^a Baltasar Bonillo,^a Maria-Mar Ortin,^a Pilar Sanchez-Andrada,^a Angel Vidal*^a and Raul-Angel Orenes^b

Received 1st June 2010, Accepted 14th July 2010 DOI: 10.1039/c0ob00193g

The ability of triarylmethane and diarylmethane fragments to behave as hydride donors participating in thermal [1,5]-H shift/ 6π -ERC tandem processes involving ketenimine and carbodiimide functions is disclosed. C-Alkyl-C-phenyl ketenimines N-substituted by a triarylmethane substructure convert into a variety of 3,3,4,4-tetrasubstituted-3,4-dihydroquinolines, as structurally related carbodiimides transform into 3,4,4-trisubstituted-3,4-dihydroquinazolines via transient ortho-azaxylylenes. The first step of these one-pot conversions, the [1,5]-H shift, is considered to be a hydride migration on the basis of the known hydricity of the tri(di)arylmethane fragment and the electrophilicity of the central heterocumulenic carbon atom, whereas the final electrocyclization involves the formation of a sterically congested C-C or C-N bond. In the cases of C,C-diphenyl substituted triarylmethane-ketenimines the usual 6π-ERC becomes prohibited by the presence of two phenyl rings at each end of the azatrienic system. This situation opens new reaction channels: (a) following the initial hydride shift, the tandem sequence continues with an alternative electrocyclization mode to give 9,10-dihydroacridines, (b) the full sequence is initiated by a rare 1,5 migration of an electron-rich aryl group, followed by a 6π -ERC which leads to 2-aryl-3,4-dihydroquinolines, or (c) a different [1,5]-H shift/ 6π -ERC sequence involving the initial migration of a hydrogen atom from a methyl group at the *ortho* position to the nitrogen atom of the ketenimine function. Diarylmethane-ketenimines bearing a methyl group at the benzylic carbon atom experience a tandem double [1,5]-H shift, the first one being the usual benzylic hydride transfer whereas the second one involves the methyl group at the initial benzylic carbon atom, the reaction products being 2-aminostyrenes. Diarylmethane-ketenimines lacking such a methyl group convert into 3,4-dihydroquinolines by the habitual tandem [1,5]-H shift/ 6π -ERC processes.

Introduction

C–H Bond functionalization, the direct and selective replacement of C–H bonds with new C–C or C–heteroatom bonds, is an important and long-standing goal in organic chemistry.¹ A series of transformations for C–H bond functionalization initiated by an intramolecular hydride transfer and followed by cyclization of the resulting intermediates have been recently disclosed.² Such hydride shift/cyclization sequences, serving for the efficient buildup of complex polycyclic systems, are in some sense related to those based upon the well-known *tert*-amino effect.³ In this context, while investigating new reactions of nitrogenated heterocumulenes, we recently found that 1,3-dioxolane-ketenimines 1 (X = O; Y = CR²R³), 1,3-dioxolane-carbodiimides 1 (X = O; Y = NAr) and 1,3-dithiolane-ketenimines 1 (X = S; Y = CR^2R^3) transform, under mild thermal treatment, into spiroquinolines 3 (X = O,S; $Y = CR^2R^3$) and spiroquinazolines 3 (X = O; Y = NAr), via a tandem sequence consisting of a [1,5]-H shift followed by a 6π electrocyclic ring closure (6π -ERC) (Scheme 1).⁴ On the other hand, when 1,3-oxathiolane-ketenimines 4 (Y = CR^2R^3) and 1,3-oxathiolane-carbodiimides 4 (Y = NAr) were submitted to mild heating the 2,1-benzisothiazol-3-ones 6 were formed, a transformation which seems to occur through a formal [1,5]-H shift/1,5 electrocyclization/[3+2] cycloreversion tandem process, with the concomitant formation of ethylene (Scheme 1).⁵ The [1,5]-H shifts from the acetalic functions to the electrophilic central carbon atom of the ketenimine or carbodiimide moieties, presumably providing the transient ortho-azaxylylenes 2 and 5, were characterized as intramolecular hydride transfers by means of computational DFT studies. In fact, the computed NBO analysis of heterocumulenes 1 and 4 shows the weakening and polarization of the acetalic C-H bond by hyperconjugative interactions of its σ^* C–H orbital with the lone pair electrons at the vicinal heteroatoms. These results demonstrated the hydride donor ability (hydricity) of the 2-monosubstituted 1,3-dioxolane, 1,3-oxathiolane and 1,3-dithiolane functions. Additionally, the experimental work and the computational calculations showed that the hydricity-imparting character of these acetalic functions decreases in the order: 1,3-dioxolane > 1,3-oxathiolane > 1.3-dithiolane.

^aDepartamento de Quimica Organica, Facultad de Quimica, Universidad de Murcia, Campus de Espinardo, 30100, Murcia, Spain. E-mail: alajarin@ um.es, vidal@um.es; Fax: + 34-868 364149; Tel: + 34-868887497, + 34-868887418

^bServicio Universitario de Instrumentacion Científica, Universidad de Murcia, Campus de Espinardo, 30100, Murcia, Spain. E-mail: raorenes@um.es † Dedicated to Professor José Elguero with respect and admiration.

[‡] Electronic supplementary information (ESI) available: Comprehensive experimental section. ¹H and ¹³C NMR spectra of compounds **11**, **13**, **18**, **20**, **22**, **25**, **27**, **30**, **32** and **34**. ³¹P NMR spectra of compounds **11**, **27** and **32**. Cif files of **13e**, **13k**, **18a** and **20b**. CCDC reference numbers 779905– 779908. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c0ob00193g



Scheme 1 Proposed mechanisms for the conversions $1 \rightarrow 3$ and $4 \rightarrow 6$.

The main class of compounds possessing hydricity is the metal hydrides.⁶ Nevertheless several types of organic compounds are also considered to exhibit this property. For example, NADH model compounds,⁷ triarylmethanes⁸ and some 1,3-dinitrogenated heterocyclic systems, such as *N*,*N*dimethylbenzimidazoles,⁹ 2-benzoyl-*N*,*N*-dimethylperhydropyrimidine¹⁰ and orthoformamides,¹¹ are generally appreciated as good hydride donors.

Based on the known hydricity of the triarylmethanes, we targeted the preparation of a series of these compounds bearing a ketenimine or carbodiimide function linked by its nitrogen atom to the *ortho* position of one of the three aryl groups (Fig. 1). Our aim was to examine the viability of the [1,5]-H transfer from the sp³ methine carbon to the electron-deficient central carbon atom of the ketenimine or carbodiimide fragment, and therefore the participation of this type of heterocumulene in



Y = CR³Ph, NA

Fig. 1 General structure of triarylmethane-ketenimines and triarylmethane-carbodiimides.

[1,5]-H shift/ 6π -ERC tandem processes closely related to those described for compounds 1. As result of this experimental work we here disclose that such triarylmethane-ketenimines converted under thermal conditions into substituted quinolines or acridines through tandem processes that involve as the first mechanistic step either the expected [1,5]-H transfer or an unusual [1,5] sigmatropic migration of an aryl group, and as the second step different modes of electrocyclic ring closure in the resulting 3-azahexatriene intermediates. The electrocyclization mode at this second step seems to depend on the nature of the substituents at the aryl rings (R¹ and R²) and at the *C*-terminal carbon atom of the starting ketenimine function (R³). In addition, we report that under similar conditions triarylmethane-carbodiimides simply undergo 1,5 migration of the hydrogen at the sp³ methine carbon atom and subsequent 6π -ERC to afford quinazolines.

Results and discussion

The synthetic strategy for the preparation of the triarylmethaneketenimines and triarylmethane-carbodiimides starts with a Friedel-Crafts acylation in which two equivalents of benzene or a monosubstituted benzene react with one equivalent of 2-nitrobenzaldehyde 7a or 3-methyl-2-nitrobenzaldehyde 7b, to provide nitrotriarylmethanes 8 (36-73%). These reactions were run under different conditions depending on the nature of R². In the cases where $\mathbf{R}^2 = \mathbf{H} (\mathbf{8a,b})$ the acylation reaction was carried out by treating benzene solutions of the 2-nitrobenzaldehydes 7 with anhydrous aluminium chloride.¹² When $R^2 = N(CH_3)_2$ (8c,d) or N-morpholino (8e) mixtures of the 2-nitrobenzaldehydes 7 and N,N-dimethylaniline or N-phenylmorpholine were treated with anhydrous zinc chloride.13 The synthesis of the nitro derivatives **8f** and **8g** in which $R^2 = OCH_3$ was achieved by reaction of the 2-nitrobenzaldehydes 7 with anisole in the presence of aluminium chloride.¹⁴ Next, the reduction of compounds 8 with molecular hydrogen in the presence of Pd on activated charcoal, using ethanol as solvent, yielded the anilines 9 (65-88%). The diazotation of anilines 9 followed by the addition of sodium azide provided azides 10. For this latter conversion different reaction conditions were also used in accordance with the nature of R^2 . For $R^2 = H$ or OCH₃ the corresponding azides **10a,b,f,g** were readily obtained (80–92%) by sequential treatment of sulfuric acid/1,4-dioxane solutions of amines 9a,b,f,g with sodium nitrite and sodium azide.13 For the cases where $R^2 = N(CH_3)_2$ or *N*-morpholino suspensions of the amines 9c-e in 6 N hydrochloric acid were first treated with a solution of sodium nitrite in water, and then with an aqueous solution of sodium azide and sodium acetate (45-72%).¹⁵ The imination reaction of triphenylphosphine with the azides 10, in anhydrous diethyl ether at room temperature, provided the triphenyliminophosphoranes 11 (86–99%) (Scheme 2).

The aza-Wittig reaction of iminophosphoranes 11 with ethylphenylketene and methylphenylketene, in dichloromethane solution at room temperature, led to the formation of the triarylmethane-ketenimines 12 (Scheme 3, Table 1), which were separated from the triphenylphosphine oxide by column chromatography on a short pad of silica gel, checked by IR (strong absorption near 2000 cm⁻¹) and immediately submitted to the next reaction step for preventing its partial hydrolysis.¹⁶ It is relevant to note that compounds 12 bear a phenyl group and an ethyl or methyl substituent at the sp² terminal carbon atom of the

15

99

61

63

80

78

75

65

60

57

52

85



Scheme 2 Reagents and conditions: (a) $R^2 = H C_6 H_6$, AlCl₃, 60 °C, 6 h; $R^2 = N(CH_3)_2$, *N*-morpholino $C_6 H_3 R^2$, ZnCl₂, 80 °C, 6 h; $R^2 = OCH_3$ $C_6 H_3 OCH_3$, AlCl₃, r.t., 12 h (36–73%). (b) H₂, Pd/C, EtOH, r.t., 24 h (65–88%). (c) $R^2 = H$, OCH₃ (1) NaNO₂, H₂SO₄ (4 N)/1,4-dioxane, 0 °C, 30 min (2) NaN₃, H₂SO₄ (4 N)/1,4-dioxane, r.t., 16 h; $R^2 = N(CH_3)_2$, *N*-morpholino (1) NaNO₂, HCl/H₂O, 0 °C, 30 min (2) NaN₃, NaOAc, HCl/H₂O, r.t., 1 h (45–92%). (d) PPh₃, Et₂O, r.t., 16 h (86–99%).

ketenimine function. When toluene or *ortho*-xylene solutions of ketenimines **12** were heated at reflux temperature, during periods of time ranging from 5 to 144 h, the 3-ethyl(methyl)-3,4,4-triaryl-3,4-dihydroquinolines **13** were formed, and isolated in moderate to good yields (Scheme 3, Table 1).

The structural determination of the 3-alkyl-3,4,4-triaryl-3,4dihydroquinolines **13** was carried out following their analytical and spectral data. In their ¹H NMR spectra the C(2)H proton resonates as a singlet at $\delta = 8.21-8.68$ ppm. Relevant ¹³C NMR data of compounds **13** are the chemical shifts of the quaternary carbon C3 at $\delta = 48.0-52.0$ ppm, the quaternary carbon C4 at $\delta = 59.7-$ 62.9 ppm and the methine C2 at $\delta = 166.2-169.4$ ppm. Moreover, the structures of compounds **13e** (R¹ = H; R² = N(CH₃)₂; R³ = CH₃) and **13k** (R¹ = H; R² = OCH₃; R³ = CH₃) were determined by Xray crystal diffraction (see Electronic supplementary information). The main feature of these two structures is the relatively long bond distance of the C3–C4 bond (C8–C9 bond in the crystallographic study), 1.598 Å in compound **13e** and 1.604 Å in **13k**. These long distances are most probably due to the steric repulsion between the bulky substituents at the C3 and C4 carbon atoms.¹⁷

 Image: new particular
 Image: new particular

CH₂CH₃

CH₂CH₃

CH₂CH₃

CH₂CH₃

CH₂CH₃

CH₂CH₃

CH₃

CH₂

CH₃

CH.

CH-

CH₃

80

40

38

92

71

90

67 79

97

74

86

70

Ketenimines 12 and Quinolines 13

Table 1

b

c

d

e

f

g h

i

i

k

1

m

Н

н

Η

CH₃

CH,

н

Η

Η

Η

CH₃

CH₃

CH₃

Н

Η

Η

 $N(CH_3)_2$

 $N(CH_3)_2$

 $N(CH_3)_2$

 $N(CH_3)_2$

OCH₂

OCH₃

OCH₃

N-morpholino

N-morpholino



Scheme 3 *Reagents and conditions*: (a) PhR³CCO, CH₂Cl₂, r.t., 30 min. (b) Toluene or *ortho*-xylene, reflux, 5–144 h.

The transformation $12 \rightarrow 13$ should proceed by a mechanism involving as the first step the cleavage of the C-H bond of the triarylmethane fragment *via* a 1,5 hydride shift to give the transient *ortho*-azaxylylene 14 (Scheme 4). Subsequent 6π electrocyclic ring closure should afford the sterically congested 3,4-dihydroquinolines 13. The denotation of the first mechanistic step as a hydride shift relies on the known hydricity of the triarylmethane C-H bond and the electrophilicity of the central ketenimine carbon atom.

The presence of a methyl substituent at the *ortho* position to the keteniminic nitrogen atom ($R^1 = CH_3$) has a significant influence on the reactivity of the triarylmethane-ketenimines 12: in general, compounds 12 with $R^1 = CH_3$ afforded the corresponding 3,4-dihydroquinolines 13 in the best chemical yields and in the shortest reaction times. This enhancement of reactivity could be rationalized by considering that the presence of this methyl group shifts the conformational equilibrium of compounds 12 to the

Ρĥ



Scheme 4 Proposed mechanism for the conversion $12 \rightarrow 13$.

Carvi-N rotational isomer that locates the electrophilic central carbon atom of the ketenimine function close to the hydrogen atom that is going to translate in the 1,5 hydride migration.¹⁸ In addition, we have observed that the presence of strong electronreleasing groups ($R^2 = N(CH_3)_2$, N-morpholino and OCH₃) at the para position of the two other aryl rings facilitates the conversion of ketenimines 12 by shortening the reaction time. These groups are known to increase the hydricity of the triarylmethane C-H bond.8 In most of the cases, placing an ethyl group instead of a methyl one at the terminal carbon atom of the ketenimine function results in a lower yield of the tandem process converting ketenimines 12 into quinolines 13, a fact that could be attributed to the slightly higher steric hindrance that makes more difficult the electrocyclization step.

We next approached the study of similar tandem processes in which the future C3–C4 bond of the putative final quinolines is more sterically congested by the presence of four aryl substituents at the extremes of that bond. To this end we prepared the C,Cdiphenyl ketenimines 15 (60–92%) by reacting iminophosphoranes 11a-d and 11f,g with diphenylketene, in dichloromethane solution at room temperature (Scheme 5).

When toluene solutions of ketenimines 15 were heated at reflux temperature for several days these heterocumulenes remained unaltered. Under more severe thermal conditions (ortho-xylene, 180 °C, sealed tube) and longer reaction times we were able to achieve the transformation of 15 into different heterocyclic compounds depending on the nature of the substituents R¹ and \mathbf{R}^2 .

Thus, the thermal treatment of C,C-diphenyl ketenimines 15a $(R^1 = R^2 = H)$ and 15f $(R^1 = H; R^2 = OCH_3)$ under such reaction conditions (5-7 days) yielded 9,10-dihydroacridines 18 (Scheme 6). The structural determination of compounds 18 was achieved following their spectral data, and confirmed by X-ray diffraction of a monocrystal of compound 18a ($R^1 = R^2 = H$) (see Electronic supplementary information).



18b R² = OCH₃ (61%)

Scheme 6 Reagents and conditions: (a) ortho-Xylene, 180 °C, sealed tube, 5–7 d.

The formation of dihydroacridines 18 could be explained by assuming the initial formation of the ortho-azaxylylene intermediates 16 via a 1,5 hydride shift. The further cyclization of these intermediates through a 6π electrocyclic ring closure involving the N3-C4-C5-C6-C7-C8 1-azatriene fragment should afford the final species 18. The alternative 6π -electrocyclization of the ortho-azaxylylenes 16 that would involve the closure of the C1–C2–N3–C4–C5–C6 3-azatriene system *via* C1–C6 bond formation, that was initially expected and is similar to those presented above, is probably made difficult by the presence of two phenyl rings at each end (C1 and C6) of the azatrienic system, which should result in a severe steric congestion at the transition state of such an electrocyclization thus preventing its occurrence.¹⁹

In the cases of the C, C-diphenyl ketenimines **15c** and **15d**, where the R² substituent is a dimethylamino group, the reaction products obtained from their thermal treatment were the 2-aryl substituted 3,4-dihydroquinolines **20** (Scheme 7).



Scheme 7 Reagents and conditions: (a) ortho-Xylene, $180 \degree C$, sealed tube, $1-2 \ d$.

Quinolines **20** were characterized following their spectral data, IR, ¹H and ¹³C NMR and mass spectrometry, and additionally by an X-ray crystal structure of example **20b** ($R^1 = CH_3$) (see Electronic supplementary information). In the solid state the C3– C4 bond distance of compound **20b** (C7–C8 in the crystallographic numbering) is also larger (1.588 Å) than a standard Csp³–Csp³ bond, and slightly shorter than those shown by quinolines **13e** and **13k**.

The formation of compounds **20** is apparently explained as occurring by a tandem process in which the first step is the rare 1,5 migration of one electron-rich 4-dimethylaminophenyl group²⁰ to the central carbon atom of the ketenimine function leading to the transient *ortho*-azaxylylene **19**, followed by the habitual 6π -ERC process.

C,*C*-Diphenyl ketenimines **15b** ($\mathbf{R}^1 = \mathbf{CH}_3$; $\mathbf{R}^2 = \mathbf{H}$) and **15g** ($\mathbf{R}^1 = \mathbf{CH}_3$; $\mathbf{R}^2 = \mathbf{OCH}_3$), both bearing a methyl group in the *ortho* position to the keteniminic nitrogen atom ($\mathbf{R}^1 = \mathbf{CH}_3$), when heated under the standard reaction conditions for extended periods of time (14–21 days), converted into the new 4-unsubstituted 3,4-dihydroquinolines **22a** ($\mathbf{R}^2 = \mathbf{H}$) and **22b** ($\mathbf{R}^2 = \mathbf{OCH}_3$), respectively

(Scheme 8). The spectral data of quinolines **22** are fully consistent with their proposed structures.



Scheme 8 *Reagents and conditions:* (a) *ortho-Xylene, 180* °C, sealed tube, 14–21 d.

The conversions $15 \rightarrow 22$ should be initiated by a 1,5 migration of one of the hydrogen atoms of the methyl group to give the transient ortho-azaxylylenes 21, which undergo subsequent 6π electrocyclization to quinolines 22 (Scheme 8). Amongst the various [1,5]-H shift/6π-ERC tandem processes that could undergo ketenimines 15b and 15g the one leading to 22 became the most viable reaction pathway, probably because it involves a facile 6π -electrocyclization step *via* a non congested transition state as the 3-azatriene intermediate experiencing the cyclization is unsubstituted at one of its carbon termini. Although the alternative [1,5]-H shift/ 6π -ERC sequences initiated by the [1,5]-H transfer from the methine carbon atom of the triarylmethane fragment should reasonably have a quite lower activation energy for this first step, when compared with the [1,5]-H shift from the methyl group, the further putative electrocyclizations must have notably higher activation barriers, instead.

The ability of triarylmethane fragments as hydride donors participating in [1,5]-H shift/ 6π -ERC tandem processes involving carbodiimide functions as the acceptor units was also evaluated. The reaction of iminophosphoranes **11c,d,f,g** with aryliso-cyanates, in dichloromethane solution at room temperature, gave triarylmethane-carbodiimides **23**. The heating at 180 °C in a sealed tube of *ortho*-xylene solutions of carbodiimides **23**, during periods of time ranging from 2 to 5 days, yielded the 3,4,4-triaryl-3,4-dihydroquinazolines **25** (Scheme 9, Table 2).

The structural characterization of quinazolines 25 relies on their spectroscopic data. Their IR spectra show an absorption band at $1606-1614 \text{ cm}^{-1}$ attributed to the vibration of the N–C bond. In

Table 2 Triarylmethane-carbodiimides 23 and quinazolines 25



Scheme 9 Reagents and conditions: (a) $4 \cdot R^3 \cdot C_6 H_4 NCO$, $CH_2 Cl_2$, r.t., 30 min. (b) ortho-Xylene, 180 °C, sealed tube, 2–5 d.

their ¹H NMR spectra the signal of the C(2)H proton appears as a singlet at $\delta = 7.64-7.73$ ppm. Their ¹³C NMR spectra show the signals of the aliphatic quaternary carbon atom C4 and the methine carbon C2 at $\delta = 71.8-71.9$ ppm and $\delta = 146.3-147.4$ ppm, respectively.

Obviously, the formation of the dihydroquinazolines **25** from carbodiimides **23** can be rationalized as resulting from a [1,5]-H shift/ 6π -ERC sequence that involves the diazatriene intermediates **24**.

It is worth noting that the thermal activation of triarylmethanecarbodiimides 23 yields a single type of products, the quinazolines 25, in contrast with the variety of heterocyclic compounds formed in the thermal cyclization of *C*,*C*-diphenyl ketenimines 15. This difference is explained by the relatively easy electrocyclization of the 1,3-diazatriene fragment of intermediates 24, which should occur through a less sterically congested transition state than the similar process of the C1–C2–N3–C4–C5–C6 3-azatriene fragment in species 16. Moreover, it is conceivable that the lone pair at the terminal nitrogen atom of the 1,3-diazatriene moiety of 24 may also assist the 6π -ERC step by conferring a certain degree of pseudopericyclic character to its transition state.²¹

At this point, we were also interested in exploring whether the C–H bond of a diarylmethane fragment would be similarly activated as a formal hydride donor in related tandem processes. With this aim and following the usual sequence of reactions, as shown in Scheme 10, we prepared diarylmethane-ketenimines **28**



and **33a,b**, the first one bearing a methyl group at the benzylic carbon atom, and the second one lacking this substituent.

The heating of ketenimine **28** in refluxing toluene provided the substituted 2-aminostyrene **30**, isolated as the unique reaction product in 91% yield. In contrast, the thermal activation of ketenimines **33** in toluene solution at 160 °C, in a sealed tube, led to the quinolines **34** (Scheme 10). Quinoline **34b**, in which \mathbb{R}^3 is a methyl group was obtained as a pair of *cis* and *trans* diastereoisomers in a ratio close to 1:1.5.

The conversion $28 \rightarrow 30$ probably proceeds by a tandem double [1,5]-H shift, the first one being the usual benzylic hydride transfer

whereas the second one involves the migration of one of the methyl protons towards the nitrogen atom of the intermediate orthoazaxylylene 29. This result demonstrated the ability of -(Ar)₂C-H hydrogen atoms to engage in intramolecular hydride shifts but also reveals a limitation of the tandem [1,5]-H shift/6π-ERC methodology here developed for the preparation of polynuclear heterocycles: a primary (or secondary) alkyl group placed at the benzylic carbon atom from which the hydride departs in the first step interrupts the habitual tandem sequence at the 6π -ERC stage, as their H atoms become involved in a competitive, and clearly more successful, [1,5]-H shift for the final recovery of the aromaticity at the benzene nucleus of the ortho-azaxylylene. Notwithstanding, in the absence of such an alkyl group the habitual tandem [1,5]-H shift/ 6π -ERC sequence is fully operative when starting from diarylmethane hydride-donor fragments, as the transformation of ketenimines 33 into quinolines 34 clearly proves.

Conclusions

We here disclosed a novel synthetic protocol for converting C-alkyl-C-phenyl ketenimines N-substituted by triarylmethane substructures into a variety of 3,3,4,4-tetrasubstituted-3,4-dihydroquinolines, which is also valid for transforming structurally related carbodiimides into 3,4,4-trisubstituted-3,4dihydroquinazolines. This protocol is based on the hydride donor ability of the triarylmethane C-H fragment which, by thermal activation in solution, promotes a tandem [1,5]-H shift/ 6π -ERC sequence leading from the heterocumulenes to the heterocyclic products. In the first step, a hydrogen atom translates from the benzylic position to the central carbon atom of the heterocumulenic function. The last mechanistic step, the 6π electrocyclization, is also noteworthy as it implies the formation of sterically congested C-C or C-N bonds. Only when the new putative C-C bond would be tetraaryl substituted the usual 6π -ERC is prevented and the tandem sequence continues with a second 6π -electrocyclization mode leading to 9,10-dihydroacridines. Alternatively, the full sequence either becomes a new tandem process of the type [1,5]-Ar shift/ 6π -ERC which leads to 2-aryl substituted 3,4-dihydroquinolines initiated by the rare migration of an electron-rich aryl group, or is replaced by another [1,5]-H shift/ 6π -ERC process involving the initial migration of a hydrogen atom from the methyl group ortho to the N atom of the ketenimine. We have also demonstrated the ability of diarylmethane fragments for promoting similar transformations and the interruption of the final electrocyclization in the cases where a methyl group is placed at the benzylic H-donor carbon atom. New investigations are currently underway in our laboratories aimed to extend the applicability of this synthetic methodology by replacing the acceptor heterocumulene units by other functions bearing electrophilic carbon atoms.

Experimental

All melting points are uncorrected. Infrared (IR) spectra were recorded neat or as Nujol emulsions. ¹H NMR spectra were recorded in CDCl₃ at 300 or 400 MHz. ¹³C NMR spectra were recorded in CDCl₃ at 75 or 100 MHz. The chemical shifts are expressed in ppm, relative to Me₄Si at $\delta = 0.00$ ppm for ¹H, while

the chemical shifts for ¹³C are reported relative to the resonance of CDCl₃ δ = 77.1 ppm. *J* values are given in Hz.

Preparation of the quinolines 13

To a solution of iminophosphorane **11** (1 mmol) in anhydrous dichloromethane (15 mL) ethylphenylketene (0.15 g, 1 mmol) or methylphenylketene (0.13 g, 1 mmol) in the same solvent (5 mL) was added. The reaction mixture was stirred at room temperature for 30 min. Then, the solvent was removed under reduced pressure and the resulting crude material was chromatographed on a silica gel column, using hexanes/diethyl ether (9:1, v/v) as eluent, to give the corresponding ketenimine **12**.

A solution of the ketenimine 12(0.5 mmol) in anhydrous toluene (20 mL) was heated at reflux temperature under an atmosphere of nitrogen for 5–144 h. After cooling at room temperature, the solvent was removed under reduced pressure and the resulting material was purified by column chromatography on silica gel.

3-Methyl-3,4,4-triphenyl-3,4-dihydroquinoline 13a. Yield = 0.11 g, 60%; mp 194–195 °C (colourless prisms from diethyl ether–hexane); v_{max} (Nujol)/cm⁻¹ 1633; δ_{H} (300 MHz; CDCl₃) 1.83 (3 H, s), 6.61 (2 H, d, *J* 7.5), 6.90 (4 H, t, *J* 7.2), 6.98–7.06 (4 H, m), 7.10–7.21 (6 H, m), 7.45 (1 H, dd, *J* 7.5 and 1.2), 7.55–7.58 (2 H, m), 8.21 (1 H, s); δ_{C} (75 MHz; CDCl₃) 20.2, 48.5 (s), 61.4 (s), 126.2, 126.3, 126.6, 126.7, 127.2, 127.5, 127.6, 127.9, 128.4, 128.6, 129.2, 131.8, 135.1 (s), 138.2 (s), 140.8 (s), 141.7 (s), 146.1 (s), 167.8; HRMS (ESI): Calcd for C₂₈H₂₄N [M + H]⁺ 374.1903, found 374.1907.

3-Ethyl-3,4,4-triphenyl-3,4-dihydroquinoline 13b. Yield = 0.03 g, 15%; yellow oil; v_{max} (neat)/cm⁻¹ 1625; δ_{H} (300 MHz; CDCl₃) 0.80 (3 H, t, *J* 7.2), 2.00–2.10 (1 H, m), 2.53–2.62 (1 H, m), 6.61 (2 H, d, *J* 7.6), 6.93–7.25 (14 H, m), 7.41–7.46 (3 H, m), 8.63 (1 H, s); δ_{C} (75 MHz; CDCl₃) 8.8, 24.8, 51.6 (s), 62.6 (s), 126.5, 126.6, 126.7, 126.8, 127.2, 127.3, 127.4, 127.8, 128.3, 129.0, 130.4, 130.5, 131.7, 135.2 (s), 135.3 (s), 141.8 (s), 142.3 (s), 144.0 (s), 167.7; HRMS (ESI): Calcd for C₂₉H₂₆N [M + H]⁺ 388.2060, found 388.2066.

3,8-Dimethyl-3,4,4-triphenyl-3,4-dihydroquinoline 13c. Yield = 0.19 g, 99%; mp 195–196 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1633; δ_{H} (400 MHz; CDCl₃) 1.81 (3 H, s), 2.56 (3 H, s), 6.65 (2 H, d, *J* 10.0), 6.86–7.04 (10 H, m), 7.09–7.24 (4 H, m), 7.54–7.57 (2 H, m), 8.25 (1 H, s); δ_{c} (100 MHz; CDCl₃) 18.5, 20.6, 48.0 (s), 61.8 (s), 126.3, 126.6, 126.7, 127.1, 127.4, 127.5, 128.0, 128.8, 128.9, 132.1, 135.0 (s), 135.9 (s), 138.5 (s), 140.1 (s), 141.4 (s), 146.2 (s), 167.4; HRMS (ESI): Calcd for C₂₉H₂₆N [M + H]⁺ 388.2060, found 388.2066.

3-Ethyl-8-methyl-3,4,4-triphenyl-3,4-dihydroquinoline Yield = 0.12 g, 61%; mp 145–147 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1628; δ_{H} (400 MHz; CDCl₃) 0.78 (3 H, t, *J* 9.6), 1.96–2.08 (1 H, m), 2.54 (3 H, s), 2.58–2.63 (1 H, m), 6.57 (2 H, d, *J* 9.0), 6.91–7.20 (14 H, m), 7.43 (2 H, d, *J* 9.0), 8.68 (1 H, s); δ_{C} (100 MHz; CDCl₃) 8.80, 18.4, 24.9, 50.9 (s), 62.9 (s), 126.4, 126.5, 126.7, 126.8, 126.9, 127.1, 127.2, 127.3, 129.0, 130.4, 130.6, 131.9, 134.9 (s), 135.5 (s), 140.6 (s), 142.0 (s), 144.2 (s), 166.2; HRMS (ESI): Calcd for C₃₀H₂₈N [M + H]⁺ 402.2216, found 402.2219.

4,4-Bis[4-(*N*,*N*-dimethylamino)phenyl]-3-methyl-3-phenyl-3,4dihydroquinoline 13e. Yield = 0.14 g, 63%; mp 190–191 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1613;
$$\begin{split} &\delta_{\rm H}(300~{\rm MHz};~{\rm CDCl_3})~1.77~(3~{\rm H},~{\rm s}),~2.90~(12~{\rm H},~{\rm s}),~6.37~(2~{\rm H},~{\rm d},~J~9.0),~6.50~(2~{\rm H},~{\rm d},~J~9.0),~6.65~(2~{\rm H},~{\rm d},~J~8.1),~6.72~(2~{\rm H},~{\rm d},~J~8.1),~6.93~(2~{\rm H},~{\rm d},~J~9.0),~6.65~(2~{\rm H},~{\rm d},~J~8.1),~6.72~(2~{\rm H},~{\rm d},~J~8.1),~6.93~(2~{\rm H},~{\rm t},~J~7.5),~6.99-7.04~(2~{\rm H},~{\rm m}),~7.10-7.20~(2~{\rm H},~{\rm m}),~7.37-7.42~(3~{\rm H},~{\rm m}),~8.23~(1~{\rm H},~{\rm s});~\delta_{\rm C}(75~{\rm MHz};~{\rm CDCl_3})~20.6,~40.3,~40.7,~49.0~({\rm s}),~59.9~({\rm s}),~110.5,~111.2,~126.4,~126.6,~127.3,~127.8,~128.2,~129.0,~129.8,~132.6,~134.0~({\rm s}),~136.6~({\rm s}),~139.1~({\rm s}),~141.8~({\rm s}),~148.8~({\rm s}),~148.9~({\rm s}),~169.4~({\rm s});~{\rm HRMS}~({\rm ESI}):~{\rm Calcd}~{\rm for}~{\rm C}_{\rm {32}}{\rm H}_{\rm {34}}{\rm N}_{\rm 3}~[{\rm M}~{\rm H}]^+~460.2747,~{\rm found}~460.2751. \end{split}$$

4,4-Bis[4-(*N*,*N***-dimethylamino) phenyl]-3-ethyl-3-phenyl-3,4-dihydroquinoline 13f.** Yield = 0.19 g, 80%; mp 171–172 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1610; $\delta_{\rm H}$ (400 MHz; CDCl₃) 0.74 (3 H, t, *J* 7.3), 1.98–2.07 (1 H, m), 2.45–2.54 (1 H, m), 2.85 (6 H, s), 2.91 (6 H, s), 6.37 (2 H, d, *J* 9.1), 6.53 (2 H, d, *J* 9.1), 6.70 (2 H, d, *J* 7.7), 6.76 (2 H, d, *J* 7.7), 7.00–7.09 (4 H, m), 7.14–7.20 (2 H, m), 7.24–7.26 (2 H, m), 7.40 (1 H, dd, *J* 7.6 and 1.3), 8.63 (1 H, s); $\delta_{\rm C}$ (100 MHz; CDCl₃) 9.0, 25.6, 40.4, 40.5, 52.0 (s), 60.9 (s), 110.5, 110.8, 126.4, 126.7, 127.1, 127.5, 128.0, 128.9, 129.9 (s), 130.6, 131.3 (s), 131.7, 132.2, 136.4 (s), 136.5 (s), 142.3 (s), 148.9 (s), 149.0 (s), 168.5; HRMS (ESI): Calcd for C₃₃H₃₆N₃ [M + H]⁺ 474.2904, found 474.2910.

4,4-Bis[4-(*N*, *N*-dimethylamino)phenyl]-3,8-dimethyl-3-phenyl-3,4-dihydroquinoline 13g. Yield = 0.18 g, 78%; mp 170–171 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1610; $\delta_{\rm H}$ (400 MHz; CDCl₃) 1.75 (3 H, s), 2.54 (3 H, s), 2.87 (6 H, s), 2.88 (6 H, s), 6.36 (2 H, d, *J* 8.8), 6.48–6.51 (2 H, m), 6.62 (2 H, d, *J* 7.6), 6.71 (2 H, d, *J* 8.8), 6.88–7.04 (6 H, m), 7.37–7.39 (2 H, m), 8.27 (1 H, s); $\delta_{\rm C}$ (100 MHz; CDCl₃) 18.5, 20.9, 40.4, 40.7, 48.3 (s), 60.3 (s), 110.5, 111.2, 126.4, 126.8, 127.1, 127.3, 128.4, 129.1, 129.6 (s), 130.1, 132.7, 134.0 (s), 135.4 (s), 136.4 (s), 139.3 (s), 140.1 (s), 148.8 (s), 148.9 (s), 167.9; HRMS (ESI): Calcd for C₃₃H₃₆N₃ [M + H]⁺ 474.2904, found 474.2909.

4,4-Bis[4-(*N*,*N***-dimethylamino)phenyl]-3-ethyl-8-methyl-3-phenyl-3,4-dihydroquinoline 13h.** Yield = 0.18 g, 75%; mp 165–167 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1610; $\delta_{\rm H}$ (300 MHz; CDCl₃) 0.72 (3 H, t, *J* 7.2), 1.94–2.06 (1 H, m), 2.45–2.50 (1 H, m), 2.53 (3 H, s), 2.86 (6 H, s), 2.92 (6 H, s), 6.36 (2 H, d, *J* 9.0), 6.53 (2 H, d, *J* 9.0), 6.67 (2 H, d, *J* 8.1), 6.76 (2 H, d, *J* 8.1), 6.91–7.08 (6 H, m), 7.23–7.26 (2 H, m), 8.68 (1 H, s); $\delta_{\rm C}$ (75 MHz; CDCl₃) 15.4, 18.4, 25.7, 40.5, 40.6, 51.4 (s), 61.2 (s), 110.4, 110.7, 126.3, 126.7, 126.9, 127.1, 128.4, 130.2 (s), 130.6, 131.7 (s), 131.8, 132.4, 135.1 (s), 136.2 (s), 136.8 (s), 140.6 (s), 148.9 (s), 149.0 (s), 167.0; HRMS (ESI): Calcd for C₃₄H₃₈N₃ [M + H]⁺ 488.3060, found 488.3065.

3-Methyl-4,4-bis[**4-**(*N*-morpholino)phenyl]-**3**-phenyl-**3**,**4**-dihydroquinoline 13i. Yield = 0.18 g, 65%; mp 221–222 °C (orange prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1608 and 1512; $\delta_{\rm H}$ (400 MHz; CDCl₃) 1.79 (3 H, s), 3.06–3.09 (8 H, m), 3.79–3.85 (8 H, m), 6.55 (2 H, d, *J* 8.4), 6.65–6.67 (4 H, m), 6.78 (2 H, d, *J* 8.4), 6.93 (2 H, t, *J* 7.6), 7.03 (2 H, t, *J* 8.4), 7.17 (2 H, t, *J* 7.6), 7.45 (3 H, m), 8.23 (1 H, s); $\delta_{\rm C}$ (100 MHz; CDCl₃) 20.1, 48.2, 48.4 (s), 48.8, 59.7 (s), 66.4, 66.5, 112.9, 113.6, 126.2, 126.5, 127.0, 127.5, 127.9, 128.3 (s), 128.4, 128.5, 129.3, 131.8 (s), 132.2, 135.6 (s), 136.7 (s), 138.3 (s), 141.4 (s), 148.9 (s), 149.1 (s), 168.6; HRMS (ESI): Calcd for C₃₆H₃₈N₃O₂ [M + H]⁺ 544.2959, found 544.2963.

3-Ethyl-4,4-bis[4-(N-morpholino)phenyl]-3-phenyl-3,4-dihydroquinoline 13j. Yield = 0.17 g, 60%; mp 184–185 °C (orange prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1713 and 1608; $\delta_{\rm H}$ (400 MHz; CDCl₃) 0.76 (3 H, t, *J* 6.8), 1.94–2.04 (1 H, m), 2.45–2.52 (1 H, m), 3.07–3.16 (8 H, m), 3.81–3.86 (8 H, m), 6.54 (2 H, d, *J* 8.8), 6.67–6.72 (4 H, m), 6.80 (2 H, d, *J* 7.6), 7.00–7.11 (4 H, m), 7.15–7.21 (2 H, m), 7.29 (2 H, d, *J* 8.8), 7.41 (1 H, d, *J* 7.6), 8.60 (1 H, s); $\delta_{\rm C}$ (100 MHz; CDCl₃) 8.4, 22.0, 48.3, 48.5, 51.4 (s), 60.6 (s), 66.4, 66.5, 112.7, 113.1, 126.1, 126.5, 126.8, 127.2, 127.6, 128.4, 130.1, 131.2, 131.8, 132.7 (s), 134.1 (s), 135.4 (s), 135.7 (s), 141.8 (s), 148.9 (s), 149.0 (s), 167.7; HRMS (ESI): Calcd for C₃₇H₄₀N₃O₂ [M + H]⁺ 558.3115, found 558.3122.

4,4-Bis(4-methoxyphenyl)-3-methyl-3-phenyl-3,4-dihydroquinoline 13k. Yield = 0.12 g, 57%; mp 168–170 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1606; δ_{H} (400 MHz; CDCl₃) 1.78 (3 H, s), 3.72 (3 H, s), 3.75 (3 H, s), 6.54 (2 H, d, *J* 9.0), 6.62 (2 H, d, *J* 7.7), 6.67 (2 H, d, *J* 9.0), 6.78 (2 H, d, *J* 8.4), 6.93 (2 H, t, *J* 7.7), 7.01–7.06 (2 H, m), 7.15–7.19 (2 H, m), 7.43–7.46 (3 H, m), 8.21 (1 H, s); δ_{C} (100 MHz; CDCl₃) 20.5, 48.8 (s), 55.2, 55.3, 60.2 (s), 111.8, 112.9, 126.7, 127.1, 127.6, 128.0, 128.5, 128.8, 128.9, 130.0, 132.9, 133.1 (s), 136.0 (s), 138.3 (s), 138.6 (s), 141.8 (s), 158.0 (s), 158.1 (s), 169.0; HRMS (ESI): Calcd for C₃₀H₂₈NO₂ [M + H]⁺ 434.2115, found 434.2119.

3-Ethyl-4, 4-bis (4-methoxyphenyl)-3-phenyl-3, 4-dihydroquinoline 131. Yield = 0.12 g, 52%; mp 187–188 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1606; δ_{H} (400 MHz; CDCl₃) 0.77 (3 H, t, *J* 7.3), 1.96–2.05 (1 H, m), 2.45–2.55 (1 H, m), 3.73 (3 H, s), 3.78 (3 H, s), 6.55 (2 H, d, *J* 9.1), 6.67 (2 H, d, *J* 7.7), 6.71 (2 H, d, *J* 9.1), 6.82 (2 H, d, *J* 7.7), 7.01–7.05 (2 H, m), 7.07–7.10 (2 H, m), 7.15 (1 H, dd, *J* 7.8 and 1.3), 7.19 (1 H, td, *J* 7.4 and 1.4), 7.31 (2 H, d, *J* 8.5 Hz), 7.42 (1 H, dd, *J* 7.7 and 1.2), 8.63 (1 H, s); δ_{C} (100 MHz; CDCl₃) 8.9, 25.4, 51.8 (s), 55.2, 55.3, 61.2 (s), 111.8, 112.3, 126.7, 127.1, 127.3, 127.8, 128.2, 128.8, 130.4, 131.8, 132.6, 134.0, 135.7 (s), 135.8 (s), 135.9 (s), 142.2 (s), 158.1 (s), 158.3 (s), 168.1; HRMS (ESI): Calcd for C₃₁H₃₀NO₂ [M + H]⁺ 448.2271, found 448.2273.

4,4-Bis(4-methoxyphenyl)-3,8-dimethyl-3-phenyl-3,4-dihydroquinoline 13m. Yield = 0.19 g, 85%; mp 158–160 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1629; δ_{H} (300 MHz; CDCl₃) 1.76 (3 H, s), 2.55 (3 H, s), 3.72 (3 H, s), 3.74 (3 H, s), 6.53 (2 H, d, *J* 8.4), 6.59 (2 H, d, *J* 7.8), 6.67 (2 H, d, *J* 8.4), 6.76 (2 H, d, *J* 7.8), 6.92–7.02 (6 H, m), 7.44 (2 H, d, *J* 8.4), 8.26 (1 H, s); δ_{C} (75 MHz; CDCl₃) 18.5, 20.7, 48.1 (s), 55.1, 55.2, 60.5 (s), 111.6, 112.7, 126.6, 126.7, 127.3, 127.5, 128.8, 128.9, 130.1, 133.0, 133.5 (s), 135.7 (s), 135.8 (s), 138.2 (s), 138.7 (s), 140.0 (s), 157.9 (s), 158.0 (s), 167.6; HRMS (ESI): Calcd for C₃₁H₃₀NO₂ [M + H]⁺ 448.2271, found 448.2276.

3-Ethyl-4,4-bis(4-methoxyphenyl)-8-methyl-3-phenyl-3,4-dihydroquinoline 13n. Yield = 0.23 g, 99%; colourless oil; v_{max} (neat)/cm⁻¹ 1606; δ_{H} (400 MHz; CDCl₃) 0.75 (3 H, t, *J* 7.2), 1.99 (1 H, m), 2.49 (1 H, m), 2.53 (3 H, s), 3.71 (3 H, s), 3.76 (3 H, s), 6.54 (2 H, d, *J* 8.8), 6.64 (2 H, d, *J* 7.6), 6.70 (2 H, d, *J* 8.8), 6.81 (2 H, d, *J* 8.0), 6.95–7.09 (6 H, m), 7.31 (2 H, d, *J* 8.0), 8.67 (1 H, s); δ_{C} (100 MHz; CDCl₃) 8.9, 18.3, 25.5, 51.1 (s), 55.1, 55.2, 61.4 (s), 111.7, 112.2, 126.5, 126.6, 127.1, 127.3, 128.9, 130.4, 131.9, 132.7, 134.2 (s), 135.4 (s), 135.5 (s), 135.9 (s), 136.2 (s), 140.5 (s), 158.0 (s), 158.2 (s), 166.6; HRMS (ESI): Calcd for C₃₂H₃₂NO₂ [M + H]⁺ 462.2428, found 462.2436.

Preparation of acridines 18, 2-arylquinolines 20 and quinolines 22

To a solution of iminophosphorane **11** (1 mmol) in anhydrous dichloromethane (15 mL) a solution of diphenylketene (0.19 g, 1 mmol) in the same solvent (5 mL) was added. The reaction mixture was stirred at room temperature under nitrogen for 30 min. Next the solvent was removed under reduced pressure and the resulting material was purified by column chromatography on silica gel, using hexanes/diethyl ether (9:1; v/v) as eluent, to provide the expected ketenimine **15**.

A deoxygenated solution of the ketenimine **15** (0.5 mmol) in anhydrous *ortho*-xylene (20 mL) was heated in a sealed tube at 180 $^{\circ}$ C during 2–21 d. After cooling, the solvent was removed under reduced pressure. The resulting material was purified by column chromatography on silica gel.

10-(2,2-Diphenylethenyl)-9-phenyl-9,10-dihydroacridine Yield = 0.21 g, 95%; mp 191–192 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1589; δ_{H} (300 MHz; CDCl₃) 5.22 (1 H, s), 6.61 (1 H, s), 6.83–6.88 (4 H, m), 7.07–7.21 (14 H, m), 7.37–7.41 (3 H, m), 7.48–7.51 (2 H, m); δ_{C} (75 MHz; CDCl₃) 47.9, 114.0, 121.5, 124.0, 124.7 (s), 126.1, 127.2, 127.5, 127.8, 128.2, 128.3, 128.5, 128.6, 129.0, 129.6, 137.7 (s), 140.1 (s), 145.5 (s), 147.3 (s); HRMS (ESI): Calcd for $C_{33}H_{26}N$ [M + H]⁺ 436.2060, found 436.2064.

10-(2,2-Diphenylethenyl)-3-methoxy-9-(4-methoxyphenyl)-9,10dihydroacridine 18b. Yield = 0.15 g, 61%; mp 150–151 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1595 and 1508; $\delta_{\rm H}$ (400 MHz; CDCl₃) 3.72 (3 H, s), 3.73 (3 H, s), 5.12 (1 H, s), 6.42 (1 H, dd, *J* 8.4 and 2.8), 6.59 (1 H, s), 6.62–6.65 (2 H, m), 6.74 (1 H, d, *J* 2.4), 6.76–6.78 (2 H, m), 6.85 (1 H, m), 6.98 (1 H, d, *J* 8.0), 7.08 (2 H, d, *J* 7.2), 7.10–7.21 (6 H, m), 7.38–7.41 (3 H, m), 7.46–7.49 (2 H, m); $\delta_{\rm C}$ (100 MHz; CDCl₃) 46.3, 55.3, 55.4, 100.5, 106.4, 113.9, 114.0, 117.8, 121.5, 123.9, 125.4 (s), 127.0, 127.8, 128.1, 128.2, 128.4, 128.5, 128.6, 129.0, 129.5, 130.2, 137.8 (s), 139.9 (s), 140.0 (s), 140.3 (s), 141.0 (s), 145.7 (s), 157.8 (s), 158.9 (s); HRMS (ESI): Calcd for C₃₅H₃₀NO₂ [M + H]⁺ 496.2271, found 496.2275.

2,4-Bis[4-(*N*,*N*-dimethylamino)phenyl]-3,3-diphenyl-3,4-dihydroquinoline **20a.** Yield = 0.13 g, 50%; colourless oil; $v_{max}(neat)/cm^{-1}$ 1608; $\delta_{H}(400 \text{ MHz}; \text{CDCl}_{3})$ 2.82 (6 H, s), 2.89 (6 H, s), 4.56 (1 H, s), 6.31 (2 H, d, *J* 7.6), 6.40–6.43 (4 H, m), 6.57 (2 H, d, *J* 8.8), 6.79 (2 H, t, *J* 8.4), 6.93–6.98 (3 H, m), 7.08–7.18 (2 H, m), 7.21–7.23 (2 H, m), 7.31 (2 H, d, *J* 9.2), 7.40 (1 H, d, *J* 8.0), 7.63 (2 H, d, *J* 8.0); $\delta_{C}(100 \text{ MHz}; \text{CDCl}_{3})$ 40.1, 40.6, 54.6, 59.0 (s), 110.5, 112.2, 125.6, 125.9, 126.5, 126.7, 126.9, 127.2, 127.3 (s), 127.6, 127.9, 128.8 (s), 129.7, 131.1, 131.2, 131.4, 131.8, 142.0 (s), 142.1 (s), 144.1 (s), 149.6 (s), 150.7 (s), 171.7 (s); HRMS (ESI): Calcd for C₃₇H₃₆N₃ [M + H]⁺ 522.2904, found 522.2908.

2,4-Bis[4-(*N*,*N*-dimethylamino)phenyl]-8-methyl-3,3-diphenyl-3,4-dihydroquinoline 20b. Yield = 0.25 g, 93%; mp 234–235 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1610 and 1519; $\delta_{\rm H}$ (300 MHz; CDCl₃) 2.46 (3 H, s), 2.80 (6 H, s), 2.89 (6 H, s), 4.51 (1 H, s), 6.32 (2 H, d, *J* 7.5), 6.41 (4 H, t, *J* 8.7), 6.54 (2 H, d, *J* 8.7), 6.76–6.84 (4 H, m), 6.93–6.98 (2 H, m), 7.11–7.23 (3 H, m), 7.40 (2 H, d, *J* 9.0), 7.61 (2 H, d, *J* 7.5); $\delta_{\rm C}$ (75 MHz; CDCl₃) 17.9, 40.2, 40.7, 55.0, 58.4 (s), 110.3, 112.3, 125.1, 125.6, 125.9, 126.2, 126.8, 127.7 (s), 127.8, 128.5, 129.2 (s), 129.7, 131.1, 131.3, 131.4, 134.8 (s), 142.3 (s), 142.4 (s), 142.5 (s), 149.5 (s), 150.5 (s), 168.6 (s); HRMS (ESI): Calcd for $C_{38}H_{38}N_3$ [M + H]⁺ 536.3060, found 536.3062.

3,3-Diphenyl-8-diphenylmethyl-3,4-dihydroquinoline22a.Yield = 0.11 g, 51%; mp 159–161 °C (colourless prisms
from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1599, 1581 and 1493;
 $\delta_{\rm H}(400 \text{ MHz; CDCl}_3)$ 3.47 (2 H, s), 6.55 (1 H, s), 6.81 (1 H, d, J
7.6), 6.95 (1 H, d, J 7.6), 7.00–7.07 (9 H, m), 7.17–7.28 (12 H,
m), 8.22 (1 H, s); $\delta_{\rm C}(100 \text{ MHz; CDCl}_3)$ 38.1, 48.7 (s), 50.4, 125.9,
126.1, 126.5 (s), 127.4, 127.7, 127.8, 128.1, 128.7, 129.8, 140.0
(s), 140.8 (s), 143.8 (s), 144.3 (s), 166.7; HRMS (ESI): Calcd for
C34H28N [M + H]⁺ 450.2216, found 450.2222.

8-Bis(4-methoxyphenyl)methyl-3,3-diphenyl-3,4-dihydroquinoline 22b. Yield = 0.13 g, 50%; mp 147–149 °C (colourless prisms from diethyl ether–hexane); v_{max} (Nujol)/cm⁻¹ 1610; δ_{H} (400 MHz; CDCl₃) 3.43 (2 H, s), 3.78 (6 H, s), 6.39 (1 H, s), 6.75–6.78 (4 H, m), 6.90–6.95 (5 H, m), 6.99 (1 H, d, *J* 7.6), 7.01–7.04 (5 H, m), 7.16–7.24 (6 H, m), 8.21 (1 H, s); δ_{C} (100 MHz; CDCl₃) 38.1, 48.6 (s), 48.7, 55.3, 113.5, 125.9, 126.4 (s), 126.7, 127.3, 127.8, 128.4, 128.5, 130.6, 136.9 (s), 140.6 (s), 140.7 (s), 143.8 (s), 157.7 (s), 166.6; HRMS (ESI): Calcd for C₃₆H₃₂NO₂ [M + H]⁺ 510.2428, found 510.2433.

Preparation of quinazolines 25

To a solution of iminophosphorane **11** (1 mmol) in anhydrous dichloromethane (15 mL) a solution of the aryl isocyanate (1 mmol) in the same solvent (5 mL) was added. The reaction mixture was stirred at room temperature for 30 min. The solvent was removed under reduced pressure, and the oily residue was chromatographed on a silica gel column using hexanes/diethyl ether (7:3, v/v) as eluent to give the corresponding carbodiimide **23**.

A solution of the carbodiimide **23** (0.5 mmol) in anhydrous *o*-xylene (20 mL) was heated a 180 °C, in a sealed tube, for 2–5 d. After cooling at room temperature, the solvent was removed under reduced pressure and the resulting material was purified by silica gel column chromatography.

3-(4-Bromophenyl)-4,4-bis[4-(*N*,*N***-dimethylamino)phenyl]-3,4-dihydroquinazoline 25a.** Yield = 0.22 g, 82%; mp 193–194 °C (colourless prisms from diethyl ether–hexane); v_{max} (Nujol)/cm⁻¹ 1608; δ_{H} (400 MHz; CDCl₃) 2.89 (12 H, s), 6.54–6.56 (4 H, m), 6.76–6.79 (3 H, m), 6.98–7.01 (1 H, m), 7.13–7.24 (8 H, m), 7.64 (1 H, s); δ_{C} (100 MHz; CDCl₃) 40.3, 71.9 (s), 111.3, 118.5 (s), 124.1, 125.3, 127.1, 127.2, 127.6, 129.0, 130.4, 131.2, 133.0 (s), 140.7 (s), 142.0 (s), 147.4, 149.3 (s); HRMS (ESI): Calcd for C₃₀H₃₀BrN₄ [M + H]⁺ 525.1648, found 525.1655.

3-(4-Chlorophenyl)-4,4-bis[4-(*N*,*N*-dimethylamino)phenyl]-8methyl-3,4-dihydroquinazoline 25b. Yield = 0.21 g, 84%; mp 230– 231 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1614 and 1574; $\delta_{\rm H}$ (300 MHz; CDCl₃) 2.41 (3 H, s), 2.90 (12 H, s), 6.52–6.57 (4 H, m), 6.61 (1 H, d, *J* 7.8), 6.83–6.87 (2 H, m), 6.91 (1 H, d, *J* 7.8), 7.00–7.04 (3 H, m), 7.11–7.16 (4 H, m), 7.70 (1 H, s); $\delta_{\rm C}$ (75 MHz; CDCl₃) 17.8, 40.4, 71.9 (s), 111.3, 124.7, 125.1, 126.6, 128.3, 129.0, 129.2 (s), 130.3 (s), 130.4, 131.9 (s), 133.0 (s), 139.1 (s), 141.7 (s), 146.5, 149.3 (s); HRMS (ESI): Calcd for C₃₁H₃₂ClN₄ [M + H]⁺ 495.2310, found 495.2317. **4,4-Bis[4-**(*N*,*N*-dimethylamino)phenyl]-8-methyl-3-(4-methylphenyl)-3,4-dihydroquinazoline 25c. Yield = 0.18 g, 76%; mp 255–257 °C (yellow prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1606; δ_{H} (400 MHz; CDCl₃) 2.19 (3 H, s), 2.42 (3 H, s), 2.88 (12 H, s), 6.53–6.56 (4 H, m), 6.59 (1 H, d, *J* 7.6), 6.79–6.81 (2 H, m), 6.85–6.89 (3 H, m), 7.01 (1 H, d, *J* 7.6), 7.14–7.18 (4 H, m), 7.73 (1 H, s); δ_{C} (75 MHz; CDCl₃) 17.9, 20.8, 40.5, 71.8 (s), 111.3, 124.3, 125.1, 125.5, 128.8, 128.9, 129.9 (s), 130.6, 131.7 (s), 133.0 (s), 134.4 (s), 139.5 (s), 140.6 (s), 147.3, 149.2 (s); HRMS (ESI): Calcd for $C_{32}H_{35}N_{4}$ [M + H]⁺ 475.2856, found 475.2860.

3-(4-Chlorophenyl)-4,4-bis(4-methoxyphenyl)-3,4-dihydroquinazoline 25d. Yield = 0.18 g, 78%; mp 96–97 °C (colourless prisms from diethyl ether–hexane); v_{max} (Nujol)/cm⁻¹ 1608; δ_{H} (300 MHz; CDCl₃) 3.71 (6 H, s), 6.72–6.82 (7 H, m), 7.01–7.06 (3 H, m), 7.19–7.26 (6 H, m), 7.66 (1 H, s); δ_{C} (75 MHz; CDCl₃) 55.3, 71.9 (s), 113.2, 124.4, 125.7, 126.9, 127.0, 128.1, 128.5, 130.8, 131.1 (s), 132.3 (s), 133.4 (s), 140.6 (s), 141.1 (s), 147.4, 158.8 (s); HRMS (ESI): Calcd for C₂₈H₂₄ClN₂O₂ [M + H]⁺ 455.1521, found 455.1527.

3-(4-Bromophenyl)-4,4-bis(4-methoxyphenyl)-8-methyl-3,4-di-hydroquinazoline 25e. Yield = 0.16 g, 61%; mp 83–85 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1608; $\delta_{\rm H}$ (400 MHz; CDCl₃) 2.42 (3 H, s), 3.76 (6 H, s), 6.55 (1 H, d, J 7.6), 6.73–6.77 (6 H, m), 6.93 (1 H, t, J 7.6), 7.06 (1 H, d, J 7.6), 7.17–7.22 (6 H, m), 7.70 (1 H, s); $\delta_{\rm C}$ (100 MHz; CDCl₃) 15.3, 55.3, 71.9 (s), 113.2, 118.6 (s), 124.9, 125.1, 126.8, 129.5, 130.8, 131.4, 132.3 (s), 132.4 (s), 133.4 (s), 139.1 (s), 141.8 (s), 146.3, 158.8 (s); HRMS (ESI): Calcd for C₂₉H₂₆BrN₂O₂ [M + H]⁺ 513.1172, found 513.1174.

Preparation of 4-chloro-2-[1-(4-methoxyphenyl)vinyl]-*N*-(2,2-diphenylvinyl)aniline 30

To a solution of iminophosphorane **27** (0.8 g, 1.5 mmol) in anhydrous dichloromethane (15 mL) a solution of diphenylketene (0.24 g, 1.5 mmol) in the same solvent (5 mL) was added. The reaction mixture was stirred at room temperature under nitrogen for 30 min. Next the solvent was removed under reduced pressure and the resulting material was purified by column chromatography on silica gel, using hexanes/diethyl ether (9.5:0.5; v/v) as eluent, to provide the ketenimine **28** (81% yield).

A solution of the ketenimine **28** (0.33 g, 0.75 mmol) in anhydrous *ortho*-xylene (20 mL) was heated at reflux temperature under nitrogen for 3 h. After cooling, the solvent was removed under reduced pressure and the resulting material was purified by column chromatography on silica gel, using hexanes/diethyl ether (9:1, v/v) as eluent.

4-Chloro-2-[1-(4-methoxyphenyl)vinyl]-*N*-(**2**,**2**-diphenylvinyl)aniline **30.** Yield = 0.30 g, 91%; orange oil; v_{max} (neat)/cm⁻¹ 1659; $\delta_{\rm H}$ (300 MHz; CDCl₃) 3.79 (3 H, s), 5.07 (1 H, d, *J* 0.9), 5.53 (1 H, d, *J* 0.9), 6.10 (1 H, d, *J* 12.0), 6.71–6.75 (2 H, m), 6.88–6.96 (3 H, m), 6.99–7.04 (2 H, m), 7.08–7.13 (3 H, m), 7.18–7.25 (7 H, m), 7.29–7.38 (1 H, m); $\delta_{\rm C}$ (75 MHz; CDCl₃) 55.3, 112.7, 114.1, 115.4, 118.4 (s), 123.9, 124.0 (s), 125.6, 126.2, 126.8, 127.6, 128.3, 128.8, 129.0, 129.8, 130.5, 130.7 (s), 137.9 (s), 138.7 (s), 141.6 (s), 144.5 (s), 159.8 (s); HRMS (ESI): Calcd for C₂₉H₂₅CINO [M + H]⁺ 438.1619, found 438.1619.

Preparation of quinolines 34

To a solution of 1-phenylmethyl-2-triphenylphosphoranylideneaminobenzene **32** (0.5 g, 1.1 mmol) in anhydrous dichloromethane (15 mL) a solution of diphenylketene (1.1 mmol) or methylphenylketene (1.1 mmol) in the same solvent (5 mL) was added. The reaction mixture was stirred at room temperature under nitrogen for 30 min. Next the solvent was removed under reduced pressure and the resulting material was purified by column chromatography on silica gel, using hexanes/diethyl ether (9.5:0.5; v/v) as eluent, to provide ketenimine **33a** (91% yield) or ketenimine **33b** (70% yield), respectively.

A deoxygenated solution of ketenimine **33** (0.5 mmol) in anhydrous toluene (20 mL) was heated in a sealed tube at 160 °C during 36 h. After cooling, the solvent was removed under reduced pressure. The resulting material was purified by column chromatography on silica gel, using hexanes/diethyl ether (4:1) as eluent.

3,3,4-Triphenyl-3,4-dihydroquinoline 34a. Yield = 0.17 g, 97%; mp 192–194 °C (colourless prisms from diethyl ether); v_{max} (Nujol)/cm⁻¹ 1614; δ_{H} (400 MHz; CDCl₃) 4.75 (1 H, s), 6.56 (2 H, d, J 7.2), 6.76–6.78 (2 H, m), 6.92–7.00 (3 H, m), 7.03–7.06 (3 H, m), 7.21–7.23 (4 H, m), 7.25–7.29 (2 H, m), 7.30–7.34 (2 H, m), 7.45 (1 H, d, J 7.2), 8.54 (1 H, s); δ_{C} (100 MHz; CDCl₃) 50.8, 54.2 (s), 126.3, 126.4, 127.0, 127.6, 127.7, 127.8, 127.9, 128.2, 128.4, 128.5, 128.9, 129.1, 130.1 (s), 139.6 (s), 141.8 (s), 143.1 (s), 144.3 (s), 165.5; HRMS (ESI): Calcd for C₂₇H₂₂N [M + H]⁺ 360.1747, found 360.1751.

3-Methyl-3,4-diphenyl-3,4-dihydroquinoline 34b. Yield = 0.10 g, 70%; colourless oil; v_{max} (neat)/cm⁻¹ 1622; δ_{H} (300 MHz; CDCl₃) 1.30 (3 H, s, minor isomer), 1.62 (3 H, s, major isomer), 4.02 (1 H, s, major isomer), 4.28 (1 H, s, minor isomer), 6.47–6.49 (3 H, m), 6.89–7.36 (23 H, m), 7.49 (1 H, d, *J* 7.5, minor isomer), 7.56 (1 H, d, *J* 7.8, major isomer), 7.90 (1 H, s, minor isomer), 8.05 (1 H, s, major isomer), 45.4 (s, minor isomer), 46.3 (s, major isomer), 54.0 (minor isomer), 54.7 (major isomer), 126.6, 126.9, 127.0, 127.1, 127.3, 127.4, 127.6, 127.7, 127.89, 127.94, 128.0, 128.1, 128.4, 128.5, 128.8, 129.6, 129.8, 138.5 (s), 139.1 (s), 140.7 (s), 142.1 (s), 142.2 (s), 144.1 (s), 168.8; HRMS (ESI): Calcd for $C_{22}H_{20}N$ [M + H]⁺ 298.1590, found 298.1594.

Acknowledgements

This work was supported by the Ministerio de Ciencia e Innovacion of Spain (Project CTQ2008-05827/BQU) and Fundacion Seneca-CARM (Project 08661/PI/08). B. B. and M.-M. O. thank Fundacion Seneca-CARM and Fundacion CajaMurcia for their respective fellowships.

Notes and references

- 1 (a) K. Godula and D. Sames, *Science*, 2006, **312**, 67–72; (b) For a series of recent monographic reviews, see: *Chem. Rev.*, 2010, Issue 2.
- 2 (a) I. D. Junberg, Y. Odabachian and F. Gagosz, J. Am. Chem. Soc., 2010, **132**, 3543–3552; (b) P. A. Vadola and D. Sames, J. Am. Chem. Soc., 2009, **131**, 16525–16528; (c) K. M. McQuaid, J. Z. Long and D. Sames, Org. Lett., 2009, **11**, 2972–2975; (d) K. M. McQuaid and D. Sames, J. Am. Chem. Soc., 2009, **131**, 402–403; (e) S. J. Pastine and D. Sames, Org. Lett., 2005, **7**, 5429–5431; (f) S. J. Pastine, K. M.

McQuaid and D. Sames, *J. Am. Chem. Soc.*, 2005, **127**, 12180–12181. No substrate possessing a heterocumulenic functionality as hydride acceptor was involved in these reports.

- 3 (a) S. Murarka, I. Deb, C. Zhang and D. Seidel, J. Am. Chem. Soc., 2009, 131, 13226–13227; (b) C. Zhang, S. Murarka, S. and D. Seidel, J. Org. Chem., 2009, 74, 419–422; (c) S. Murarka, C. Zhang, M. D. Konieczynska and D. Seidel, Org. Lett., 2009, 11, 129–132, and references cited in these reports.
- 4 M. Alajarin, B. Bonillo, M.-M. Ortin, P. Sanchez-Andrada and A. Vidal, *Org. Lett.*, 2006, **8**, 5645–5648.
- 5 (a) M. Alajarin, B. Bonillo, P. Sanchez-Andrada, A. Vidal and D. Bautista, Org. Lett., 2009, **11**, 1365–1368; (b) M. Alajarin, B. Bonillo, P. Sanchez-Andrada and A. Vidal, J. Org. Chem., 2010, **75**, 3737–3750.
- 6 (a) C. Creutz and M. H. Chou, J. Am. Chem. Soc., 2009, 131, 2794–2795; (b) M. R. Nimlos, C. H. Chang, C. J. Curtis, A. Miedaner, H. M. Pilath and D. L. DuBois, Organometallics, 2008, 27, 2715–2722; (c) X.-J. Qi, Y. Fu, L. Liu and Q.-X. Guo, Organometallics, 2007, 26, 4197–4203; (d) W. W. Ellis, A. Miedaner, C. J. Curtis, D. H. Gibson and D. L. DuBois, J. Am. Chem. Soc., 2002, 124, 1926–1932; (e) T.-Y. Cheng, B. S. Brunschwig and R. M. Bullock, J. Am. Chem. Soc., 1998, 120, 13121–13137; (f) T.-Y. Cheng and R. M. Bullock, Organometallics, 1995, 14, 4031–4033.
- 7 (a) D. Richter and H. Mayr, Angew. Chem., Int. Ed., 2009, 48, 1958–1961; (b) R. Lavilla, Curr. Org. Chem., 2004, 8, 715–737; (c) X.-Q. Zhu, H.-R. Li, Q. Li, T. Ai, J.-Y. Lu, Y. Yang and J.-P. Cheng, Chem.–Eur. J., 2003, 9, 871–880; (d) S. Fukuzumi, Q. Inada and T. Suenobu, J. Am. Chem. Soc., 2003, 125, 4808–4816; (e) X.-Q. Zhu, Y.-C. Liu and J.-P. Cheng, J. Org. Chem., 1999, 64, 8980–8981.
- 8 (a) W. W. Ellis, J. W. Raebiger, C. J. Curtis, J. Bruno and D. L. DuBois, J. Am. Chem. Soc., 2004, 126, 2738–2743; (b) W. W. Ellis, R. Ciancanelli, S. M. Miller, J. W. Raebiger, M. R. DuBois and D. L. DuBois, J. Am. Chem. Soc., 2003, 125, 12230–12236; (c) E.-U. Würthwein, G. Lang, L. H. Schappele and H. Mayr, J. Am. Chem. Soc., 2002, 124, 4084– 4092; (d) X.-M. Zhang, J. W. Bruno and E. Enyinnaya, J. Org. Chem., 1998, 63, 4671–4678; (e) K. L. Handoo, J.-P. Cheng and V. D. Parker, J. Am. Chem. Soc., 1993, 115, 5067–5072; (f) J.-P. Cheng, K. L. Handoo and V. D. Parker, J. Am. Chem. Soc., 1993, 115, 2655–2660.
- 9 (a) D. E. Schwarz, T. M. Cameron, P. J. Hay, B. L. Scott, W. Tumas and D. L. Thorn, *Chem. Commun.*, 2005, 5919–5921; (b) I.-S. Lee and E. H. Jeoung, *J. Org. Chem.*, 1998, **63**, 7275–7279; (c) F. Montgrain, S. M. Ramos and J. D. Wuest, *J. Org. Chem.*, 1988, **53**, 1489–1492.
- 10 X.-M. Zhang, J. W. Bruno and E. Enyinnaya, J. Org. Chem., 1998, 63, 4671–4678.

- 11 P. Brunet and J. D. Wuest, J. Org. Chem., 1996, 61, 2020–2026 and references cited therein.
- 12 H. E. Ungnade and E. W. Crandall, J. Am. Chem. Soc., 1949, 71, 2209– 2210.
- 13 K. Fukui, Y. Inamoto, H. Gitano and C. Nagata, J. Am. Chem. Soc., 1959, 81, 5954–5957.
- 14 X. Wang, Y. Wang, D.-M. Du and J. Xu, J. Mol. Catal. A: Chem., 2006, 255, 31–35.
- 15 R. N. Carde, G. Jones, W. H. McKinley and C. Price, J. Chem. Soc., Perkin Trans. 1, 1978, 1211–1218.
- 16 All the heterocumulenes, ketenimines and carbodiimides synthesized in this work were similarly treated due to their hydrolytic sensitivity.
- 17 The standard Csp³–Csp³ bond distance is 1.54 Å, see: F. H. Allen, D. G. Kennard, L. Watson, A. G. Barmmer, A. G. Orpen and R. Taylor, *J. Chem. Soc., Perkin Trans.* 2, 1987, S1–S19.
- 18 For a related example of enhancement of reactivity by an *ortho* substituent, see: K. Mori, T. Kawasaki, S. Sueoka and T. Akiyama, *Org. Lett.*, 2010, **12**, 1732–1735.
- 19 The results of a preliminary DFT computation study of the transformation 15a → 18a are fully consistent with the reaction path outlined in Scheme 6. The alternative C1-C6 electrocyclization of azaxylylene 16 into the corresponding 3,3,4,4-tetraphenyl-3,4-dihydroquinoline has been also calculated and shown to be non competitive, on kinetic and thermodynamic grounds, with the experimentally-observed channel leading to 18.
- 20 For an example of a 1,3 migration of a phenyl group in ketenimines, see: (a) C. O. Kappe, G. Kollenz, R. Leung-Toung and C. Wentrup, J. Chem. Soc., Chem. Commun., 1992, 487–488; (b) For some examples of reactions involving 1,5-aryl shifts, see: Z.-Y. Chen and M.-J. Wu, Org. Lett., 2005, 7, 475–477; (c) B. Schulte, R. Froehlich and A. Studer, Tetrahedron, 2008, 64, 11852–11859; (d) A. Studer, M. Bossart and T. Vasella, Org. Lett., 2000, 2, 985–988; (e) P. J. Battye and D. W. Jones, J. Chem. Soc., Chem. Commun., 1984, 990–992; (f) P. T. Lansbury and R. L. Letsinger, J. Am. Chem. Soc., 1959, 81, 940–943.
- 21 (a) A. R. De Lera and F. P. Cossio, Angew. Chem., Int. Ed., 2002, 41, 1150–1152; (b) J. Rodriguez-Otero and E. M. Cabaleiro-Largo, Chem.– Eur. J., 2003, 9, 1837–1843; (c) M. Alajarin, P. Sanchez-Andrada, F. P. Cossio, A. Arrieta and B. Lecea, J. Org. Chem., 2001, 66, 8470–8377; (d) M. Alajarin, P. Sanchez-Andrada, A. Vidal and F. Tovar, J. Org. Chem., 2005, 70, 1340–1349; (e) M. Alajarin, A. Vidal, P. Sanchez-Andrada, F. Tovar and G. Ochoa, Org. Lett., 2000, 2, 965–968; (f) S. Sakai, Theor. Chem. Acc., 2008, 120, 177–183; (g) J. A. Duncan, D. E. G. Calkins and M. Chavartha, J. Am. Chem. Soc., 2008, 130, 6740–6748.