



Contents lists available at ScienceDirect

## Bioorganic &amp; Medicinal Chemistry

journal homepage: [www.elsevier.com/locate/bmc](http://www.elsevier.com/locate/bmc)

# Facile access to modified and functionalized PNAs through Ugi-based solid phase oligomerization

Jacques Saarbach, Daniela Masi, Claudio Zambaldo, Nicolas Winssinger \*

Faculty of Science, Department of Organic Chemistry, NCCR Chemical Biology, University of Geneva, 30 quai Ernest Ansermet, Geneva, Switzerland

## ARTICLE INFO

## Article history:

Received 29 March 2017

Revised 30 May 2017

Accepted 31 May 2017

Available online xxxxx

## Keywords:

Peptide nucleic acid

Ugi reaction

Solid phase synthesis

Biosupramolecular chemistry

## ABSTRACT

Peptide nucleic acids (PNAs) derivatized with functional molecules are increasingly used in diverse biosupramolecular applications. PNAs have proven to be highly tolerant to modifications and different applications benefit from the use of modified PNAs, in particular modifications at the  $\gamma$  position. Herein we report simple protocols to access modified PNAs from iterative Ugi couplings which allow modular modifications at the  $\alpha$ ,  $\beta$  or  $\gamma$  position of the PNA backbone from simple starting materials. We demonstrate the utility of the method with the synthesis of several bioactive small molecules (a peptide ligand, a kinase inhibitor and a glycan)-PNA conjugates.

© 2017 Elsevier Ltd. All rights reserved.

## 1. Introduction

Peptide nucleic acids (PNAs) are artificial oligomers in which the phosphoribose of DNA or RNA has been replaced by a peptidic backbone (Fig. 1).<sup>1–3</sup> PNAs stand out for their ability to form more stable duplex with DNA or RNA than the natural homoduplex, for their high sequence discrimination, as well as for their biological and chemical robustness. These remarkable properties of PNA have attracted tremendous attention for diagnostic and therapeutic applications.<sup>4</sup> Over the past decade the field of applications of PNA has grown significantly with applications of this oligomer as a biosupramolecular tag for programmable assemblies and reactions,<sup>5–8</sup> in DNA-based nanotechnologies,<sup>9,10</sup> to direct folding of protein-like structures,<sup>11–13</sup> and as chloride ion sensor.<sup>14</sup> For these purposes, PNAs bearing modification in the backbone to introduce various functionalities or ligands are frequently used. Several modifications to the N-(2-aminoethyl)glycine backbone have been reported as beneficial or well tolerated in the hybridization.<sup>15–18</sup> Substitution at the  $\alpha$  position (replacement of glycine with other amino acids)<sup>19</sup> are tolerated while a *gem*-dimethyl substitution at the  $\alpha$  position provides enhanced duplex stability with DNA.<sup>20</sup> An important modification is the introduction of a chiral  $\gamma$ -substituent ((*S*) stereochemistry defined based on a methyl substituent) which generates a steric bias that favors right-handed helicity with base stacking, resulting in enhanced duplex stability with DNA.<sup>21–25</sup> The enhanced properties of this backbone have




recently been used to target genomic DNA.<sup>26</sup> A lysine<sup>22,27,28</sup> or cysteine<sup>29,30</sup> side chain at the  $\gamma$ -position has also proven useful for the conjugation of small molecules. The combination of chirality at the  $\alpha$  and  $\gamma$  positions within a single PNA unit provides additive duplex stability with an overriding influence of the  $\gamma$  stereochemistry.<sup>31</sup> More recently, modification at the  $\beta$  position has also been reported to be tolerated (*S*-methyl).<sup>32</sup> Furthermore, using the opposite stereochemistry at the  $\gamma$  positions has been used to create PNA with orthogonal hybridization to DNA or RNA.<sup>33,34</sup> In addition to modifications of the PNA backbone, many alternative nucleobases beyond the canonical A, C, G, and T have been designed and can provide interesting properties: enhanced hybridization (2,6-diaminopurine,<sup>35</sup> analogs of thymine (7-Cl-bT)<sup>36</sup> and cytosine (G-clamp)<sup>37</sup>, promiscuous hybridization (hypoxanthine)<sup>38</sup> or fluorescent probes (thiazole orange).<sup>39</sup>

## 2. Results and discussion

PNAs are typically synthesized in the C to N direction by iterative cycles of coupling/ deprotection with monomers protected orthogonally on the nucleobases and N-terminus (Fig. 1). However, the exponential synthetic demand to access modified PNA monomers in the various combinations of backbones and nucleobases is limiting their implementation. Herein, we report a straightforward method based on reiterative Ugi reactions<sup>40</sup> to access PNA oligomers with modifications at the  $\alpha$ ,  $\beta$ ,  $\gamma$  positions in combination with any nucleobase. The methodology also lends itself to the synthesis of PNA bearing ligand at defined positions within the backbone. While the Ugi 4-CR reaction has been previously

\* Corresponding author.

E-mail address: [nicolas.winssinger@unige.ch](mailto:nicolas.winssinger@unige.ch) (N. Winssinger).

**Standard PNA synthesis: C to N synthesis****Ugi-based PNA synthesis****Method 1: C to N synthesis****Method 2: N to C synthesis**

**Fig. 1.** General strategy for PNA synthesis by standard reiterative coupling (top); C to N Ugi couplings (middle) and N to C Ugi couplings (bottom). NB: nucleobase; Pg: protecting group.

used to synthesize PNA monomers or dimers,<sup>41–45</sup> these previous methods had not been implemented for oligomer synthesis with functionalization of the PNAs.

We began our investigation with the evaluation of the Ugi reaction to prepare PNA in the C to N direction (Method 1, Fig. 1). To facilitate the LC-MS analysis of the reactions, the synthesis was initiated with a resin loaded with a lysine naphthoate prior to the introduction of the first residue by standard Fmoc-based chemistry (Fig. 1). The naphthyl group has a UV absorption at 283 nm, which provides a distinct window to identify the product and any side product arising from subsequent reactions. This resin loaded with a first amino acid was treated with 4 equivalents of nucleobase (2, thymine acetic acid), aldehyde 3 and benzyl isocyanide in EtOH. The reaction proceeded to completion for all 19 natural primary amino acids after 1 h. While there are a few examples of chiral induction in Ugi-4CR,<sup>46,47</sup> they are performed at low temperature under anhydrous conditions and we did not expect notable diastereoselectivity. Performing the reaction with resin 1 loaded with 1-Aminocyclopropanecarboxylic acid afforded a 1:1 mixture of diastereoisomers resolvable by HPLC. To further ascertain the diastereoselectivity, the reaction was performed in solution with the methyl ester of valine instead of resin 1 followed by Fmoc deprotection with concomitant cyclization to the ketopiperazine clearly indicating a mixture of diastereoisomers as judged by LC-MS and NMR. Based on the efficient Ugi-4CR, we synthesized an 8-mer by reiteration of the cycle (amino acid coupling, Fmoc deprotection, Ugi reaction, Fmoc deprotection) using glycine and the four canonical nucleobases and benzyl isocyanide to obtain the PNA as a single product thus establishing the viability of this Ugi-based oligomerization albeit without control of the stereochemistry at the  $\beta$  position (Fig. 2).

We next evaluated the Ugi reaction to prepare PNAs in the N to C direction (Fig. 1). This second Ugi variation provides inherent control over the most important stereochemistry at  $\beta$  or  $\gamma$  positions (either stereochemistry at the  $\alpha$  position is tolerated in the

hybridization). To this end, three different isocyanides (9<sup>48</sup>, 12, 14) were prepared as shown in Fig. 3 to ultimately yield unmodified PNA,  $\gamma$  or  $\beta$ -modified PNAs respectively.<sup>49,50</sup> Chiral isocyanides 12 and 14 were obtained in three steps from intermediate 10.<sup>51</sup> For isocyanide 12, intermediate 10 was Fmoc protected and the Mtt was removed with formic acid and acetic anhydride to directly obtain 11. Dehydration of the formamide using Burgess's reagent or POCl<sub>3</sub> afforded 12 in excellent yield. A similar sequence starting with the formylation of 10 was used to obtain 14 in equally good yield. Investigation into the Ugi reaction starting from resin 5 with isocyanide 9, thymine acetic acid 2 and formaldehyde (Fig. 4) led to the formation of two peaks by LC-MS, the expected product 16 and a side product with a mass corresponding to 17. This side product was assumed to arise from reaction intermediate 15 that engaged in a second imine condensation (reaction path b, Fig. 4) rather than undergo the acyl transfer of the Ugi condensation (reaction path a) to afford 16. Attempts to optimize the reaction conditions (solvent, equivalents of formaldehyde, temperature) failed to completely shut down this side reaction. This side reaction precludes the use of formaldehyde for reiterative Ugi coupling towards PNA oligomers. However, this side reaction was not observed with other aldehydes and the use of diverse aldehydes afforded the desired product in >95% yield in <30 min reaction time at room temperature. The best conditions employed THF as a solvent with, 4 equivalents of isocyanide, 4 equivalents of a nucleobase (added as a NMP solution) and 50 equivalents of hexafluoroisopropanol (HFIP)<sup>52</sup> for 30 min. Notably, the reaction could be used to introduce diverse substituent at the  $\alpha$  position (PEG 18a for solubility<sup>53</sup>, 2-nitrobenzyl 18b for photocleavable PNA or azide functionalized side chain 18c for further conjugation) to obtain PNAs 19a–d. Performing the reaction with chiral isocyanides 12 or 14 proceeded equally well affording the products as a mixture of diastereoisomers at the  $\alpha$  position (ca. 1:1, based on LC-MS integration of compound 19d-2), however, the lack of stereochemical control at the  $\alpha$  position has minimal impact on PNA hybridization based on the




Fig. 2. PNA synthesis by C to N Ugi couplings, method 1.




Fig. 3. Synthesis of isocyanide 9–14.

overriding influence of the defined stereochemistry at the  $\gamma$  position, which is defined by the isocyanide.<sup>54</sup> Nonetheless, we sought an alternative to obtain PNAs without substitution at the  $\alpha$  position. As an alternative to the use of formaldehyde, we reasoned that performing the Ugi with aldehyde **18e** would afford a product that would readily undergo a decarboxylation, following cleavage and global deprotection. It was found that aldehyde **18e** could be used in the Ugi reaction affording product **20e** which indeed underwent a decarboxylation under classical conditions (10% aq. acetic acid, 90 °C). Thus, the use of aldehyde **18e** affords PNAs unmodified at the  $\alpha$  position. Reaction with cyclic ketone **21** afforded the desired product with a *gem*-disubstitution at the  $\alpha$  position. The latter two reactions with aldehyde **18e** or ketone **21** where slower and required 50 °C to reach completion within 1 h. To illustrate the utility of this Ugi-based PNA synthesis method, we prepared various examples combining different nucleobases and backbone modifications (four examples of 8mer PNA using isobutyraldehyde, isocyanide **9** or **12a** and five nucleobases; one example of 6mer PNA alternating PEG and isobutyraldehyde with isocyanide **9**). All PNAs were obtained as a single major product showing the expected mass by MALDI and LC-MS (one example shown in Fig. 4, see SI for additional examples).

We then turned our attention to the application of this methodology to prepare PNAs modified at defined position with ligands. We anticipated that peptides or other ligands could be derivatized with a Fmoc-*D*-Dap(Mtt)OH residue that would result in the desired stereochemistry at the  $\gamma$  position, while providing orthogonally protected amino terminus to synthesize the PNA via regular Fmoc chemistry then deprotect the Mtt<sup>51</sup> and continue with Ugi-based PNA synthesis. As shown in Fig. 5, this strategy was used to synthesize an Abl kinase ligand<sup>55</sup> (11 mer peptide, **26**) which was derivatized with a PEG followed Fmoc-*D*-Dap(Mtt)OH to obtain **27**. The Fmoc was removed and 8 cycles of regular Fmoc-based PNA synthesis were performed. The Mtt was then deprotected and 4 cycles of N to C Ugi oligomerization were performed. Cleavage of the product from the resin afford the desired compound **28** as the major product. Alternately, the trifunctional linker **29** can be used to achieve ligand coupling after completion of PNA synthesis with CuAAC conjugation of the ligand. To illustrate this strategy, resin **29** was Mtt-deprotected and engaged in 4 cycles of Ugi-based PNA synthesis. Then, the azide was reduced (TCEP)<sup>56</sup> and the 8 cycles of regular Fmoc-based PNA synthesis were performed. Ultimately, two different ligands were coupled, a kinase inhibitor (dasatinib)<sup>57</sup> or a glycan (cellobiose) to obtain **30** and





Fig.4. Ugi oligomerization in the N to C direction, method 2.



**Fig. 5.** Application of bi-directional synthesis for the preparation of PNA-ligand conjugates at defined position in the oligomer backbone Decarboxylation of a bi-directional PNA 8-mer.

**31.**<sup>58</sup> This strategy provided the desired conjugate with high purity. Finally, to check the efficiency of the decarboxylation of PNAs prepared with aldehyde **18e**, a bidirectional 8-mer PNA **32** (4 cycles of Ugi-based synthesis and 4 cycles of regular Fmoc-based PNA synthesis) was prepared and decarboxylated to yield **33** (10% acetic acid in water, 90 °C, 1 h). The decarboxylated PNA **33** was lyophilized and LC-MS analysis indicated a complete decarboxylation without any degradation, which establishes that Ugi-based strategy can be used in combination with classical PNA synthesis.

### 3. Experimental section

Experimental procedures for reagents and resins are detailed in the supplemental information

### 3.1. General procedure for C to N Ugi reaction

Representative example with aldehyde **3** and benzyl isocyanide (other reactions performed under the same conditions with

different aldehydes or isocyanides): To a solution of nucleobase acetic acid (8.8  $\mu\text{mol}$ , 4.0 equiv) in 5  $\mu\text{L}$  of NMP was added Fmoc-acetaldehyde **3** (2.4 mg, 8.8  $\mu\text{mol}$ , 4.0 equiv) and benzyl isocyanide (1.2  $\mu\text{L}$ , 8.8  $\mu\text{mol}$ , 4.0 equiv) followed by 25  $\mu\text{L}$  of MeOH. The solution was first sonicated in an ultra-sound bath then added to resin **1** (5 mg, 2.5  $\mu\text{mol}$ , 1.0 equiv) which was shaken for 1 h at room temperature. Subsequently the resin was first washed with DMF (6 $\times$ ) and  $\text{CH}_2\text{Cl}_2$  (6 $\times$ ).

### 3.2. General procedure for N to C Ugi reaction

Representative example with isocyanide **9** and thymine acetic acid **2** and aldehyde **18** (other reactions performed under the same conditions with different aldehydes or isocyanides): Isocyanide **9** (4.0 equiv, 10.0  $\mu\text{mol}$ , 2.8 mg) was dissolved in 10.0  $\mu\text{L}$  of a 1 M solution of thymine acetic acid **2** in NMP (4.0 equiv., 10.0  $\mu\text{mol}$ ), the suspension was sonicated until the solution became clear, then HFIP (50.0 equiv, 110.0  $\mu\text{mol}$ , 12.0  $\mu\text{L}$ ) and 80  $\mu\text{L}$  of THF were added. Aldehyde **18** (4.0 equiv, 10.0  $\mu\text{mol}$ , 1.2  $\mu\text{L}$ ) was added to the previous solution and the mixture was transferred to the resin. The reaction was shaken for 30 min at room temperature. Subsequently the resin was first washed with DMF (6 $\times$ ) and  $\text{CH}_2\text{Cl}_2$  (6 $\times$ ).

### 3.3. General decarboxylation procedure

The PNA was cleaved from resin using 100% TFA (1 h, room temperature) and precipitated in diethylether (10 $\times$  volume of TFA). The solution was centrifuged to obtain a pellet, which was decanted and taken up in 10% acetic acid in water. The solution was heated to 90  $^\circ\text{C}$  for 1 h. The decarboxylation can be observed by formation of bubbles inside the vial and monitored by LC-MS. The final PNA derivative was obtained by lyophilizing the solution.

## 4. Conclusion

The methodologies reported herein greatly facilitates the synthesis of functionalized PNAs. The N to C Ugi oligomerization enable bi-directional PNA synthesis with conjugation of ligands to the PNA at a given position in the oligomer. The current interest in the use of ODNs to display small molecules with defined topology or use of PNA as sensor and foldamers should greatly benefit from the synthetic methods reported herein. The fact that any combination of backbone and nucleobase modifications can be accessed without lengthy synthesis of monomers should open new horizons in PNA applications.

## Acknowledgments

The authors thank the Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung (grant: 200020\_157106; 200020\_169141) and the NCCR Chemical Biology for financial support.

## A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bmc.2017.05.064>.

## References

- Egholm M, Buchardt O, Christensen L, et al. PNA hybridizes to complementary oligonucleotides obeying the Watson-Crick hydrogen-bonding rules. *Nature*. 1993;365:566–568.
- Nielsen PE, Egholm M, Berg RH, Buchardt O. Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. *Science*. 1991;254:1497–1500.
- Nielsen PE. Peptide nucleic acid. A molecule with two identities. *Acc Chem Res*. 1999;32:624–630.
- Hyrup B, Nielsen PE. Peptide nucleic acids (PNA): synthesis, properties and potential applications. *Bioorg Med Chem*. 1996;4:5–23.
- Gorska K, Winssinger N. Reactions templated by nucleic acids: more ways to translate oligonucleotide-based instructions into emerging function. *Angew Chem Int Ed Engl*. 2013;52:6820–6843.
- Percivalle C, Bartolo JF, Ladame S. Oligonucleotide-templated chemical reactions: pushing the boundaries of a nature-inspired process. *Org Biomol Chem*. 2013;11:16–26.
- Shi B, Zhou Y, Huang Y, Zhang J, Li X. Recent advances on the encoding and selection methods of DNA-encoded chemical library. *Bioorg Med Chem Lett*. 2017;27:361–369.
- Zambaldo C, Barluenga S, Winssinger N. PNA-encoded chemical libraries. *Curr Opin Chem Biol*. 2015;26:8–15.
- Barluenga S, Winssinger N. PNA as a biosupramolecular tag for programmable assemblies and reactions. *Acc Chem Res*. 2015;48:1319–1331.
- Ranallo S, Rossetti M, Plaxco KW, Vallee-Belisle A, Ricci F. A modular, DNA-based beacon for single-step fluorescence detection of antibodies and other proteins. *Angew Chem Int Ed Engl*. 2015;54:13214–13218.
- Machida T, Dutt S, Winssinger N. Allosterically regulated phosphatase activity from peptide-PNA conjugates folded through hybridization. *Angew Chem Int Ed*. 2016;55:8595–8598.
- Brudno Y, Birnbaum ME, Kleiner RE, Liu DR. An in vitro translation, selection and amplification system for peptide nucleic acids. *Nat Chem Biol*. 2010;6:148–155.
- Rapireddy S, Nhon L, Meehan RE, et al. RID-1Mimic containing gamma PNA scaffold exhibits broad-spectrum antibacterial activities. *J Am Chem Soc*. 2012;134:4041–4044.
- Prakash V, Saha S, Chakraborty K, Krishnan Y. Rational design of a quantitative, pH-insensitive, nucleic acid based fluorescent chloride reporter. *Chem Sci*. 2016;7:1946–1953.
- Sugiyama T, Kittaka A. Chiral peptide nucleic acids with a substituent in the N-(2-aminoethyl)glycine backbone. *Molecules*. 2012;18:287–310.
- Dragulescu-Andrasi A, Zhou P, He G, Ly DH. Cell-permeable GPNA with appropriate backbone stereochemistry and spacing binds sequence-specifically to RNA. *Chem Commun (Camb)*. 2005;244–246.
- Zhou P, Wang M, Du L, Fisher GW, Waggoner A, Ly DH. Novel binding and efficient cellular uptake of guanidine-based peptide nucleic acids (GPNA). *J Am Chem Soc*. 2003;125:6878–6879.
- Puschl A, Sforza S, Haaime G, Dahl O, Nielsen PE. Peptide nucleic acids (PNAs) with a functional backbone. *Tetrahedron Lett*. 1998;39:4707–4710.
- Haaime G, Lohse A, Buchardt O, Nielsen PE. Peptide nucleic acids (PNAs) containing thymine monomers derived from chiral amino acids: hybridization and solubility properties of D-lysine PNA. *Angew Chem Int Ed*. 1996;35:1939–1942.
- Gourishankar A, Ganesh KN. (alpha, alpha-dimethyl)glycyl (dmG) PNAs: achiral PNA analogs that form stronger hybrids with cDNA relative to isosequential RNA. *Artif DNA PNA XNA*. 2012;3:5–13.
- Dragulescu-Andrasi A, Rapireddy S, Frezza BM, Gayathri C, Gil RR, Ly DH. A simple gamma-backbone modification preorganizes peptide nucleic acid into a helical structure. *J Am Chem Soc*. 2006;128:10258–10267.
- Englund EA, Appella DH. Synthesis of gamma-substituted peptide nucleic acids: a new place to attach fluorophores without affecting DNA binding. *Org Lett*. 2005;7:3465–3467.
- Yeh JI, Shivachiev B, Rapireddy S, et al. Crystal structure of chiral gammaPNA with complementary DNA strand: insights into the stability and specificity of recognition and conformational preorganization. *J Am Chem Soc*. 2010;132:10717–10727.
- Vieville JM, Barluenga S, Winssinger N, Delsuc MA. Duplex formation and secondary structure of gamma-PNA observed by NMR and CD. *Biophys Chem*. 2016;210:9–13.
- Verona MD, Verdolino V, Palazzesi F, Corradini R. Focus on PNA Flexibility and RNA binding using molecular dynamics and metadynamics. *Sci Rep*. 2017;7.
- Bahal R, Quijano E, McNeer NA, et al. Single-stranded gamma PNAs for in vivo site-specific genome editing via Watson-Crick recognition. *Curr Gene Ther*. 2014;14:331–342.
- Englund EA, Appella DH. Gamma-substituted peptide nucleic acids constructed from L-lysine are a versatile scaffold for multifunctional display. *Angew Chem Int Ed Engl*. 2007;46:1414–1418.
- Englund EA, Wang D, Fujigaki H, et al. Programmable multivalent display of receptor ligands using peptide nucleic acid nanoscaffolds. *Nat Commun*. 2012;3:614.
- de Koning MC, Petersen L, Weterings JJ, Overhand M, van der Marel GA, Filippov DV. Synthesis of thiol-modified peptide nucleic acids designed for post-assembly conjugation reactions. *Tetrahedron*. 2006;62:3248–3258.
- Scheibe C, Bujotzek A, Darnedde J, Weber M, Seitz O. DNA-programmed spatial screening of carbohydrate-lectin interactions. *Chem Sci*. 2011;2:770–775.
- Sforza S, Tedeschi T, Corradini R, Marchelli R. Induction of helical handedness and DNA binding properties of peptide nucleic acids (PNAs) with two stereogenic centres. *Eur J Org Chem*. 2007;5879–5885.
- Sugiyama T, Imamura Y, Demizu Y, Kurihara M, Takano M, Kittaka A. Beta-PNA: peptide nucleic acid (PNA) with a chiral center at the beta-position of the PNA backbone. *Bioorg Med Chem Lett*. 2011;21:7317–7320.
- Sacui J, Hsieh WC, Manna A, Sahu B, Ly DH. Gamma peptide nucleic acids: as orthogonal nucleic acid recognition codes for organizing molecular self-assembly. *J Am Chem Soc*. 2015;137:8603–8610.

34. Chang D, Lindberg E, Winssinger N. Critical analysis of rate constants and turnover frequency in nucleic acid-templated reactions: reaching terminal velocity. *J Am Chem Soc.* 2017;139:1444–1447.
35. Haaime G, Hansen HF, Christensen L, Dahl O, Nielsen PE. Increased DNA binding and sequence discrimination of PNA oligomers containing 2,6-diaminopurine. *Nucleic Acids Res.* 1997;25:4639–4643.
36. Eldrup AB, Christensen C, Haaime G, Nielsen PE. Substituted 1,8-naphthyridin-2 (1H)-ones are superior to thymine in the recognition of adenine in duplex as well as triplex structures. *J Am Chem Soc.* 2002;124:3254–3262.
37. Ausin C, Ortega JA, Robles J, Grandas A, Pedroso E. Synthesis of amino- and guanidino-G-clamp PNA monomers. *Org Lett.* 2002;4:4073–4075.
38. Sanders JM, Wampole ME, Chen CP, et al. Effects of hypoxanthine substitution in peptide nucleic acids targeting KRAS2 oncogenic mRNA molecules: theory and experiment. *J Phys Chem B.* 2013;117:11584–11595.
39. Kohler O, Seitz O. Thiazole orange as fluorescent universal base in peptide nucleic acids. *Chem Commun.* 2003;39:2938–2939.
40. Domling A. Recent developments in isocyanide based multicomponent reactions in applied chemistry. *Chem Rev.* 2006;106:17–89.
41. Domling A, Chi KZ, Barrere M. A novel method to highly versatile monomeric PNA building blocks by multi component reactions. *Bioorg Med Chem Lett.* 1999;9:2871–2874.
42. Maison W, Schlemminger I, Westerhoff O, Martens J. Multicomponent synthesis of novel amino acid-nucleobase chimeras: a versatile approach to PNA-monomers. *Bioorg Med Chem.* 2000;8:1343–1360.
43. Huang PC, Hsu GJ, Zhuang BR, Sung K. Novel synthesis of alpha-PNA monomers by U-4CR. *Amino Acids.* 2008;34:449–453.
44. Ovadia R, Lebrun A, Barvik I, Vasseur JJ, Baraguey C, Alvarez K. Synthesis and structural characterization of monomeric and dimeric peptide nucleic acids prepared by using microwave-promoted multicomponent reactions. *Org Biomol Chem.* 2015;13:11052–11071.
45. Ovadia R, Mondielli C, Vasseur JJ, Baraguey C, Alvarez K. Contribution to PNA-RNA chimera synthesis: one-pot microwave-assisted ugi reaction to obtain dimeric building blocks. *Eur J Org Chem.* 2017;469–475.
46. Kunz H, Pfrenkle W. Asymmetric-synthesis on carbohydrate templates - stereoselective ugi synthesis of alpha-amino-acid derivatives. *J Am Chem Soc.* 1988;110:651–652.
47. Hashimoto T, Kimura H, Kawamata Y, Maruoka K. A catalytic asymmetric ugi-type reaction with acyclic azomethine imines. *Angew Chem Int Ed.* 2012;51:7279–7281.
48. Sureshbabu VV, Narendra N, Nagendra G. Chiral N-Fmoc-beta-amino alkyl isonitriles derived from amino acids: first synthesis and application in 1-substituted tetrazole synthesis. *J Org Chem.* 2009;74:153–157.
49. Wang X, Wang QG, Luo QL. Synthesis of isonitriles from N-substituted formamides using triphenylphosphine and iodine. *Synthesis-Stuttgart.* 2015;47:49–54.
50. Narendra N, Vishwanatha TM, Nagendra G, Sureshbabu VV. Chiral N-beta-Fmoc-amino alkyl isonitriles in Ugi-4CR: an assembly of novel 1,1'-iminodicarboxylated peptidomimetics. *Tetrahedron.* 2012;68:1992–2000.
51. Chouikhi D, Ciobanu M, Zambaldo C, Duplan V, Barluenga S, Winssinger N. Expanding the scope of PNA-encoded synthesis (PES): Mtt-protected PNA fully orthogonal to fmoc chemistry and a broad array of robust diversity-generating reactions. *Chem Eur J.* 2012;18:12698–12704.
52. Katsuyama A, Matsuda A, Ichikawa S. Revisited mechanistic implications of the joullie-ugi three-component reaction. *Org Lett.* 2016;18:2552–2555.
53. Sahu B, Sacui I, Rapireddy S, et al. Synthesis and characterization of conformationally preorganized, (R)-diethylene glycol-containing gamma-peptide nucleic acids with superior hybridization properties and water solubility. *J Org Chem.* 2011;76:5614–5627.
54. Manicardi A, Calabretta A, Bencivenni M, et al. Affinity and selectivity of C2- and C5-substituted "chiral-box" PNA in solution and on microarrays. *Chirality.* 2010;22:E161–E172.
55. Songyang Z, Carraway 3rd KL, Eck MJ, et al. Catalytic specificity of protein-tyrosine kinases is critical for selective signalling. *Nature.* 1995;373:536–539.
56. Ciobanu M, Huang KT, Daguer JP, et al. Selection of a synthetic glycan oligomer from a library of DNA-templated fragments against DC-SIGN and inhibition of HIV gp120 binding to dendritic cells. *Chem Commun (Camb).* 2011;47:9321–9323.
57. Li NS, Gossai N, Naumann JA, Gordon PM, Piccirilli JA. Efficient synthetic approach to linear Dasatinib-DNA conjugates by click chemistry. *Bioconjugate Chem.* 2016.
58. Novoa A, Machida T, Barluenga S, Imberty A, Winssinger N. PNA-encoded synthesis (PES) of a 10000-member hetero-glycoconjugate library and microarray analysis of diverse lectins. *ChemBioChem.* 2014;15:2058–2065.