

A Journal of the Gesellschaft Deutscher Chemiker A Deutscher Chemiker GDCh International Edition www.angewandte.org

Accepted Article

Title: Quantitative Chirality and Concentration Sensing of Alcohols, Diols, Hydroxy Acids, Amines and Amino Alcohols using Chlorophosphite Sensors in a Relay Assay

Authors: Christian Wolf, F. Yushra Thanzeel, and Kaluvu Balaraman

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Angew. Chem. Int. Ed. 10.1002/anie.202005324

Link to VoR: https://doi.org/10.1002/anie.202005324

WILEY-VCH

COMMUNICATION

WILEY-VCH

Quantitative Chirality and Concentration Sensing of Alcohols, Diols, Hydroxy Acids, Amines and Amino Alcohols using Chlorophosphite Sensors in a Relay Assay

F. Yushra Thanzeel, Kaluvu Balaraman and Christian Wolf*^[a]

[a] Ms. F. Y. Thanzeel, Prof. Dr. K. Balaraman, Prof. Dr. C. Wolf Department of Chemistry Georgetown University 37th and O Streets, Washington, DC 20057, USA E-mail: cw27@georgetown.edu

Supporting information for this article is given via a link at the end of the document

Abstract: Analytical methods that allow simultaneous determination of the concentration and enantiomeric composition of small sample amounts and are also compatible with high-throughput multi-well plate technology have received increasing attention in recent years. We now introduce a new class of broadly useful small-molecule probes and a relay sensing strategy that together accomplish these tasks with five classes of compounds including the challenging group of monoalcohols - a scope that stands out among previously reported UV, fluorescence and CD assays. Several chlorophosphite probes and aniline indicators have been evaluated and used for on-the-fly CD/UV sensing following a continuous workflow. The wide application range of the readily available sensors is highlighted with almost 30 alcohols, diols, hydroxy acids, amines and amino alcohols, and the accuracy of the stereochemical analysis is showcased with samples covering a wide range of concentrations and enantiomeric ratios.

The introduction high-throughput experimentation of methodologies has greatly accelerated scientific discoveries and streamlined efforts in numerous academic and industrial laboratories aimed at solving complex chemical and biological tasks under strict time constraints.¹ Despite the undisputable need for quantitative screening methods that can take full advantage of generally available multiwell plate technology with parallel data acquisition and processing capabilities it is still routine to analyze one sample at a time with serial techniques.^{2,3} The shortage of experimental advance with these endeavors can at least be partially attributed to the difficulty with simultaneous determination of the concentration and enantiomeric ratio (er) of chiral samples, and these tasks are commonly accomplished separately by gravimetric analysis and chiral chromatography, respectively. Recent progress with chiroptical sensing technologies has shown that high-throughput screening of an increasing variety of chiral compounds is now possible.⁴ The most impressive examples have been achieved with chiral amines, amino alcohols, amino acids, hydroxy acids and diols.⁵ But with regard to other compound classes, for example mono-alcohols, chiroptical on-the-fly concentration and er determination needs to be demonstrated. In fact, stereochemical analysis of chiral monoalcohols by NMR spectroscopy,6 mass spectrometry7 or optical methods⁸ has remained challenging.⁹

We now wish to introduce a chiral alcohol relay sensing strategy that can accomplish three tasks, i.e. determination of the absolute configuration, sample concentration and enantiomeric composition, altogether. We have studied a series of chromophoric phosphite and amidophosphite probes that expedite comprehensive stereochemical analysis of chiral monoalcohols via simultaneous UV/CD analysis from a single sample. In addition, we prove the usefulness of this new class of sensors by demonstrating a wide application spectrum that includes alcohols but also extends to diols, hydroxy acids, amines and amino alcohols. The unique molecular recognition and chiroptical sensing features are based on irreversible formation of alkyl (amido)phosphite products exhibiting characteristic UV and circular dichroism (CD) signals that allow combined concentration and *er* analysis. While most chirality sensors introduced to date have a narrow substrate scope we show that our phosphite probes are very broadly useful.

To date, a wide range of optical chirality assays operating on the principles of dynamic covalent chemistry, in particular systems involving reversible Schiff base formation, or metal coordination, multicomponent assemblies, host-quest complexation, hydrogen bonding interactions and irreversible substrate binding have surfaced.^{4c} To the best of our knowledge, stereochemical analysis with a phosphite or amidophosphite probe has not been demonstrated. We envisioned that fast binding of alcohol substrates with aromatic chlorophosphite probes 1-5 should be possible and enable the complicated stereochemical sensing tasks outlined above, Figure 1. Although initial focus was placed on the challenging group of alcohols 6-18, the possibility of optical sensing of diols and hydroxy acids 19-22, amino alcohols 23-28 and amines 29-33 was also investigated.

The chlorobenzodioxaphosphite **1** was commercially available and the analogues **2-4** were prepared in a single step from the corresponding diols and phosphorous trichloride (see SI). The N,N-dibenzyl chlorobenzodiazaphophite **5** was synthesized in three high-yielding steps as shown in Scheme 1. The ability of the sensors **1-5** to differentiate between the enantiomers of 1-phenylethanol, **6**, and 1-phenylbutanol, **7**, was then tested under various conditions.

COMMUNICATION

Chiroptical concentration and er sensing

- On-the-fly sensing with CD/UV relay assay
- Ø Determination of sample er and total concentration
- Broadly applicable, fast & accurate
- Sensing of alcohols, diols, hydroxy acids, amino alcohols, amines
- a, amines
- Stoichiometric (1:1) sensing, no analyte excess required
 Amenable to automation and HTS equipment
- Probe structures





Amino alcohols



Figure 1. Structures of chlorophosphite probes 1-5 and chiral alcohol, diol, hydroxy acid, amino alcohol and amine target compounds 6-33. Only one enantiomer is shown.

We were pleased to find that all probes generate CD signals that are easily obtained by a mixing stoichiometric amounts of the chlorophosphite and the analyte in the presence of diisopropylethylamine, see Scheme 1 and SI. The phosphite formation was verified with several substrates by ESI-MS and NMR analysis. Importantly, the sensing reaction is very fast, it is complete within 3 minutes and readily occurs in common organic solvents, including CHCl₃, CH₂Cl₂, THF and ACN, at room temperature which underscores the potential of this sensing assay in high-throughput screening applications.



Scheme 1. Synthesis of sensor 5 (top) and CD signatures of the phosphites derived from probes 1 and 5, respectively, and the enantiomers of 1-phenylethanol, 6 (bottom). For comparison, the CD spectrum of the free alcohol (R)-6 under the same conditions is shown in orange. The CD measurements were conducted in chloroform at 0.75 mM (left) and 0.22 mM (right). See SI for details.

Encouraged by these initial findings, we decided to test the suitability of the chloroamidophosphite **5** for enantioselective recognition and analysis of the enantiomeric ratio of samples with various amounts of **6**, Table 1 and SI. In all cases, the formation of the amidophosphite **34** allowed correct identification of the absolute configuration of the major enantiomer by comparison with a reference sample and determination of the enantiomeric composition of the nonracemic samples with high accuracy. For example, the sensing of the samples containing the (*R*)- and (*S*)-**6** in a 95.0:5.0 and 35.0: 65.0 ratio gave 94.6:5.4 and 34.9:65.1, respectively, entries 1 and 4.

Table 1. Chiroptical determination of the enantiomeric ratio and absolute configuration of samples of 1-phenylethanol, 6, using probe 5.

Bn N P-Cl 5 Bn	OH 6 DIPEA, CHCl ₃	Bn N P-O 34 Bn

	San	nple comp	Sensin	g results	
Entry	Conc. (mM)	Abs. config.	er	Abs. config.ª	er ^b
1	5	R	95.0:5.0	R	94.6:5.4
2	10	S	15.0:85.0	s	18.8:81.2
3	15	R	75.0:25.0	R	78.3:21.7
4	18	S	35.0:65.0	s	34.9:65.1
5	19	R	95.0:5.0	R	95.5:4.5

^aBased on the sign of the induced Cotton effects. ^bThe *er* was calculated based on the CD signal at 300 nm. See SI for details.

COMMUNICATION



Figure 2. Representative CD responses of probes 1 and 5 to alcohols, hydroxy acids, amino alcohols and amines. All measurements were performed at 0.13-0.40 mM in chloroform. For more details, see SI.

We then continued with the evaluation of the substrate scope by applying the standard sensing protocol with probes 1 and 5 to a large variety of alcohols 6-18 including aliphatic substrates and natural products such as menthol and isopulegol. In all cases, we found that the alcohol binding induces a characteristic CD signal that can be utilized for enantioselective analysis. Representative sensing results with 11, 12, 14 and 18 are shown in Figure 2. Based on the general usefulness with mono-alcohols, we expected that the application spectrum of our sensors would extend to other important target compounds and we therefore applied our chiroptical assay to the diols, hydroxy acids, amino alcohols and amines 19-33. Again, we observed smooth phosphite formation under mild conditions in the presence of base and simultaneous induction of strong chiroptical signals at submillimolar concentrations, Figure 2. The successful testing with a total of 28 structurally diverse analytes demonstrates that chirality sensing with aromatic chlorophosphites is broadly useful and a practical tool for the determination of the absolute configuration and enantiomeric composition of several compound classes.

To develop a robust optical assay that can accomplish on-thefly sensing of the enantiomeric ratio and total concentration of chiral compounds we explored the tandem use of sensor 1 and aniline derived UV indicators. The general concept of this relay assay is shown in Scheme 2. First, the phosphite formation of an unknown amount of the alcohol substrate that may be present in up to 25.0 mM with a full equivalent of 1 matching the maximal possible analyte concentration is used for the determination of the absolute configuration and enantiomeric ratio via CD analysis as described above. The remaining excess of unreacted 1 is then captured with an aniline indicator to generate a quantifiable UV signal that is correlated to the original analyte concentration. Several candidates were screened for this purpose and the shortest reaction time together with a distinct UV change that can be used for accurate determination of the original alcohol amount were obtained with para-anisidine 38 (SI).



Scheme 2. a) Workflow of the comprehensive UV/CD sensing of 1-phenylethanol using probe 1 and *p*-anisidine as indicator. b) Chiroptical probe responses. Left: CD response of probe 1 at 273 nm to scalemic samples of 1-phenylethanol. Right: UV response at 308 nm to different concentrations of 1-phenylethanol in the presence of 1 and the indicator. See SI for details.

COMMUNICATION

This chiroptical sensing workflow was first tested with samples containing the alcohol 6 in varying amounts and enantiomeric ratios to quantitatively correlate the variation in the corresponding chiroptical readouts to the change of the analyte concentration and er. Having established the relay flow we then attempted the comprehensive sensing analysis of several alcohol mixtures. The samples were subsequently treated with the chlorophosphite and the aniline probe and the resulting mixtures containing 37 and 39 in varying quantities were then subjected to CD/UV analysis, Table 2. The results prove that the optical relay sensing concept is very reliable and generates accurate concentration and er data. The analysis of a nonracemic solution of 6 with an S:R ratio of 83.4:16.6 and a total concentration (both enantiomers combined) of 6.0 mM gave 84.1:15.9 and 6.2 mM, see entry 1. The sensing of other mixtures with vastly different amounts and enantiomeric compositions was also successful and we obtained relatively small error margins that are acceptable for high-throughput screening purposes, entries 2-7. The recording of CD and UV spectra can be accomplished simultaneously by modern CD spectrophotometers which takes full advantage of the inherent speed of the underlying reactions and the continuous relay workflow. The reproducibility and robustness of this approach were validated with samples containing similar amounts of acetophenone, see SI. We like to point out that chirality sensing with chlorophosphites is not restricted to one particular probe scaffold although the general scope and usefulness were evaluated in depth using 1 and 5. Overall, these proved equally valuable. However, 1 is commercially available which is certainly advantageous.

 Table 2. Simultaneous chiroptical sensing of the concentration, absolute configuration and enantiomeric ratio of nonracemic samples of 6.

	Sample composition			Sensing results		
Entry	Conc. (mM)	Abs. config.	er	Conc.ª (mM)	Abs. config. ^b	erc
1	6.0	S	83.4:16.6	6.2	S	84.1:15.9
2	10.0	S	85.0:15.0	9.9	S	82.9:17.1
3	13.0	R	25.0:75.0	13.8	R	22.3:77.7
4	16.0	S	71.9:28.1	15.6	S	67.5:32.5
5	18.0	R	33.3:66.7	18.6	R	34.1:65.9
6	20.0	S	62.5:37.5	20.5	S	61.8:38.2
7	24.0	S	97.9:2.1	22.5	S	96.9:3.1

^aBased on the UV change at 308 nm. ^bBased on the sign of the induced Cotton effects. ^cThe *er* was calculated based on the CD amplitude measured at 273 nm. See SI for details.

Finally, we examined if one could use an atropisomeric BINOL derived chlorophosphite probe for sensing, too. We expected that (P)-40 would have a strong CD signal originating from the BINOL moiety that may be selectively altered upon binding of either (R)- or (S)-1-phenylethanol, 6, because diastereomeric products 41 are formed. This would be fundamentally different from the use of the CD-silent chlorophosphites 1-5 which are achiral or tropos-type agents that give enantiomeric products with the chiral target compounds 6-33 thus generating induced CD effects. Interestingly, we found that sensing of enantiopure 6 and its racemate with (P)-40 yields essentially the same CD spectrum and therefore does not allow chiroptical enantiodifferentiation and ee determination which

altogether underscores the value and practicality of the sensing with **1-5** described herein, Scheme 3.



Scheme 3. Sensing of the enantiomers and a racemic mixture of alcohol **6** with (P)-**40**. The reactions were conducted at 20.0 mM and the CD spectra were collected at 0.20 mM in chloroform.

In summary, we have introduced a new class of chiroptical probes and a relay assay sensing strategy that enable accurate on-the-fly stereochemical analysis of mono-alcohols, diols, hydroxy acids, amines and amino alcohols – a variety that stands out among previously reported UV, fluorescence and CD assays. The chlorophosphite CD probes **1-5** and several aniline derived UV indicators were evaluated and the commercially available chlorobenzodioxaphosphite and para-anisidine were combined into a continuous sensing workflow. The practicality of this approach and the accuracy of the chiroptical concentration and enantiomeric ratio analysis were demonstrated with seven 1-phenylethanol samples. It is envisioned that this assay can be adapted to high-throughput equipment and multiwell plate technology which would allow fully automated operation and simultaneous screening of hundreds of samples in parallel.

Acknowledgements

We gratefully acknowledge financial support from the U.S. National Science Foundation (CHE-1764135).

Keywords: Optical sensing • chirality • circular dichroism • UV spectroscopy • chlorophosphites

- a) A. McNally, C. K. Prier, D. W. C. MacMillan, *Science* 2011, 334, 1114-1117. b) A. B. Santanilla, E. L. Regalado, T. Pereira, M. Shevlin, K. Bateman, L.-C. Campeau, J. Schneeweis, S. Berritt, Z. -C. Shi, P. Nantermet, Y. Liu, R. Helmy, C. J. Welch, P. Vachal, I. W. Davies, T. Cernak, S. D. Dreher, *Science* 2015, 347, 49-53. c) K. D. Collins, T. Gensch, F. Glorius, *Nat. Chem.* 2014, *6*, 859-871.
- [2] For examples of fast chiral HPLC enantioseparations: a) D. Kotoni, A. Ciogli, C. Molinaro, I. D'Acquarica, J. Kocergin, T. Szczerba, H. Ritchie, C. Villani, F. Gasparrini, *Anal. Chem.* 2012, *84*, 6805-6813. (b) C. L. Barhate, L. A. Joyce, A. A. Makarov, K. Zawatzky, F. Bernardoni, W. A. Schafer, D. W. Armstrong, C. J. Welch, E. L. Regalado, *Chem. Commun.* 2017, *53*, 509-512.
- [3] For recent advances with NMR chiral solvating agents, see a) L. Yang, T. Wenzel, R. T. Williamson, M. Christensen, W. Schafer, C. J. Welch, ACS Cent. Sci. 2016, 2, 332-340. b) G. Storch, M. Haas, O. Trapp, Chem. Eur. J. 2017, 23, 5414-5418. c) Q. H. Luu, K. G. Lewis, A. Banerjee, N. Bhuvanesh, J. A. Gladysz, Chem. Sci. 2018, 9, 5087-5099 and references therein.
- [4] a) D. Leung, S. O. Kang, E. V. Anslyn, *Chem. Soc. Rev.* 2012, *41*, 448-479. b) C. Wolf, K. W. Bentley, *Chem. Soc. Rev.* 2013, *42*, 5408-5424.
 For a recent Perspective on this topic: c) B. T. Herrera, S. L. Pilicer, E. V. Anslyn, L. A. Joyce, C. Wolf, *J. Am. Chem. Soc.* 2018, *140*, 10385-10401.
- [5] a) S. H. Shabbir, J. R. Clinton, E. V. Anslyn, *Proc. Natl. Acad. Sci. USA* 2009, *106*, 10487-10492. b) S. Nieto, J. M. Dragna, E. V. Anslyn, *Chem.*

COMMUNICATION

Eur. J. 2010, 16, 227-232. c) K. W. Bentley, D. Proano, C. Wolf, Nat. Commun. 2016, 7, 12539. d) E. G. Shcherbakova, E. G.; V. Brega, V. M. Lynch, T. D. James, P. Anzenbacher, Chem. Eur. J. 2017, 23, 10222-10229. e) K. W. Bentley, P. Zhang, C. Wolf, C. Science Advances 2016, 2, e1501162. f) F. Biedermann, W. M. Angew. Chem. Int. Ed. 2014, 53, 5694-5699. g) T. A. Feagin, D. P. Olsen, Z. C. Headman, J. M. Heemstra, J. Am. Chem. Soc. 2015, 137, 4198-4206. h) Z. A. De los Santos, C. Wolf, J. Am. Chem. Soc. 2016, 138, 13517-13520. i) L. A. Joyce, E. C. Sherer, C. J. Welch, Chem. Sci. 2014, 5, 2855-2861. j) F. Y. Thanzeel, K. Balaraman, C. Wolf, Nat. Comm. 2018, 9, 5323. (k) Thanzeel, F. Y. Sripada, A.; Wolf, C. J. Am. Chem. Soc. 2019, 141, 16382-16387. I) M. E. Shirbhate, S. Kwon, A. Song, S. Kim, D. Kim, H. Huang, Y. Kim, H. Lee, S.-J. Kim, M.-H. Baik, J. Yoon, K. M. Kim, J. Am. Chem. Soc. 2020, 142, 4975-4979.

- [6] a) C. Wolf, A. M. Cook, J. E. Dannatt, *Tetrahedron: Asymm.* 2014, *25*, 163-169. b) L. Yang, T. Wenzel, R. T. Williamson, M. Christensen, W. Schafer, C. J. Welch, *ACS Central Sci.* 2016, *3*, 332-340. c) G. Bian, S. Yang, H. Huang, H. Zong, L. Song, H. Fan X. Sun, *Chem. Sci.* 2016, *7*, 932-938. d) M.-S. Seo, S. Jang, H. Kim, *Chem. Commun.* 2018, *54*, 6804-6808.
- a) J. Guo, J. Wu, G. Siuzdak, M. G. Finn, *Angew. Chem. Int. Ed.* **1999**, 38, 1755-1758. b) M. T. Reetz, M. H. Becker, H. W. Klein, D. Stöckigt, *Angew. Chem. Int. Ed.* **1999**, 38, 1758-1761. Asymmetric reaction MS analysis: c) C. Markert, A. Pfaltz, *Angew. Chem. Int. Ed.* **2004**, *43*, 2498-2500. d) C. Markert, A. Pfaltz, *J. Am. Chem. Soc.* **2008**, *130*, 3234-3235. A review on this topic: e) S. Piovesana, R. Samperi, A. Lagana, M. Bella, *Chem. Eur. J.* **2013**, *19*, 11478-11494.
- [8] Examples of optical methods that allow determination of the absolute configuration of mono-alcohols: a) K. Kobayashi, Y. Asakawa, Y. Kikuchi, H. Toi, Y. Ayoma, J. Am. Chem. Soc. 1993, 115, 2648-2654. b) J. M. Lintuluoto, V. V. Borovkov, Y. Inoue J. Am. Chem. Soc. 2002, 124, 13676-13677. c) S. Havashi, M. Yotsukura, M. Noii, T. Takanami, Chem. Commun. 2015, 51, 11068-11071. d) S. Shimo, K. Takahashi, N. Iwasawa, Chem. Eur. J. 2019, 25, 3790-3794. e) T. Madry, A. Czapik, M. Kwit, ACS Omega 2019, 4, 3244-3256. Methods that allow er analysis of mono-alcohols: f) L. You, J. S. Berman, E. V. Anslyn, Nat. Chem. 2011, 3, 943–948. g) L. You, J. S. Berman, A. Lucksanawichien, . E. V. Anslyn, J. Am. Chem. Soc. 2012, 134, 7117-7125. h) L. You, G. Pescitelli, E. V. Anslvn, L. Di Bari, J. Am. Chem. Soc. 2012, 134, 7126-7134, i) C. Ni, D. Zha, H. Ye, Y. Hai, Y. Zhou. E. V. Anslyn, L. You, Angew. Chem. Int. Ed. 2018, 57, 1300-1305. j) H. H. Jo, X. Gao, L. You, E. V. Anslyn, M. J. Krische, Chem. Sci. 2015, 6, 6747-6753.
- [9] For a general overview of analytical methods: C. Wolf in *Dynamic Stereochemistry of Chiral Compounds Principles and Applications*. RSC Publishing, Cambridge, UK, **2008**, pp. 136-179.



COMMUNICATION

Entry for the Table of Contents

Chiroptical concentration and er relay sensing

- 🧹 On-the-fly sensing with CD/UV relay assay
- Ø Determination of sample er and total concentration
- 🧹 Broadly applicable, fast & accurate
- Alcohols, diols, hydroxy acids, amino alcohols, amines
- Stoichiometric (1:1) sensing, no analyte excess required



Achiral chlorophosphite probes enable accurate optical analysis of the concentration and enantiomeric ratio of chiral alcohols, diols, hydroxy acids, amino alcohols and amines. The broad utility of these readily available chiroptical sensors stands out among previously reported UV, fluorescence and CD probes which is highlighted with almost 30 examples including natural compounds.