

Photolysis of Cyclopentadienyltin Compounds: a New Source of Tin-centred Radicals

By ALWYN G. DAVIES* and MAN-WING TSE

(Christopher Ingold Laboratories, University College London, 20 Gordon Street, London WC1H 0AJ)



Summary η^1 -Cyclopentadienyltin(IV) compounds and bis-(η^5 -cyclopentadienyl)tin(II) are photolysed to give the cyclopentadienyl radical, and the former process provides a useful new route to tin(III) radicals.

RECENT work has shown that whereas the diamagnetic metallocenes, (η^5 -C₅H₅)₂M (M = Fe, Ru, or Co⁺) are photo-stable,¹ the alkylmetallocenes of the titanium group, (η^5 -C₅H₅)₂MR₂ (M = Ti, Zr, or Hf)² and (η^5 -C₅H₅)₃ThPr¹³ are photolysed with homolysis of the metal-alkyl σ -bond. We report here that, in contrast, both η^1 -C₅H₅Sn^{IV} and η^5 -C₅H₅Sn^{II} compounds are readily photolysed with preferential loss of the cyclopentadienyl group, and that the former reaction provides a useful new route to a series of Sn^{III} radicals.

If the η^1 -C₅H₅ compounds (C₅H₅)₄Sn, (C₅H₅)₂SnBu₂, C₅H₅SnBu₃, and (C₅H₅)₃SnCl are photolysed in an inert solvent in the cavity of an e.s.r. spectrometer, they all show a strong spectrum of the cyclopentadienyl radical C₅H₅·, $a(5H)$ 5.9 G (*e.g.*, Figure 1). No spectrum of the


butyl radical could be detected with the compounds (C₅H₅)₂SnBu₂ and C₅H₅SnBu₃.

The e.s.r. spectra of non-persistent tin(III) radicals in solution are difficult to observe, and attempts to use the high power conditions necessary were unsuccessful with the present compounds because the spectrum of the cyclopentadienyl radical, which overlies that of the tin radicals, is difficult to saturate. However, all these systems show the characteristic reactions of tin(III) radicals. For example, photolysis of (C₅H₅)SnBu₃ in ethylene as solvent shows the spectrum of the β -tributylstannylethyl radicals, Bu₃SnCH₂CH₂·, $a(2H_\alpha)$ 19.6, $a(2H_\beta)$ 16.0 G at -90 °C, and photolysis in the presence of alkyl bromides shows the characteristic spectra of the corresponding alkyl radicals. The reactions which are involved are illustrated in equations (1)–(3).



In reaction (3), unless the concentration of the alkyl bromide is very low, the spectrum of the cyclopentadienyl radical is broadened presumably by an exchange process, and the spectrum of the radical R'· can be observed in isolation. An example with an intermediate concentration of ethyl bromide is shown in Figure 2.

Reagents of this type provide a useful alternative to the hexa-alkylditins⁴ for the photolytic generation of R₃Sn· radicals, and should make accessible for the first time a variety of other radicals X₃Sn·, as the precursors C₅H₅-SnX₃ can readily be prepared by reaction (4).



E.s.r. experiments with the π -bonded compound (η^5 -C₅H₅)₂Sn^{II} are less convenient because it is only slightly soluble in non-polar solvents, and a yellow solid separates during photolysis. However, photolysis of a solution in toluene again gave a spectrum of the cyclopentadienyl




FIGURE 1. E.s.r. spectrum of the cyclopentadienyl radical formed by the photolysis of tricyclopentadienyltin chloride in cyclopentane at -60 °C.





FIGURE 2. E.s.r. spectrum of the ethyl radical formed by the photolysis of tributyl(cyclopentadienyl)tin in the presence of ethyl bromide in cyclopropane at -90°C . The broadened lines of the cyclopentadienyl radical in the centre of the spectrum are lost if the concentration of ethyl bromide is increased.

radical, but now the addition of ethyl bromide served only to broaden the spectrum, and no signals due to the ethyl radical could be detected. We conclude that again photolysis results in cleavage of the cyclopentadienyl group from tin [equation (5)], but the fate of the Sn^{I} radical is uncertain.



This reaction is parallel to the photolysis of the σ -bonded sterically-hindered dialkyltin compound $[(\text{Me}_3\text{Si})_2\text{CH}]_2\text{Sn}^{\text{II}}$ which has been studied by Lappert and his co-workers,⁵ but in that case only the spectrum of the persistent tin(III) radical $[(\text{Me}_3\text{Si})_2\text{CH}]_3\text{Sn}^{\cdot}$ is observed; again the fate of the Sn^{I} fragment is unknown.

We are grateful to the International Tin Research Council for the award of a Fellowship to M.-W.T.

(Received, 19th January 1978; Com. 051.)

¹ P. Borrell and E. Henderson, *Inorg. Chim. Acta*, 1975, **12**, 215.

² H. Alt and M. D. Rausch, *J. Amer. Chem. Soc.*, 1974, **96**, 5936; E. Samuel, P. Maillard, and C. Gianotti, *J. Organometallic Chem.*, 1977, **142**, 289.

³ M. D. Rausch, W. H. Boon, and H. G. Alt, *J. Organometallic Chem.*, 1977, **141**, 299; P. J. Krusic and J. K. Kochi, *J. Amer. Chem. Soc.*, 1971, **93**, 846.

⁴ J. Cooper, A. Hudson, and R. A. Jackson, *J.C.S. Perkin II*, 1973, 1056.

⁵ J. D. Cotton, C. S. Cundy, D. H. Harris, A. Hudson, M. F. Lappert, and P. W. Lednor, *J.C.S. Chem. Comm.*, 1974, 651; A. Hudson, M. F. Lappert, and P. W. Lednor, *J.C.S. Dalton*, 1976, 2369.