Enantioselective Reductive Coupling of 1,3-Enynes to Glyoxalates Mediated by Hydrogen: Asymmetric Synthesis of β , γ -Unsaturated α -Hydroxy Esters

Young-Taek Hong,[†] Chang-Woo Cho,^{†,‡} Eduardas Skucas,[†] and Michael J. Krische*,[†]

University of Texas at Austin, Department of Chemistry and Biochemistry, Austin, Texas 78712, and Kyungpook National University, Department of Chemistry, Daegu 702-701, Korea mkrische@mail.utexas.edu

Received July 2, 2007

LETTERS 2007 Vol. 9, No. 19 3745-3748

ORGANIC

ABSTRACT



Catalytic hydrogenation of 1,3-enynes 1a–8a in the presence of ethyl glyoxalate at ambient pressure and temperature using a rhodium catalyst modified by (R)-(3,5-'Bu-4-MeOPh)-MeO-BIPHEP results in highly regio- and enantioselective reductive coupling to furnish the corresponding α -hydroxy esters 1b–8b. As demonstrated by the elaboration of α -hydroxy ester 1b, the terminal and internal olefin moieties embodied by the diene side chain are subject to selective manipulation, one over the other.

The Fischer–Tropsch reaction^{1a,b} and alkene hydroformylation^{1c} are practiced on enormous scale and may be viewed as prototypical hydrogen-mediated C–C bond formations. Whereas these processes rely upon the use of carbon monoxide as a coupling partner, only recently has it been shown that catalytic hydrogenation may induce C–C coupling between π -unsaturated reactants and conventional electrophilic partners in the form of carbonyl compounds and imines.^{2–7} Through the use of cationic rhodium and iridium precatalysts, the hydrogenative coupling of conjugated enones,³ conjugated alkenes,⁴ and both conjugated^{5a–g} and nonconjugated alkynes^{5h,i,j,k} to carbonyl and imine^{3h,5c,j,k} partners have been devised. In addition, hydrogenation of 1,6-diynes, 1,6-enynes, and 1,6-alkynals is found to provide products of reductive carbocyclization.⁶ In the course of studies aimed at the development of methods for alkyne–carbonyl reductive coupling,^{8–10} highly enantioselective direct hydrogen-mediated couplings of 1,3-enynes to pyruvates and α -ketoesters using cationic rhodium catalysts modified by (*R*)-xylyl-WALPHOS were developed.^{5e} More recently, related couplings of trialkylsilyl-substituted 1,3-diynes to glyoxalates using cationic rhodium catalysts modified by (*R*)-Cl,MeO-BIPHEP were devised.^{5f} Parallel efforts toward the enantioselective reductive coupling of 1,3-enynes to glyoxalates were impeded by the fact that commercially available chelating chiral phosphine ligands

[†] University of Texas at Austin.

[‡] Kyungpook National University.

⁽¹⁾ Fischer, F.; Tropsch, H. Brennstoff Chem. 1923, 4, 276. (b) Fischer,

F.; Tropsch, H. *Chem. Ber.* **1923**, *56B*, 2428. (c) Roelen, O. German Patent DE 1938, 849,548.

⁽²⁾ For recent reviews, see: (a) Ngai, M.-Y.; Kong, J.-R.; Krische, M. J. J. Org. Chem. 2007, 72, 1063. (b) Iida, H.; Krische, M. J. Top. Cur. Chem. 2007, 279, 77.

⁽³⁾ For hydrogen-mediated reductive aldol and Mannich couplings, see:
(a) Jang, H.-Y.; Huddleston, R. R.; Krische, M. J. J. Am. Chem. Soc. 2002, 124, 15156. (b) Huddleston, R. R.; Krische, M. J. Org. Lett. 2003, 5, 1143. (c) Koech, P. K.; Krische, M. J. Org. Lett. 2004, 6, 691. (d) Marriner, G. A.; Garner, S. A.; Jang, H.-Y.; Krische, M. J. J. Org. Chem. 2004, 69, 1380. (e) Jung, C.-K.; Garner, S. A.; Krische, M. J. Org. Lett. 2006, 8, 5657. (g) Jung, C.-K.; Krische, M. J. Org. Lett. 2006, 8, 5657. (g) Jung, C.-K.; Krische, M. J. J. Am. Chem. Soc. 2006, 128, 17051. (h) Garner, S. A.; Krische, M. J. J. Org. Chem. 2007, 72, 5843.

⁽⁴⁾ For hydrogen-mediated couplings of alkenes to carbonyl compounds, see: (a) Jang, H.-Y.; Huddleston, R. R.; Krische, M. J. Angew. Chem., Int. Ed. 2003, 42, 4074. (b) Hong, Y.-T.; Barchuk, A.; Krische, M. J. Angew. Chem., Int. Ed. 2006, 128, 6885. Also see ref 7b.





uniformly failed to promote high levels of asymmetric induction. Here, we report that cationic rhodium complexes modified by (R)-(3,5-'Bu-4-MeOPh)-MeO-BIPHEP catalyze the reductive coupling of 1,3-enynes **1a**–**8a** to ethyl glyox-

alate under hydrogenation conditions to furnish the corresponding α -hydroxy esters **1b**-**8b** in good to excellent yield and with high levels of asymmetric induction (Scheme 1).

Preliminary studies involved a ligand assay, wherein 1,3envne 1a (200 mol %) was hydrogenated in the presence of ethyl glyoxalate (100 mol %) at ambient temperature and pressure in dichloroethane (0.2 M) using a catalyst derived from Rh(cod)₂OTf (5 mol %) and a chiral bidentate phosphine ligand (5 mol %). Using (R)-BINAP as the ligand, the desired product of reductive coupling 1b was obtained in 84% yield and 63% enantiomeric excess as a single regioisomer and with complete control of olefin geometry. This promising result stimulated further efforts toward the identification of highly enantioselective catalysts for the reductive coupling of enyne 1a to ethyl glyoxalate. Numerous commercially available chiral bidentate phosphine ligands were assayed under the standard coupling conditions specified above, which enabled a rough correlation between ligand structural features vis-à-vis conversion and enantioselection (Table 1). In general, any substitution at phosphorus is required for high levels of conversion. A single alkyl substituent at phosphorus, as in the case of WALPHOS type ligands, significantly decreases the isolated yield (Table 1, entries 7-9). In the BINAP, PHANEPHOS and WALPHOS series of ligands, the xylyl derivatives were slightly better at enforcing asymmetric induction than the corresponding phenyl derivatives (Table 1, entries 1-2, 3-4, and 7-9, respectively). Among the phenyl-substituted chiral chelating phosphine ligands assayed, (R)-MeO-BIPHEP enforced the highest levels of asymmetric induction (Table 1, entry 16). Stimulated by this result and the aforementioned trends, (R)-



^{*a*} Cited yields are of isolated material. Enantiomeric excess was determined via chiral stationary-phase HPLC analysis using a Chiralcel OD-H column. Eluant: 2% *i*-PrOH in hexane. Flow rate: 0.5 mL/min. Detection wavelength: 254 nm. Retention times: $t_{major} = 16.7 \text{ min}$, $t_{minor} = 18.6 \text{ min}$. NPOH = 2-naphthoic acid. TPPA = triphenylacetic acid. See the Supporting Information for further details.

(3,5-'Bu-4-MeOPh)-MeO-BIPHEP was investigated. Gratifyingly, α -hydroxy ester **1b** was obtained in 65% isolated yield and 91% enantiomeric excess (Table 1, entry 17). Further assay of the rhodium counterion, $[Rh(cod)_2]X$, where X =OTf, BF₄, and "BARF" (BARF = $B(3,5-(CF_3)_2C_6H_3)_4$), revealed no significant change in isolated yield or enantioselection in response to variation of counterion (Table 1, entries 17-19). In contrast, Brønsted acid additives were found to have a profound effect on both isolated yield and enantioselection. Brønsted acid additives are believed to facilitate the catalytic process by circumventing highly energetic 4-centered transition structures for σ -bond metathesis, as required for direct hydrogenolysis of metallacyclic intermediates, with 6-centered transition structures for hydrogenolysis of rhodium carboxylates derived upon protonolytic cleavage of the metallacycle.² Whereas 2-naphthoic acid (5 mol %) significantly increases the isolated yield of 1b, a dramatic decrease in enantioselection is observed (Table 1, entry 20). For couplings conducted in the presence of triphenylacetic acid, which provide α -hydroxy ester **1b** in 78% isolated yield and 95% enantiomeric excess, an increase in both isolated yield and enantioselection was observed (Table 1, entry 20). Accordingly, these latter conditions were adopted as standard conditions.

To assess scope, conjugated enynes **1a–8a** possessing structurally diverse acetylenic termini were coupled to ethyl

(6) For hydrogen-mediated carbocyclizations, see: (a) Jang, H.-Y.; Krische, M. J. J. Am. Chem. Soc. 2004, 126, 7875. (b) Jang, H.-Y.; Hughes, F. W.; Gong, H.; Zhang, J.; Brodbelt, J. S.; Krische, M. J. J. Am. Chem. Soc. 2005, 127, 6174. (c) Rhee, J.-U.; Krische, M. J. J. Am. Chem. Soc. 2006, 128, 10674. (d) Jung, I. G.; Seo, J.; Lee, S. I.; Choi, Y.; Chung, Y. K. Organometallics 2006, 25, 4240. (e) Smith, D. M.; Pulling, M. E.; Norton, J. R. J. Am. Chem. Soc. 2007, 129, 770 and ref 7a.

(7) Prior to our work, two isolated examples of hydrogen-mediated C–C bond formation that did not involve couplings to carbon monoxide were reported: (a) Molander, G. A.; Hoberg, J. O *J. Am. Chem. Soc.* **1992**, *114*, 3123. (b) Kokubo, K.; Miura, M.; Nomura, M. *Organometallics* **1995**, *14*; 4521.

(8) Indirect asymmetric alkyne-aldehyde reductive coupling has been achieved via stoichiometric alkyne hydrometallation (hydroboration or hydrozirconation) followed by transmetallation to afford vinylzinc reagents, which engage in enantioselective catalytic additions to aldehydes; see: (a) Oppolzer, W.; Radinov, R. Helv. Chim. Acta 1992, 75, 170. (b) Oppolzer, W.; Radinov, R. J. Am. Chem. Soc. 1993, 115, 1593. (c) Soai, K.; Takahashi, K. J. Chem. Soc., Perkin Trans. 1 1994, 1257. (d) Wipf, P.; Xu, W. Tetrahedron Lett. 1994, 35, 5197. (e) Oppolzer, W.; Radinov, R. N.; De Brabander, J. Tetrahedron Lett. 1995, 36, 2607. (f) Wipf, P.; Ribe, S. J. Org. Chem. 1998, 63, 6454. (g) Oppolzer, W.; Radinov, R. N.; El-Sayed, E. J. Org. Chem. 2001, 66, 4766. (h) Dahmen, S.; Bräse, S. Org. Lett.
 2001, 3, 4119. (i) Chen, Y. K.; Lurain, A. E.; Walsh, P. J. J. Am. Chem. Soc. 2002, 124, 12225. (j) Ji, J.-X.; Qiu, L.-Q.; Yip, C. W.; Chan, A. S. C. J. Org. Chem. 2003, 68, 1589. (k) Lurain, A. E.; Walsh, P. J. J. Am. Chem. Soc. 2003, 125, 10677. (1) Ko, D.-H.; Kang, S.-W.; Kim, K. H.; Chung, Y.; Ha, D.-C. Bull. Kor. Chem. Soc. 2004, 25, 35. (m) Sprout, C. M.; Richmond, M. L.; Seto, C. T. J. Org. Chem. 2004, 69, 6666. (n) Jeon, S.-J.; Chen, Y. K.; Walsh, P. J. Org. Lett. 2005, 7, 1729. (o) Jeon, S.-J.; Fisher, E. L.; Carroll, P. J.; Walsh, P. J. J. Am. Chem. Soc. 2006, 128, 9618. (p) Lauterwasser, F.; Gall, J.; Hoefener, S.; Bräse, S. Adv. Synth. Catal. 2006, 348, 2068.

glyoxalate under standard conditions (Figure 1). In all cases, C-C coupling occurs at the acetylenic terminus of the conjugated enyne to furnish α -hydroxy esters **1b-8b** as single constitutional isomers with complete control of alkene geometry. The coupling was applicable to conjugated enynes possessing aryl (**1a**, **2a**, **3a**), heteroaryl (**4a**, **8a**), primary alkyl (**6a**, **7a**), and secondary alkyl (**5a**) substituents at the acetylenic terminus. Isolated yields of the α -hydroxy esters **1b**-**8b** ranged from 70 to 82%, and the levels of enantioselection ranged from 86 to 97% ee (Figure 1). Notably, over-



Figure 1. Enantioselective catalytic reductive coupling of substituted 1,3-enynes to ethyl glyoxalate mediated by hydrogen. Cited yields are of isolated material. See the Supporting Information for detailed experimental procedures. Enantiomeric excess was determined via chiral stationary phase HPLC analysis.

reduction of the diene side chain of adducts 1b-8b was not observed under the conditions of hydrogen-mediated coupling. Absolute stereochemical assignment of α -hydroxy esters 1b-8b is based upon X-ray diffraction analysis of the amide derived from 1b and (*R*)-1-(2-naphthyl)ethylamine.

Each olefin of the diene side chain of the reductive coupling products 1b-8b is subject to selective manipulation. To illustrate, α -hydroxy ester 1b was subjected to the

⁽⁵⁾ For hydrogen-mediated couplings of alkynes to carbonyl compounds and imines, see: (a) Huddleston, R. R.; Jang, H.-Y.; Krische, M. J. J. Am. Chem. Soc. 2003, 125, 11488. (b) Jang, H.-Y.; Huddleston, R. R.; Krische, M. J. J. Am. Chem. Soc. 2004, 126, 4664. (c) Kong, J.-R.; Cho, C.-W.; Krische, M. J. J. Am. Chem. Soc. 2005, 127, 11269. (d) Cho, C.-W.; Krische, M. J. Org. Lett. 2006, 8, 891. (e) Kong, J.-R.; Ngai, M.-Y.; Krische, M. J. J. Am. Chem. Soc. 2006, 128, 718. (f) Cho, C.-W.; Krische, M. J. Org. Lett. 2006, 8, 3873. (g) Komanduri, V.; Krische, M. J. J. Am. Chem. Soc. 2006, 128, 16448. (h) Kong, J.-R.; Krische, M. J. J. Am. Chem. Soc. 2006, 128, 16448. (h) Kong, J.-R.; Krische, M. J. J. Am. Chem. Soc. 2006, 128, 16448. (h) Kong, J.-R.; Kong, J.-R.; Krische, M. J. J. Am. Chem. Soc. 2006, 129, 280. (j) Skucas, E.; Kong, J.-R.; Krische, M. J. J. Am. Chem. Soc. 2007, 129, 242. (k) Barchuk, A.; Ngai, M.-Y.; Krische, M. J. J. Am. Chem. Soc. 2007, 129, 8432.

⁽⁹⁾ For indirect catalytic enantioselective alkyne-ketone reductive couplings analogous to those described in the preceding reference, see: (a) Li, H.; Walsh, P. J. J. Am. Chem. Soc. **2004**, *126*, 6538. (b) Li, H.; Walsh, P. J. J. Am. Chem. Soc. **2005**, *127*, 8355. (c) Jeon, S.-J.; Li, H.; Garcia, C.; La Rochelle, L. K.; Walsh, P. J. J. Org. Chem. **2005**, *70*, 448.

⁽¹⁰⁾ Direct alkyne-aldehyde reductive coupling to furnish allylic alcohols has been achieved under the conditions of nickel catalysis; see: (a) Oblinger, E.; Montgomery, J. J. Am. Chem. Soc. **1997**, *119*, 9065. (b) Huang, W.-S.; Chan, J.; Jamison, T. F. Org. Lett. **2000**, *2*, 4221. (c) Mahandru, G. M.; Liu, G.; Montgomery, J. J. Am. Chem. Soc. **2004**, *126*, 3698. (d) For enyne coupling, see: Miller, K. M.; Luanphaisarnnont, T.; Molinaro, C.; Jamison, T. F. J. Am. Chem. Soc. **2004**, *126*, 4130. (e) Miller, K. M.; Jamison, T. F. Org. Lett. **2005**, *7*, 3077.

following transformations. Hydroxy-directed epoxidation of **1b** catalyzed by $OV(acac)_2^{11}$ stereoselectively delivers vinyl epoxide **1c**. Oxidative cleavage of the diene terminus using the Lemieux–Johnson protocol furnishes the conjugated enal **1e**¹² Exhaustive reduction of the diene side chain to form the saturated α -hydroxy ester **1d** is achieved upon hydrogenation of **1b** using Crabtree's catalyst.¹³ Finally, exposure of diene **1b** to conditions for ruthenium catalyzed cross metathesis with 1,4-diacetoxy-2-butene¹⁴ enables formation of allylic acetate **1f**. The structural assignment of **1c-1f** is made in analogy to the corresponding pyruvate coupling products (Figure 2).^{5e}



Figure 2. Elaboration of coupling product 1b.

In summary, we report a regio- and enantioselective hydrogenative coupling of substituted 1,3-enynes to ethyl glyoxalate catalyzed by rhodium. Good to excellent levels of asymmetric induction are achieved using (R)-(3,5-'Bu-4-MeOPh)-MeO-BIPHEP as chiral ligand with triphenylacetic acid as a Brønsted acid cocatalyst. A stereochemical model accounting for the high levels of enantioselection enforced by (R)-(3,5-'Bu-4-MeOPh)-MeO-BIPHEP will be disclosed in due course. These studies serve as a prelude to our longterm goal of developing hydrogenation catalysts enabling reductive C-C coupling of simple, commercially available nonconjugated alkynes to diverse aldehydes en route to optically enriched allylic alcohols.

Acknowledgment is made to the Robert A. Welch Foundation, Johnson & Johnson, Merck, and the NIH-NIGMS (RO1-GM69445) for partial support of this research. Drs. Hans Ulrich Bläser, Matthias Lotz, and Marc Thommen of Solvias are thanked for a generous donation of (R)-3,5-'Bu-4-MeOBIPHEP.

Supporting Information Available: Spectral data for all new compounds (¹H NMR, ¹³C NMR, IR, HRMS). Single-crystal X-ray diffraction data corresponding to the amide derived from **1b** and (R)-(+)-1-(2-naphthyl)ethylamine. This material is available free of charge via the Internet at http://pubs.acs.org.

OL7015548

^{(11) (}a) Sharpless, K. B.; Michaelson, R. C. J. Am. Chem. Soc. **1973**, 95, 6136. (b) For a review, see: Sharpless, K. B.; Verhoeven, T. R. Aldrichim. Acta **1979**, 12, 63.

^{(12) (}a) Lemieux; R. U.; von Rudloff, E. *Can. J. Chem.* **1955**, *33*, 1701.
(b) Pappo, R.; Allen, D. S., Jr.; Lemieux;, R. U.; Johnson, W. S. *J. Org. Chem.* **1956**, *21*, 478.

^{(13) (}a) Crabtree, R. H.; Felkin, H.; Morris, G. E. J. Organomet. Chem. **1977**, 141, 205. (b) For a review, see: Crabtree, R. H. Acc. Chem. Res. **1979**, 12, 331. (c) For hydrogenation of conjugated dienes catalyzed by iridium, see: Cui, X.; Burgess, K. J. Am. Chem. Soc. **2003**, 125, 14212.

⁽¹⁴⁾ Funk, T. W.; Efskind, J.; Grubbs, R. H. Org. Lett. 2005, 7, 187.