



Cite this: *Org. Biomol. Chem.*, 2018, **16**, 9171

Syntheses and *in vitro* biological evaluation of S1PR1 ligands and PET studies of four F-18 labeled radiotracers in the brain of nonhuman primates†

Zonghua Luo,^a Junbin Han,^a Hui Liu,^a Adam J. Rosenberg,^a Delphine L. Chen,^a Robert J. Gropler,^a Joel S. Perlmutter^{a,b} and Zhude Tu^{a,*}

A series of seventeen hydroxyl-containing sphingosine 1-phosphate receptor 1 (S1PR1) ligands were designed and synthesized. Their *in vitro* binding potencies were determined using [³²P]S1P competitive binding assays. Compounds **10a**, **17a**, **17b**, and **24** exhibited high S1PR1 binding potencies with IC₅₀ values ranging from 3.9 to 15.4 nM and also displayed high selectivity for S1PR1 over other S1P receptor subtypes (IC₅₀ > 1000 nM for S1PR2–5). The most potent compounds **10a**, **17a**, **17b**, and **24** were subsequently radiolabeled with F-18 in high yields and purities. MicroPET studies in cynomolgus macaque showed that [¹⁸F]**10a**, [¹⁸F]**17a**, and [¹⁸F]**17b** but not [¹⁸F]**24** crossed the blood brain barrier and had high initial brain uptake. Further validation of [¹⁸F]**10a**, [¹⁸F]**17a**, and [¹⁸F]**17b** in preclinical models of neuroinflammation is warranted to identify a suitable PET radioligand to quantify S1PR1 expression *in vivo* as a metric of an inflammatory response.

Received 22nd October 2018,
Accepted 15th November 2018

DOI: 10.1039/c8ob02609b

rsc.li/obc

Introduction

Sphingosine 1-phosphate (S1P) is a membrane-derived lysophospholipid that plays a critical regulatory role in inflammatory diseases.^{1–3} S1P acts by modulating five highly specific G protein-coupled receptors (GPCRs, S1PR1–5) and has high binding affinities (nanomolar) with these five receptors.^{4–8} The unique patterns of cellular and temporal expression determine the specific roles of S1PR1–5 in each organ system. Modulation of these receptors is particularly important in the central nervous, cardiovascular, and immune systems. Among these five-receptor subtypes, S1PR1 is one of the most abundant receptors in the entire GPCR superfamily. S1PR1 was originally called the endothelial differentiation growth factor receptor 1 (EDG-1). It is particularly enriched in endothelial cells and vascular smooth muscle cells, and also expressed on immune cells such as lymphocytes.⁹ The relevance of S1PR1 in clinical disease has become readily apparent with the U.S. Food and Drug Administration (FDA) approval of the S1PR1 modulator fingolimod (FTY720) for treating relapsing-

remitting multiple sclerosis (RR-MS). MS is a chronic autoimmune, inflammatory disease caused by lymphocytic infiltration that leads to a demyelinating neurodegenerative disease with no cure. RR-MS is the most common form, characterized by intermittent relapses that lead to increasing functional disability that in large part may be mediated by neuroinflammatory responses in the brain.¹⁰ Fingolimod, structurally similar to S1P, functionally modulates S1PR1 on lymphocytes, leading to S1PR1 downregulation and subsequent lymphocyte trapping in lymph nodes. This reduces the peripheral blood lymphocyte count and presumably lymphocyte infiltration in the brain. The effects of fingolimod on S1PR1 expression in the central nervous system (CNS) itself and its contribution to treatment efficacy are not well understood. S1PR1 is expressed in neurons and glia in the CNS, including astrocytes, which modulate inflammatory responses throughout the gray and white matter; microglia, the resident macrophages in the CNS, and oligodendrocytes, which produce the myelin needed for nerve conduction. In the *in vitro* models of demyelination induced by lysophosphatidylcholine (LPS), fingolimod reduced demyelination and neuronal process extension in the CNS, presumably through modulation of S1PR1.^{11–13} It also inhibited microglial activation and reduced oligodendrocyte apoptosis.¹³ Specific deletion of S1PR1 in astrocytes resulted in decreased experimental autoimmune encephalomyelitis (EAE) pathology *in vivo* and a loss of fingolimod efficacy, suggesting that fingolimod also acts on S1PR1 in astrocytes.^{14,15} Fingolimod-treated MS patients exhibited reduced brain volume loss, lesion

^aDepartment of Radiology, Washington University School of Medicine, 510 South Kingshighway Boulevard, St Louis, Missouri, 63110, USA. E-mail: tuz@mir.wustl.edu; Tel: +1-314-362-8487

^bDepartments of Neurology, Neuroscience, Physical Therapy and Occupational Therapy, Washington University School of Medicine, St Louis, MO 63110, USA

†Electronic supplementary information (ESI) available. See DOI: 10.1039/c8ob02609b

numbers, and severity,^{14,16–18} confirming the *in vitro* and animal model findings that S1PR1 pathways play important roles in anti-inflammation and neuroprotection. A S1PR1-targeted positron emission tomography (PET) radiotracer would help us understand the pathophysiology of S1PR1 in the CNS. Imaging S1PR1 *in vivo* could be used to develop therapies that improve upon the initial treatment successes with fingolimod. Furthermore, S1PR1 is a therapeutic target of great interest for a number of other inflammatory diseases, including inflammatory bowel disease¹⁹ and atherosclerotic disease,^{20,21} as well as cancer, where S1PR1 expression promotes treatment resistance^{22,23} and enhances metastatic potential.^{23–26} S1PR1 imaging would enable mechanistic studies and S1PR1-specific readouts of targeted therapy in a wide range of inflammatory diseases. Using PET with the promising C-11 radiotracer [¹¹C]TZ3321, we demonstrated increased S1PR1 expression at inflammatory response sites in three animal models: the rat EAE model of MS, the ApoE^{−/−} mouse femoral artery wire-injury model of neointimal hyperplasia, and the rat carotid injury model of vascular inflammation.^{27–29} The longer half-life of F-18 ($T_{1/2}$ = 109.8 min) reduces time constraints on tracer production, distribution, and permits longer imaging sessions for higher target-to-reference ratios than is possible with C-11. Therefore, the PET radiochemistry community has focused on the development of potent and selective [¹⁸F]-labeled S1PR1 radiotracers to quantitatively measure the changes of S1PR1 expression in response to inflammation. Several [¹⁸F]-labeled small molecules have been characterized for their suitability to be [¹⁸F]-labeled PET radiotracers for imaging S1PR1 *in vitro* and *in vivo* (Fig. 1). However, either they had inadequate *in vivo* metabolic stability or did not penetrate the blood brain barrier (BBB). For instance, three [¹⁸F]-labeled fingolimod derivatives, [¹⁸F]3, [¹⁸F]4a, [¹⁸F]4b reported by Haufe's group showed fast kinetics, low selectivity, or defluorination *in vivo*, which prevented their transfer into

clinical investigation.^{30,31} Our group explored oxadiazole-containing analogues and reported two [¹⁸F]fluoroethoxy radiotracers, one having an azetidine-3-carboxylic acid ([¹⁸F]5), and another tracer having a 1H-1,2,3-triazole-4-methanol ([¹⁸F]6).^{32,33} Although the *in vivo* study suggested that the mouse liver uptake of [¹⁸F]5 had a strong positive correlation with acute liver inflammation induced by LPS injection, this radiotracer has limited penetration of the BBB for imaging neuroinflammation. Our preliminary data revealed that [¹⁸F]6 exhibited a high rat brain uptake with 0.71%ID per gram at 60 min post injection and no *in vivo* defluorination. A specific S1PR1 ligand reduced the uptake of [¹⁸F]6 binding to S1PR1 in tissue sections of the inflamed brain from an LPS-induced neuroinflammation mouse model. But follow-up microPET imaging studies of [¹⁸F]6 in cynomolgus macaque suggested this radiotracer did not penetrate the nonhuman primate (NHP) brain (seen in ESI†). Species differences in the BBB penetration ability of PET radiotracers have been previously reported for other PET neuroimaging radiotracers.³⁴ Consequently, neither [¹⁸F]5 nor [¹⁸F]6 is a suitable PET S1PR1 radiotracer to assess the neuroinflammatory responses in humans. To overcome the above-mentioned challenges in the identification of a suitable [¹⁸F]-labeled S1PR1 radiotracer for neuroinflammation response in CNS disorders, we focused on further optimization of the oxadiazole analogues by replacing the carboxylic acid moiety with hydroxyl group(s) in the terminal of substituted side chain. Our previous rat biodistribution of [¹⁸F]6 showed good brain uptake compared to carboxylic acid containing analogues. Herein, we reported the syntheses of new serial of terminal hydroxyl-containing S1PR1 compounds, *in vitro* binding assays to determine their binding potencies and selectivity for S1PR1, radiosyntheses of four F-18 radiotracers, and microPET studies of these four radiotracers in the cynomolgus macaque brain to investigate their *in vivo* kinetics and ability to penetrate the BBB.




Fig. 1 Structures of fingolimod and S1PR1 radiotracers.

Results and discussion


To identify an [^{18}F]-labeled radiotracer with high potency, high selectivity, and the capability to penetrate the BBB. Initially, the (1,2,3-triazol-4-yl)methanol moiety in compound **6** was replaced by hydroxyl methyl functional group to generate compound **10a**, we then replaced the trifluoromethyl group with a cyano or hydrogen group (Scheme 1). Next, we turned our attention to retaining the trifluoromethyl group and replaced the hydroxyl methyl group of **10a** with alkyl alcohols, amino alcohols, and ether alcohols. In total, seventeen new analogues were designed, synthesized and screened for *in vitro* and *in vivo* properties. The syntheses of the target compounds are depicted in Schemes 1–5 following reported procedures with necessary modifications.^{32,33} For compounds **10a–c**, syntheses were initiated with 4-(hydroxymethyl)-benzonitrile (**7**) treated with hydroxylamine hydrochloride in presence of NaHCO_3 to yield amidoxime **8**. Condensation and cyclization of the amidoxime **8** with aromatic carboxylic acids **9a–c** in DMF in presence of 2-(1*H*-benzotriazole-1-yl) 1,1,3,3-tetra-methylaminium tetrafluoroborate (TBTU) and a catalytic amount of 1-hydroxybenzo-triazole hydrate (HOBt) to generate compounds **10a–c** (Scheme 1).

Compounds **13a–c** were synthesized by employing a similar procedure described for compounds **10a–c**. Treatment of compounds **11a–c** with hydroxylamine hydrochloride yielded the intermediates **12a–c**, which were subjected to condensation and then cyclization with compound **9a** to yield compounds **13a–c**. Compound **13d** was prepared by deprotection of *tert*-butyloxy-carbonyl (Boc) group using 4.0 M HCl in dioxane (Scheme 2).


Compound **10a** was utilized as a key intermediate for the syntheses of analogues **15a–b** and **17a–g**. Swern oxidation of the primary alcohol in compound **10a** with oxalyl chloride in the presence of triethylamine at -78°C to yield the intermediate aldehyde **14**,³² followed by reductive amination using corresponding amines generated the amino alcohol compounds **15a–b** (Scheme 3).

Chlorination of compound **10a** was achieved by reaction with cyanuric chloride in DMF to afford the intermediate chloride **16**, followed by nucleophilic substitution with various alcohols to yield compounds **17a–e** and **17g**. Compound **17f** was generated by deprotection of Boc group of **17e** (Scheme 3).




Compounds **20a–b** were synthesized using the corresponding 4-(2-hydroxy-ethoxy)benzonitrile (**18a**) or 4-((2,3-dihydroxy-propoxy)methyl)benzonitrile (**18b**) to generate amidox-



Scheme 1 Syntheses of **10a–c**. Reagents and conditions: (a) Hydroxylamine hydrochloride, NaHCO_3 , methanol, reflux; (b) TBTU, HOBt, DIPEA, DMF, RT– 120°C .





















Scheme 2 Syntheses of **13a–d**. Reagents and conditions: (a) Hydroxylamine hydrochloride, NaHCO_3 , methanol, reflux; (b) **9a**, TBTU, HOBt, DIPEA, DMF, RT– 120°C ; (c) 4.0 M HCl in dioxane, RT.



In the third series, we PEGylated compound **10a** using different PEG units. PEGylation has been reported as an

Table 1 Structures and binding potencies (mean \pm SD) of S1P and newly synthesized compounds against S1PR1^a

							
Compound	R ¹	R ²	S1PR1 IC ₅₀ \pm SD (nM)	Compound	R ¹	R ²	S1PR1 IC ₅₀ \pm SD (nM)
S1P	—	—	1.4 \pm 0.3	17a	CF ₃		14.0 \pm 0.4
10a	CF ₃		6.7 \pm 0.7 ^b	17b	CF ₃		15.4 \pm 3.3
10b	CN		15.4 \pm 3.8	17c	CF ₃		106 \pm 25
10c	H		>1000	17d	CF ₃		125 \pm 22
(±)-13a	CF ₃		39.9 \pm 3.6	17f	CF ₃		75.6 \pm 19.3
13b	CF ₃		38.0 \pm 6.0	17g	CF ₃		130 \pm 19
13d	CF ₃		48.2 \pm 9.7	20a	CF ₃		56.6 \pm 23.8
15a	CF ₃		87.2 \pm 17.7	(±)-20b	CF ₃		23.8 \pm 5.8
15b	CF ₃		125 \pm 27	(±)-24	CF ₃		3.9 \pm 0.5

^a IC₅₀ values were determined with at least 3 independent experiments, each run was performed in duplicate. ^b Ref. 32.

effective strategy for improving the solubility, stability, pharmacodynamics, and pharmacokinetics of drug candidates interacting with biologically interesting proteins or peptides.^{36–42} The addition of a small number of PEGylaton units ($n = 2, 3, 4$) is frequently used to improve a compound's hydrophilicity which may facilitate crossing the BBB. As one example, [¹⁸F]AV45, the well-known radiotracer used for imaging β -amyloid (A β) aggregates in the AD brain contains three PEGylation units.⁴¹ Therefore, we next synthesized compounds **17a–d** with 1–4 PEGs and tested their binding potency. Our *in vitro* binding data showed that compound **17a** and **17b** with one and two PEG units had similar binding potencies with IC₅₀ values of 14.0 and 15.4 nM, respectively. Compound **17c** (IC₅₀ = 106 nM) with three PEGs, had approximately 16-fold lower potency compared to **10a**. Compound **17d** (IC₅₀ = 125 nM) with four PEGs also showed further decreased binding potency.

The *in vitro* S1P binding data of these new PEG-containing compounds suggested that the addition of one or two PEG units retained S1PR1 binding potency, but the extension with more PEG chains (>2 units) decreased S1PR1 binding potency. In addition, for compounds **17f** and **17g**, in which a secondary or tertiary amine was introduced in the PEG chains, S1PR1 binding potencies decreased with IC₅₀ values of 75.6 and 130 nM, respectively. Finally, we modified compound **20a** with a 2-hydroxyethoxyl group directly linked to a phenyl group, this approach also reduced binding potency (IC₅₀ = 56.6 nM) when compared to compound **17a** with a 2-hydroxy ethoxyl linked with a benzyl group. Further optimization of **17a** was conducted by introducing a terminal hydroxyl group using glycerin to yield compound (±)-**20b**, which showed an IC₅₀ value of 23.8 nM, which is less potent than the glycol tail compound **17a**. The alkyl di-alcohol compound (±)-**24**, which incorporated


an ethylene glycol directly linked with phenyl group was found to be the most potent S1PR1 ligand of this study with an IC₅₀ value of 3.9 nM. Because compounds **10a**, **17a**, **17b**, and (±)-**24** each had high binding potency for S1PR1 with IC₅₀ < 20 nM, these four compounds were screened to determine their binding potency to other S1P receptor subtypes S1PR2, 3, 4 and 5 to check their binding selectivity. Our *in vitro* data shown in Table 2 revealed that all four compounds had no significant binding toward S1PR2–5 (IC₅₀ > 1000 nM), indicating that they are selective for S1PR1 over other S1PR subtypes S1PR2, 3, 4, and 5.

Consequently, these four compounds were radiolabeled with F-18 to check their potential to be neuroimaging PET radiotracers for quantifying S1PR1 expression in the brain. Prior to radiolabeling, the corresponding precursors **27a–d** were synthesized by following Scheme 6. Methoxymethyl (MOM) protected hydroxybenzimidamide **25a–d** were prepared

Table 2 Binding potencies (mean \pm SD) of **10a**, **17a–b**, and (±)-**24** toward S1PR1–5 and Clog *P* values

Compd.	IC ₅₀ ^a (nM)					Clog <i>P</i> ^b
	S1PR1	S1PR2	S1PR3	S1PR4	S1PR5	
S1P	1.4 \pm 0.3	3.6 \pm 0.5	0.4 \pm 0.2	151 \pm 82	3.1 \pm 1.1	
10a	6.7 \pm 0.7	>1000	>1000	>1000	>1000	4.00
17a	14.0 \pm 0.4	>1000	>1000	>1000	>1000	4.02
17b	15.4 \pm 3.3	>1000	>1000	>1000	>1000	3.85
(±)-24	3.9 \pm 0.5	>1000	>1000	>1000	>1000	3.09

^a IC₅₀ values were determined with at least 3 independent experiments, each run was performed in duplicate; assays for compounds which showed no activity (IC₅₀ > 1000 nM) were repeated twice. ^b Clog *P* values were calculated by PerkinElmer ChemDraw® 16.0.1.4 (77).



Scheme 6 General syntheses of precursors **27a–d** and radiotracers [^{18}F]**10a**, [^{18}F]**17a**, [^{18}F]**17b**, and (±)-[^{18}F]**24**. Reagents and conditions: (a) 4-Hydroxy-3-(trifluoromethyl)benzoic acid, TBTU, HOBT, DIPEA, DMF, RT–120 °C; (b) ethylene ditosylate, K_2CO_3 , CH_3CN , 90 °C; (c) [^{18}F]KF, Kryptofix 222, K_2CO_3 , CH_3CN , 110 °C, 15 min, 6 N HCl, 5 min.

by following the general procedure described for **12a–c** with MOM protected benzonitriles. The condensation and cyclization of **25a–d** and 4-hydroxy-3-(trifluoro-methyl)benzoic acid afforded **26a–d**, which were subjected to *O*-alkylation with ethylene ditosylate to afford MOM-protected precursors **27a–d**. The radiosyntheses of [^{18}F]**10a**, [^{18}F]**17a**, [^{18}F]**17b**, and (±)-[^{18}F]**24** were successfully achieved by using nucleophilic reaction between each tosylate precursor and [^{18}F]KF in acetonitrile with Kryptofix 222, followed by deprotection of MOM group. The radioactive products were purified using a semi-preparative high performance liquid chromatography (HPLC) and then formulated using 10% of ethanol saline solution. Each dose of [^{18}F]**10a**, [^{18}F]**17a**, [^{18}F]**17b**, and (±)-[^{18}F]**24** was authenticated by co-injecting with corresponding cold standard compound **10a**, **17a**, **17b**, and (±)-**24**, respectively. These four radiotracers were achieved with radiochemical yields of 35–40%, radiochemical purities >98%, and molar activities >74 GBq μmol^{-1} (decay corrected to end of synthesis, EOS).

To check if each radiotracer had the capability to penetrate the BBB with sufficient accumulation in the animal brain, microPET studies of the brain of a cynomolgus macaque were

performed for each F-18 labeled radiotracer on different days. Macaque did not have studies more frequently than every 2 weeks to allow the subject to recover. Time activity curves (TACs) showing brain uptake and washout from our microPET studies of these four F-18 radiotracers are shown in Fig. 2, brain TACs indicated that three radiotracers [^{18}F]**10a**, [^{18}F]**17a**, and [^{18}F]**17b** were able to cross the BBB and had high initial brain uptake in the nonhuman primates.

The brain standard uptake value (SUV) for [^{18}F]**10a**, [^{18}F]**17a**, and [^{18}F]**17b** reached a maximum at 4–6 min after tracer injection. [^{18}F]**10a** had the highest initial brain uptake and quick washout kinetics from the brain while the brain washout kinetics of [^{18}F]**17b** was relatively slow. For the most potent compound (±)-**24**, with an IC_{50} value of 3.9 nM, our microPET studies of (±)-[^{18}F]**24** indicated that it did not penetrate the BBB and its brain accumulation was very low. Lipophilicity of small molecule drug is a key parameter affecting penetration of the BBB. $\log P$ value is an index of small molecular lipophilicity. Most drugs that are active in the central nervous system have $\log P$ values that range from 1.5 to 2.7,⁴³ though some have values beyond this range. The calculated $\log P$ (Clog P)




Fig. 2 Representative time-activity curves of microPET studies for [^{18}F]**10a**, [^{18}F]**17a**, [^{18}F]**17b** and (±)-[^{18}F]**24** in brains of nonhuman primates. The curves indicated that three radiotracers [^{18}F]**10a**, [^{18}F]**17a** and [^{18}F]**17b** achieved the max SUV value fast, at 4–6 min post injection. [^{18}F]**10a** had the highest initial brain uptake while [^{18}F]**17b** exhibited relatively slow washout kinetics. (±)-[^{18}F]**24** displayed the lowest initial uptake in the brain, indicating it has no capability in penetrating the BBB of nonhuman primate.

values (Table 2) of these four tracers were 4.00, 4.02, 3.85, and 3.09 for [^{18}F]**10a**, [^{18}F]**17a**, [^{18}F]**17b**, and (\pm)-[^{18}F]**24**, respectively. The most potent radiotracer (\pm)-[^{18}F]**24** (IC_{50} value of 3.9 nM for S1PR1) had a log P value of 3.09 which would predict better BBB penetration. The unanticipated low nonhuman primate brain uptake of (\pm)-[^{18}F]**24** despite its lipophilicity may have resulted from the two hydroxyl groups causing different *in vivo* pharmacological properties, such as binding to P-glycoprotein (Pgp).⁴⁴ Further investigation would be needed to understand the inability of (\pm)-[^{18}F]**24** in penetrating BBB.

Together, our microPET studies in nonhuman primates suggested that radiotracers [^{18}F]**10a**, [^{18}F]**17a**, and [^{18}F]**17b** have potential to be promising S1PR1 PET radiotracers for imaging the neuroinflammatory response *in vivo*. Further investigations in animal models of neuroinflammatory diseases are warranted to characterize their radiopharmaceutical properties.

Conclusions

In conclusion, we developed a series of new S1PR1 ligands and screened *in vitro* binding potencies for all the newly synthesized compounds. Compounds **10a**, **17a**, **17b**, and (\pm)-**24** exhibited high potency ($\text{IC}_{50} < 20$ nM) and high selectivity for S1PR1 ($\text{IC}_{50} > 1000$ nM over S1PR2, 3, 4, and 5). The radio-syntheses of [^{18}F]**10a**, [^{18}F]**17a**, [^{18}F]**17b**, and (\pm)-[^{18}F]**24** were accomplished with good yields (35–40%), high radiochemical and chemical purities (>98%), and reasonable molar activities (>74 GBq μmol^{-1}). MicroPET studies in cynomolgus macaque indicated that [^{18}F]**10a**, [^{18}F]**17a**, and [^{18}F]**17b** can cross the BBB of NHPs and have good brain uptake while (\pm)-[^{18}F]**24** was not able to penetrate the BBB. Further *in vivo* evaluation of [^{18}F]**10a**, [^{18}F]**17a**, and [^{18}F]**17b** in animal models of inflammatory disease mechanisms is needed to check the suitability of these radiotracers for assessing the neuroinflammatory response by measuring the S1PR1 expression prior to translation of a lead radiotracer to clinical investigations.

Experimental section

General information

All reagents and chemicals were obtained from standard commercial sources and used without further purification unless otherwise stated. Organic reactions were carried out under inert nitrogen and moisture-free with dry solvent. Thin layer chromatography (TLC) was used to monitor the reaction. Final organic products were purified by flash column chromatography using 230–400 mesh silica gel purchased from Silicycle. Melting points were determined on a MEL-TEMP 3.0 apparatus without correction. All deuterated solvents were purchased from Cambridge Isotope Laboratories. ^1H and ^{13}C NMR spectra were recorded on a 400 MHz Varian instrument. Chemical shifts were reported in parts per million (ppm) and were calibrated using a residual undeuterated solvent as an

internal reference (CDCl_3 : δ 7.26 ppm; CD_3OD : δ 3.31 ppm; $\text{DMSO}-d_6$: δ 2.50 ppm; $\text{acetone}-d_6$: δ 2.05 ppm). Multiplicities are indicated by s (singlet), d (doublet), t (triplet), q (quartet), p (pentet), h (hextet), m (multiplet) and br (broad). High-resolution positive ion mass was acquired by a Bruker Maxis 4G Q-TOF mass spectrometer with electrospray ionization source. The welfare of the animals conformed to the requirements of National Institutes of Health (NIH). This work was conducted at the NHP Facility of Washington University in St Louis with approval from the IACUC.

(*E*)-*N'*-Hydroxy-4-(hydroxymethyl)benzimidamide (8). Synthesis of compound **8** followed the published procedure.³³

General procedure for the synthesis of 10a–c

To a round-bottom flask equipped with a stir bar was added acid **9a–c** (1.0 eq.), HOBT (0.2 eq.), TBTU (1.0 eq.), DIPEA (3.0 eq.), and DMF (5.0 mL mmol^{-1}). The reaction mixture was stirred for 0.5 h followed by adding amidoxime **8** (1.0 eq.). The reaction mixture was stirred for 1 h at room temperature (RT), then refluxed in a pre-heated 120 °C oil-bath for 4 h and monitored by TLC. After cooling, the reaction mixture was diluted with water and extracted with ethyl acetate. The ethyl acetate layer was washed with 1 M HCl, saturated brine, and dried over anhydrous MgSO_4 . After filtration and concentration, the residue was purified on a silica gel column to give the product **10a–c**.

(4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)phenyl)methanol (**10a**). The characterization is found in our previous publication.³²

2-(2-Fluoroethoxy)-5-(3-(4-(hydroxymethyl)phenyl)-1,2,4-oxadiazol-5-yl)benzonitrile (**10b**). Compound **10b** was purified by flash chromatography, eluted with hexane/ethyl acetate (1/1, v/v) to afford white solid product. Yield: 30%, MP: 150–151 °C. ^1H NMR (400 MHz, $\text{acetone}-d_6$) δ 8.52–8.34 (m, 2H), 8.14–7.95 (m, 2H), 7.64–7.43 (m, 3H), 5.04–4.85 (m, 2H), 4.78 (d, J = 5.0 Hz, 2H), 4.73–4.60 (m, 2H), 4.49 (br, 1H). ^{13}C NMR (101 MHz, $\text{acetone}-d_6$) δ 173.8, 168.6, 163.1, 146.3, 134.2, 133.5, 127.1, 126.8, 125.1, 117.4, 114.6, 113.7, 102.9, 81.5 (d, $J_{\text{C-F}}$ = 170 Hz), 69.1 (d, $J_{\text{C-F}}$ = 18.1 Hz), 63.3. HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{F}_3\text{N}_3\text{O}_3$ [$\text{M} + \text{H}^+$] 340.1092, found 340.1087.

(4-(5-(4-(2-Fluoroethoxy)phenyl)-1,2,4-oxadiazol-3-yl)-phenyl)-methanol (**10c**). Compound **10c** was purified by flash chromatography, eluted with hexane/ethyl acetate (1/1, v/v) to afford white solid product. Yield: 45%, MP: 140–142 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.21–8.12 (m, 2H), 8.05 (d, J = 8.2 Hz, 2H), 7.54 (d, J = 8.0 Hz, 2H), 7.24 (d, J = 8.9 Hz, 2H), 5.38 (s, 1H), 4.94–4.74 (m, 2H), 4.61 (s, 2H), 4.53–4.34 (m, 2H). ^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ 175.5, 168.5, 162.4, 146.8, 130.4, 127.4, 127.3, 125.0, 116.5, 115.9, 83.2, 81.6, 68.0, 67.8, 62.9. HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{15}\text{FN}_2\text{O}_3$ [$\text{M} + \text{H}^+$] 315.1139, found 315.1139.

4-(((2,3-Dihydroxypropyl)(methyl)amino)methyl)benzonitrile (**11a**). To a round-bottom flask equipped with a stir bar was added 4-(bromomethyl)benzonitrile (3.9 g, 20.0 mmol), 3-(methylamino)propane-1,2-diol (2.5 g, 23.0 mmol), K_2CO_3 (6.9 g, 50.0 mmol), and acetonitrile (50 mL). The mixture was

stirred overnight at RT. Then, the mixture was concentrated under reduced pressure, the residue was purified by flash chromatography, eluted with hexane/ethyl acetate (1/1, v/v) to afford oil product. (3.0 g, 68%) ^1H NMR (400 MHz, CDCl_3) δ 7.64–7.56 (m, 2H), 7.44–7.36 (m, 2H), 3.88–3.79 (m, 1H), 3.77–3.66 (m, 2H), 3.60–3.44 (m, 2H), 2.70–2.60 (m, 1H), 2.43–2.34 (m, 1H), 2.25 (s, 3H).

4-(((2-Hydroxyethyl)(methyl)amino)methyl)benzonitrile (11b). To a round-bottom flask equipped with a stir bar was added 4-(bromomethyl)benzonitrile (3.0 g, 15.3 mmol), 2-(methylamino)ethan-1-ol (1.0 g, 13.3 mmol), and acetonitrile (20 mL). After cooling to 0 °C, triethylamine (2.8 g, 27.7 mmol) was added to the mixture dropwise. The mixture was warmed to RT and stirred overnight. Then, the mixture was concentrated in vacuum, and the crude residue was purified by flash chromatography, eluted with hexane/ethyl acetate (3/2, v/v) to afford the semi-solid product. (2.0 g, 79%) ^1H NMR (400 MHz, CDCl_3) δ 7.63–7.58 (m, 2H), 7.42 (d, J = 7.9 Hz, 2H), 3.68–3.59 (m, 5H), 2.63–2.57 (m, 2H), 2.23 (s, 3H).

***Tert*-Butyl (4-cyanobenzyl)(2-hydroxyethyl)carbamate (11c).** To a round-bottom flask equipped with a stir bar was added 4-formylbenzonitrile (6.0 g, 45.8 mmol), ethanolamine (2.8 g, 45.8 mmol), and methanol (80 mL). The mixture was stirred in a pre-heated 60 °C oil-bath for 6 h, then cooled to 0 °C followed by adding sodium borohydride (8.7 g, 229 mmol) portionwise. The reaction was warmed to room temperature (RT) slowly and stirred for an additional 5 h. Then, the reaction mixture was concentrated under reduced pressure, and the residue was dissolved in ethyl acetate (200 mL), washed with water (100 mL) and saturated brine (100 mL), and dried over anhydrous MgSO_4 . After filtering and concentration, the residual solid was re-dissolved in dichloromethane (100 mL). To the above solution was added di-*tert*-butyl dicarbonate (6.7 g, 30.6 mmol) and 1 N NaOH (17.0 mL). The mixture was stirred at RT overnight, then the mixture was washed with water and saturated brine, dried over anhydrous MgSO_4 . After filtering and concentration, the crude semi-solid product **11c** was obtained. (8.6 g, 68%) ^1H NMR (400 MHz, CDCl_3) δ 7.64 (d, J = 7.8 Hz, 2H), 7.38 (d, J = 8.0 Hz, 2H), 4.58 (s, 2H), 3.74 (s, 2H), 3.52–3.29 (m, 3H), 1.41 (s, 9H).

General procedure for the synthesis of 12a–c

To a round-bottom flask equipped with a stir bar was added **11a–c** (1.0 eq.), hydroxylamine hydrochloride (2.0 eq.), NaHCO_3 (4.0 eq.), and methanol (5.0 mL mmol^{-1}). The reaction was refluxed and stirred in a pre-heated 75 °C oil-bath for 6 h. The reaction mixture was cooled to room temperature, and the precipitate was filtered off and washed with methanol. The filtrate was concentrated under reduced pressure to give the product **12a–c** and used directly without further purification.

(*E*)-4-(((2,3-Dihydroxypropyl)(methyl)amino)methyl)-*N'*-hydroxybenzimidamide (12a). Yield: 51%, MP: 160–164 °C. ^1H NMR (400 MHz, $\text{MeOD-}d_4$) δ 7.60 (d, J = 7.8 Hz, 2H), 7.37 (d, J = 7.9 Hz, 2H), 3.85–3.78 (m, 1H), 3.66–3.58 (m, 3H), 3.57–3.43 (m, 3H), 2.52–2.45 (m, 2H), 2.25 (s, 3H).

(*E*)-*N'*-Hydroxy-4-(((2-hydroxyethyl)(methyl)amino)methyl)-benzimidamide (12b). Yield: 60%, MP: 162–165 °C. ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 9.57 (s, 1H), 7.61 (d, J = 7.1 Hz, 2H), 7.30 (d, J = 7.4 Hz, 2H), 5.76 (s, 2H), 4.39 (s, 1H), 3.50 (s, 4H), 2.42 (t, J = 6.2 Hz, 2H), 2.15 (s, 3H).

***Tert*-Butyl (*E*)-4-((*N'*-hydroxycarbamimidoyl)benzyl)(2-hydroxyethyl)carbamate (12c).** Yield: 71%, MP: 138–141 °C. ^1H NMR (400 MHz, CDCl_3) δ 7.66–7.53 (m, 2H), 7.35–7.26 (m, 2H), 5.30 (s, 2H), 4.86 (s, 2H), 4.52 (s, 2H), 3.71 (s, 2H), 3.47–3.31 (m, 2H), 1.46 (s, 9H).

(\pm)-3-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)(methyl)amino)propane-1,2-diol (13a). Compound **13a** was synthesized according to the general procedure as described for compound **10a–c**. Yield: 40%, white solid, MP: 101–103 °C. ^1H NMR (400 MHz, $\text{MeOD-}d_4$) δ 8.29 (d, J = 7.1 Hz, 2H), 8.01 (d, J = 8.0 Hz, 2H), 7.47 (d, J = 8.1 Hz, 2H), 7.35 (d, J = 9.4 Hz, 1H), 4.83–4.69 (m, 3H), 4.48–4.37 (m, 3H), 3.87–3.80 (m, 1H), 3.63 (s, 2H), 3.57–3.46 (m, 2H), 2.53–2.49 (m, 2H), 2.28 (s, 3H). ^{13}C NMR (101 MHz, $\text{MeOD-}d_4$) δ 174.3, 168.5, 159.8, 142.1, 133.3, 129.5, 126.9, 126.6 (q, $J_{\text{C-F}}$ = 5.4 Hz), 125.4, 122.9 (q, $J_{\text{C-F}}$ = 274 Hz), 119.3 (q, $J_{\text{C-F}}$ = 31 Hz), 116.4, 113.8, 81.2 (d, $J_{\text{C-F}}$ = 171 Hz), 68.9, 68.6 (q, $J_{\text{C-F}}$ = 20 Hz), 64.9, 62.0, 60.1, 41.7. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{23}\text{F}_4\text{N}_3\text{O}_4$ [$\text{M} + \text{H}^+$] 470.1697, found 470.1694.

2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)(methyl)amino)ethan-1-ol (13b). Compound **13b** was prepared according to the general procedure as described for compound **10a–c**. Yield: 34%, white solid, MP: 108–111 °C. ^1H NMR (400 MHz, CDCl_3) δ 8.45 (s, 1H), 8.34 (d, J = 8.7 Hz, 1H), 8.11 (d, J = 8.1 Hz, 2H), 7.45 (d, J = 8.1 Hz, 2H), 7.17 (d, J = 8.8 Hz, 1H), 4.90–4.74 (m, 2H), 4.47–4.37 (m, 2H), 3.68–3.62 (m, 4H), 2.64 (t, J = 5.3 Hz, 2H), 2.26 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ 174.2, 168.8, 159.9, 142.1, 133.3, 129.4, 127.8 (q, $J_{\text{C-F}}$ = 5.25 Hz), 127.6, 125.6, 122.7 (q, $J_{\text{C-F}}$ = 274 Hz), 120.2 (q, $J_{\text{C-F}}$ = 32 Hz), 117.1, 113.4, 81.2 (d, $J_{\text{C-F}}$ = 172 Hz), 68.4 (d, $J_{\text{C-F}}$ = 21.2 Hz), 62.0, 58.5, 58.4, 41.6. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{21}\text{F}_4\text{N}_3\text{O}_3$ [$\text{M} + \text{H}^+$] 440.1592, found 440.1588.

***Tert*-Butyl (4-(5-(4-(2-fluoroethoxy)-3-(trifluoromethyl)-phenyl)-1,2,4-oxadiazol-3-yl)benzyl)(2-hydroxyethyl)-carbamate (13c).** Compound **13c** was prepared according to the general procedure as described for compound **10a–c**. Yield: 42%, white solid, MP: 120–122 °C. ^1H NMR (400 MHz, CDCl_3) δ 8.47 (s, 1H), 8.35 (d, J = 8.8 Hz, 1H), 8.13 (d, J = 8.0 Hz, 2H), 7.38 (d, J = 7.7 Hz, 2H), 7.18 (d, J = 8.8 Hz, 1H), 4.92–4.75 (m, 2H), 4.56 (s, 2H), 4.50–4.36 (m, 2H), 3.75 (s, 2H), 3.46 (s, 2H), 2.90 (br, 1H), 1.47 (s, 9H).

2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)amino)ethan-1-ol (13d). To a solution of **13c** (250 mg, 0.48 mmol) in methanol (5.0 mL) was added 4 M HCl in dioxane (5.0 mL). The reaction was stirred at RT for 6 h, then the solvent was removed under reduced pressure and the residue was dissolved in ethyl acetate. The solution was washed with saturated NaHCO_3 , water, and dried over MgSO_4 . After filtering and concentration, the final product was obtained. (193 mg, 95%) MP: 126–130 °C. ^1H NMR (400 MHz, $\text{MeOD-}d_4$) δ 8.35 (d, J = 4.9 Hz, 2H), 8.06 (d, J = 8.1 Hz, 2H),

7.50 (d, J = 8.1 Hz, 2H), 7.39 (d, J = 9.3 Hz, 1H), 4.84–4.68 (m, 2H), 4.51–4.39 (m, 2H), 3.85 (s, 2H), 3.71–3.64 (m, 2H), 2.73 (t, J = 5.6 Hz, 2H). ^{13}C NMR (101 MHz, MeOD- d_4) δ 174.4, 168.4, 159.8, 142.8, 133.4, 128.7, 127.1, 126.7 (q, $J_{\text{C-F}}$ = 5.1 Hz), 125.4, 124.3, 121.6, 116.4, 113.9, 81.2 (d, $J_{\text{C-F}}$ = 171 Hz), 68.7 (d, $J_{\text{C-F}}$ = 20 Hz), 60.1, 52.5, 50.3. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{19}\text{F}_4\text{N}_3\text{O}_3$ [$\text{M} + \text{H}^+$] 426.1435, found 426.1432.

General procedure for the synthesis of 15a–b

To a round-bottom flask equipped with a stir bar was added aldehyde **14**^{32,33} (1.0 eq.), amine (1.5 eq.), methanol (14 mL mmol^{−1}), and acetic acid (0.5 mL mmol^{−1}). The reaction mixture was stirred for 1 h at which time sodium cyanoborohydride (1.0 eq.) was added. The reaction mixture was stirred overnight and diluted with water. The precipitate was filtered off and washed with water to give off-white solid product **15a–b**.

2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)amino)propane-1,3-diol (15a). Yield: 50%, off-white solid, MP: 135–137 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 8.47–8.41 (m, 1H), 8.34 (s, 1H), 8.04 (d, J = 8.2 Hz, 2H), 7.57 (d, J = 8.0 Hz, 3H), 4.89–4.70 (m, 2H), 4.63–4.50 (m, 2H), 4.45 (s, 2H), 3.88 (s, 2H), 3.49–3.35 (m, 4H), 2.61–2.53 (m, 1H). ^{13}C NMR (101 MHz, DMSO- d_6) δ 174.0, 168.2, 159.5, 145.6, 134.1, 128.6, 127.0, 126.6 (q, $J_{\text{C-F}}$ = 5.1 Hz), 124.2, 122.4 (q, $J_{\text{C-F}}$ = 274 Hz), 117.9, 116.0, 115.1, 109.6, 81.7 (d, $J_{\text{C-F}}$ = 162 Hz), 68.8 (q, $J_{\text{C-F}}$ = 20 Hz), 61.1, 60.5, 50.5. HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{21}\text{F}_4\text{N}_3\text{O}_4$ [$\text{M} + \text{H}^+$] 456.1541, found 456.1537.

(1-(4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)azetidin-3-yl)methanol (15b). Yield: 23%, white solid, MP: 114–116 °C. ^1H NMR (400 MHz, DMSO- d_6) δ 8.45 (d, J = 8.8 Hz, 1H), 8.36–8.32 (m, 1H), 8.12 (d, J = 8.2 Hz, 2H), 7.65–7.55 (m, 3H), 5.49 (s, 1H), 4.88–4.82 (m, 1H), 4.78–4.71 (m, 1H), 4.64–4.57 (m, 1H), 4.57–4.49 (m, 1H), 3.41–3.27 (m, 8H), 2.34–2.21 (m, 1H).

3-(4-(Chloromethyl)phenyl)-5-(4-(2-fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazole (16). To a round-bottom flask was added cyanuric chloride (253 mg, 1.37 mmol) and DMF (2.5 mL). A solution of **10c** (500 mg, 1.31 mmol) in dichloromethane (3.5 mL) was added dropwise at RT, and the reaction was continued to stir at RT for overnight. Then, the reaction mixture was diluted with dichloromethane and water. The dichloromethane layer was then separated and concentrated in vacuum to afford crude product (400 mg, 73%). The crude product was used directly for the next step. ^1H NMR (400 MHz, CDCl_3) δ = 8.47 (d, J = 2.0 Hz, 1H), 8.35 (dd, J = 8.8 Hz, 2.0 Hz, 1H), 8.17 (d, J = 8.0 Hz, 2H), 7.54 (d, J = 8.4 Hz, 2H), 7.18 (d, J = 8.8 Hz, 1H), 4.93–4.76 (m, 2H), 4.65 (s, 2H), 4.48–4.36 (m, 2H).

General procedure for the synthesis of 17a–e and 17g

To a round-bottom flask equipped with a stir bar was added alcohols (10 eq.) and THF (2 mL mmol^{−1}). After cooling to 0 °C, sodium hydride (1.8 eq.) was added portionwise, and the reaction vessel was equipped with a reflux condenser and heated to 80 °C and stirred for 30 min. Chloride **16** (1.0 eq.) was added to the flask and the mixture was stirred overnight. The reaction was cooled to room temperature and quenched

with water. The crude was extracted with ethyl acetate, washed with saturated brine, and dried over MgSO_4 . After filtering and concentrated, the residue was purified by flash chromatography on silica gel column to afford the product.

2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)oxy)ethanol (17a). Compound **17a** was purified by flash chromatography, eluted with hexane/ethyl acetate (2/1, v/v) to afford off-white solid. Yield: 35%, MP: 109–110 °C. ^1H NMR (400 MHz, CDCl_3) δ 8.42 (s, 1H), 8.30 (dd, J = 8.6, 1.4 Hz, 1H), 8.11 (d, J = 8.1 Hz, 2H), 7.46 (d, J = 8.0 Hz, 2H), 7.13 (d, J = 8.7 Hz, 1H), 4.89–4.71 (m, 2H), 4.62 (s, 2H), 4.46–4.32 (m, 2H), 3.82–3.73 (m, 2H), 3.67–3.59 (m, 2H), 2.19 (br, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 174.2, 168.7, 159.6, 141.4, 133.3, 127.9, 127.7 (q, $J_{\text{C-F}}$ = 5.2 Hz), 127.6, 126.1, 122.7 (q, $J_{\text{C-F}}$ = 271 Hz), 120.2 (q, $J_{\text{C-F}}$ = 32 Hz), 117.0, 113.4, 81.3 (d, $J_{\text{C-F}}$ = 172 Hz), 72.7, 71.7, 68.4 (d, $J_{\text{C-F}}$ = 21 Hz), 61.9. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{18}\text{F}_4\text{N}_2\text{O}_4$ [$\text{M} + \text{H}^+$] 427.1245, found 427.1240.

2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)oxy)ethoxyethanol (17b). Compound **17b** was purified by flash chromatography, eluted with hexane/ethyl acetate (1/1, v/v) to afford an off-white semi-solid. Yield: 39%. ^1H NMR (400 MHz, CDCl_3) δ 8.43 (s, 1H), 8.32 (d, J = 8.5 Hz, 1H), 7.94 (d, J = 7.8 Hz, 1H), 7.82 (d, J = 10.3 Hz, 1H), 7.71–7.51 (m, 2H), 7.16 (d, J = 8.7 Hz, 1H), 4.91–4.85 (m, 1H), 4.79–4.74 (m, 1H), 4.70 (s, 2H), 4.47–4.42 (m, 1H), 4.41–4.35 (m, 1H), 3.79–3.69 (m, 6H), 3.67–3.60 (m, 2H), 2.41 (br, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 174.4, 168.0, 159.5, 141.8, 133.3, 128.0, 127.7 (q, $J_{\text{C-F}}$ = 5.4 Hz), 127.2, 126.5, 122.8 (q, $J_{\text{C-F}}$ = 274 Hz), 119.0 (q, $J_{\text{C-F}}$ = 32 Hz), 116.8, 113.4, 81.2 (d, $J_{\text{C-F}}$ = 173 Hz), 72.5, 70.3, 70.0, 68.4 (d, $J_{\text{C-F}}$ = 21 Hz), 66.4, 61.8. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{22}\text{F}_4\text{N}_2\text{O}_5$ [$\text{M} + \text{H}^+$] 471.1583, found 471.1589.

2-(2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)oxy)ethoxy)ethoxyethanol (17c). Compound **17c** was purified by flash chromatography, eluted with hexane/ethyl acetate (1/3, v/v) to afford an off-white semi-solid. Yield: 42%. ^1H NMR (400 MHz, CDCl_3) δ 8.45 (br, 1H), 8.33 (d, J = 8.7 Hz, 1H), 8.13 (d, J = 8.1 Hz, 2H), 7.49 (d, J = 8.1 Hz, 2H), 7.16 (d, J = 8.8 Hz, 1H), 4.90–4.75 (m, 2H), 4.64 (s, 2H), 4.49–4.36 (m, 2H), 3.74–3.61 (m, 12H), 2.63 (br, 1H). ^{13}C NMR (101 MHz, CDCl_3) δ 174.2, 168.8, 159.6, 141.7, 132.2, 127.9, 127.7 (q, $J_{\text{C-F}}$ = 5.2 Hz), 127.6, 125.9, 122.7 (q, $J_{\text{C-F}}$ = 271 Hz), 120.2 (q, $J_{\text{C-F}}$ = 32 Hz), 117.1, 113.4, 81.3 (d, $J_{\text{C-F}}$ = 171 Hz), 72.7, 72.5, 70.7, 70.6, 70.3, 69.7, 68.4 (d, $J_{\text{C-F}}$ = 21 Hz), 61.7. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{26}\text{F}_4\text{N}_2\text{O}_6$ [$\text{M} + \text{H}^+$] 515.1800, found 515.1806.

1-(4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)phenyl)-2,5,8,11-tetraoxatridecan-13-ol (17d). Compound **17d** was purified by flash chromatography, eluted with hexane/ethyl acetate (1/3, v/v) to afford an off-white semi-solid. Yield: 25%. ^1H NMR (400 MHz, CDCl_3) δ 8.33 (d, J = 5.7 Hz, 1H), 8.25–8.18 (m, 1H), 8.07–8.00 (m, 2H), 7.40 (d, J = 7.9 Hz, 2H), 7.06 (d, J = 8.7 Hz, 1H), 4.80–4.64 (m, 3H), 4.59–4.52 (m, 2H), 4.36–4.27 (m, 2H), 3.69–3.54 (m, 14H), 3.54–3.48 (m, 2H). ^{13}C NMR (101 MHz, CDCl_3) δ 173.9, 168.3,

159.3, 141.8, 132.2, 127.9, 127.7 ($q, J_{C-F} = 5.2$ Hz), 127.6, 125.9, 122.7 ($q, J_{C-F} = 271$ Hz), 120.2 ($q, J_{C-F} = 32$ Hz), 117.1, 113.4, 81.3 ($d, J = 172.5$ Hz), 72.7, 72.5, 70.6, 70.5, 70.4, 70.3, 70.2, 69.7, 68.4 ($d, J_{C-F} = 21$ Hz), 61.7. HRMS (ESI) calcd for $C_{26}H_{30}F_4N_2O_7 [M + H^+]$ 559.2062, found 559.2066.

Tert-Butyl (2-((4-(5-(4-(2-fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)oxy)ethyl)(2-hydroxy-ethyl) carbamate (17e). Compound 17e was synthesized according to the general procedure. The crude product was used directly for the next step reaction without any further purification.

2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)oxy)ethyl)amino)ethan-1-ol (17f). To a solution of 17e (284 mg, 0.50 mmol) in methanol (5.0 mL) was added 4 M HCl in dioxane (5.0 mL). The reaction was stirred at RT for 6 h, then the solvent was concentrated in vacuum and the residue was dissolved in ethyl acetate. The solution was washed with saturated $NaHCO_3$, water, and dried over $MgSO_4$. After filtering and concentration, the final white solid product was obtained. (199 mg, 85%) MP: 126–130 °C. 1H NMR (400 MHz, $CDCl_3$) δ 8.45 (s, 1H), 8.34 (d, $J = 8.7$ Hz, 1H), 8.14 (d, $J = 8.1$ Hz, 2H), 7.46 (d, $J = 8.1$ Hz, 2H), 7.17 (d, $J = 8.8$ Hz, 1H), 4.91–4.75 (m, 2H), 4.60 (s, 2H), 4.47–4.38 (m, 2H), 4.35–4.29 (m, 2H), 3.74–3.67 (m, 4H), 3.52 (t, $J = 5.0$ Hz, 2H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 174.3, 168.7, 158.6, 141.3, 133.3, 127.8, 127.7 ($q, J_{C-F} = 5.25$ Hz), 127.6, 126.1, 122.7 ($q, J_{C-F} = 275$ Hz), 120.3 ($q, J_{C-F} = 31$ Hz), 117.0, 113.4, 86.3 ($d, J_{C-F} = 173$ Hz), 72.6, 68.9, 68.4 ($d, J_{C-F} = 21.1$ Hz), 62.0, 46.0, 44.2. HRMS (ESI) calcd for $C_{22}H_{23}F_4N_3O_4 [M + H^+]$ 470.1697, found 470.1694.

2-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)oxy)ethyl)(methyl)amino)ethan-1-ol (17g). Compound 17g was purified by flash chromatography, eluted with ethyl acetate/methanol (50/1, v/v) to afford a pale yellow semi-solid. Yield: 26%. 1H NMR (400 MHz, $CDCl_3$) δ 8.40 (d, $J = 1.8$ Hz, 1H), 8.29 (dd, $J = 8.7, 2.0$ Hz, 1H), 8.07 (d, $J = 8.1$ Hz, 2H), 7.42 (d, $J = 8.1$ Hz, 2H), 7.11 (d, $J = 8.8$ Hz, 1H), 4.88–4.79 (m, 1H), 4.75–4.65 (m, 1H), 4.54 (s, 2H), 4.44–4.25 (m, 2H), 4.12 (s, 3H), 3.55 (s, 2H), 2.73–2.60 (m, 4H), 2.29 (s, 3H). ^{13}C NMR (101 MHz, $CDCl_3$) δ 174.8, 168.7, 160.2, 159.0, 141.3, 133.3, 128.0, 127.8 ($q, J_{C-F} = 5.25$ Hz), 127.4, 125.9, 122.8 ($q, J_{C-F} = 275$ Hz), 120.4 ($q, J_{C-F} = 31$ Hz), 116.8, 113.4, 72.8, 60.4, 59.9, 57.7, 43.0. HRMS (ESI) calcd for $C_{17}H_{21}F_4N_3O_4 [M + H^+]$ 408.1541, found 408.1536.

4-(2-Hydroxyethoxy)benzonitrile (18a). Compound 18a was synthesized according to the published procedure.⁴⁵

4-((2,3-Dihydroxypropoxy)methyl)benzonitrile (18b). Commercially available DL-1,2-isopropylidene glycerol (3.0 g, 22.5 mmol) was added to a suspension of NaH (0.6 g, 25.8 mmol) in THF (30 mL) at 0 °C under nitrogen. The mixture was stirred for 30 min during which time 4-cyanobenzyl bromide (4.0 g, 20.4 mmol) was added dropwise over 5 min followed by warming of the reaction mixture to RT. After 2 h, the reaction mixture was partitioned between ammonium chloride solution and ethyl acetate. The aqueous phase was extracted with ethyl acetate and the combined organics dried over anhydrous $MgSO_4$ and concentrated in vacuum. The

crude residue was purified by flash chromatography on silica gel, eluted with hexane/ethyl acetate (1/1, v/v) to afford an off-white solid product. (3.8 g, 90%) MP: 80–83 °C. 1H NMR (400 MHz, $CDCl_3$) δ 7.63 (d, $J = 8.1$ Hz, 2H), 7.43 (d, $J = 7.9$ Hz, 2H), 4.61 (s, 2H), 3.97–3.89 (m, 1H), 3.76–3.69 (m, 1H), 3.66–3.56 (m, 3H), 2.75 (s, 1H), 2.29 (s, 1H).

(E)-N'-Hydroxy-4-(2-hydroxyethoxy)benzimidamide (19a). Compound 19a was synthesized according to the general procedure as described for compound 12a–c. The crude product was used directly for the next step reaction without further purification. 1H NMR (400 MHz, $DMSO-d_6$) δ 9.41 (s, 1H), 7.56 (d, $J = 8.7$ Hz, 2H), 6.89 (d, $J = 8.7$ Hz, 2H), 5.68 (s, 2H), 4.83 (s, 1H), 3.97 (t, $J = 4.7$ Hz, 2H), 3.68 (d, $J = 4.7$ Hz, 2H).

(E)-4-((2,3-Dihydroxypropoxy)methyl)-N'-hydroxybenzimidamide (19b). Compound 19b was synthesized according to the general procedure as described for compound 12a–c. Yield: 45%, MP: 142–144 °C. 1H NMR (400 MHz, $MeOD-d_4$) δ 7.61 (d, $J = 8.0$ Hz, 2H), 7.39 (d, $J = 7.9$ Hz, 2H), 4.57 (s, 2H), 3.83–3.75 (m, 1H), 3.64–3.45 (m, 4H).

2-(4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)phenoxy)ethan-1-ol (20a). Compound 20a was synthesized according to the general procedure as described for compound 10a–c. Yield: 37%, a white solid. MP: 148–149 °C. 1H NMR (400 MHz, $acetone-d_6$) δ 8.45–8.39 (m, 2H), 8.08 (d, $J = 8.9$ Hz, 2H), 7.56 (d, $J = 8.7$ Hz, 1H), 7.13 (d, $J = 8.9$ Hz, 2H), 4.95–4.80 (m, 2H), 4.68–4.56 (m, 2H), 4.18 (t, $J = 4.9$ Hz, 2H), 4.05 (br, 1H), 3.92 (d, $J = 4.3$ Hz, 2H). ^{13}C NMR (101 MHz, $acetone-d_6$) δ 174.1, 168.4, 161.8, 159.9, 133.7, 128.9, 126.8 ($q, J_{C-F} = 5.1$ Hz), 123.2 ($q, J_{C-F} = 273$ Hz), 119.3, 119.0, 116.8, 114.9, 114.4, 81.6 ($d, J_{C-F} = 170$ Hz), 69.9, 68.9 (d, $J_{C-F} = 20$ Hz), 60.3. HRMS (ESI) calcd for $C_{19}H_{16}F_4N_2O_4 [M + H^+]$ 413.1119, found 413.1113.

(±)-3-((4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)benzyl)oxy)propane-1,2-diol (20b). Compound 20b was synthesized according to the general procedure as described for compound 10a–c. Yield: 38%, white solid, MP: 102–105 °C. 1H NMR (400 MHz, $MeOD-d_4$) δ 8.34–8.28 (m, 2H), 8.04 (d, $J = 8.1$ Hz, 2H), 7.50 (d, $J = 8.0$ Hz, 2H), 7.35 (d, $J = 8.5$ Hz, 1H), 4.85–4.69 (m, 2H), 4.61 (s, 2H), 4.48–4.38 (m, 2H), 3.88–3.81 (m, 1H), 3.67–3.51 (m, 4H). ^{13}C NMR (101 MHz, $MeOD-d_4$) δ 174.3, 168.4, 159.8, 142.1, 133.3, 127.6, 126.9, 126.6 ($q, J_{C-F} = 5.35$ Hz), 125.6, 122.9 ($q, J_{C-F} = 273$ Hz), 119.3 ($q, J_{C-F} = 31$ Hz), 116.3, 113.8, 81.2 (d, $J_{C-F} = 170$ Hz), 72.3, 71.6, 70.9, 68.6 ($q, J_{C-F} = 20.1$ Hz), 63.1. HRMS (ESI) calcd for $C_{21}H_{20}F_4N_2O_5 [M + H^+]$ 457.1381, found 457.1385.

(E)-N'-Hydroxy-4-vinylbenzimidamide (22). Compound 22 was synthesized according to the general procedure as described for compound 12a–c. Yield: 95%. The detailed procedure was described in the published literature.⁴⁶

5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-3-(4-vinylphenyl)-1,2,4-oxadiazole (23). Compound 23 was synthesized according to the general procedure as described for compound 10a–c. The crude product was used directly for the next step reaction without further purification.

(±)-1-(4-(5-(4-(2-Fluoroethoxy)-3-(trifluoromethyl)phenyl)-1,2,4-oxadiazol-3-yl)phenyl)ethane-1,2-diol (24). To a 50 mL of

round bottle flask was added **18** (327 mg, 0.86 mmol), 4-methylmorpholine *N*-oxide (122 mg, 1.03 mmol) and THF/H₂O (3/1, 12.0 mL). The mixture was stirred for 5 min before adding OsO₄ (2.5 wt% in *tert*-butanol, 545 μ L). The reaction was stirred continuously at RT overnight. Then, the reaction mixture was diluted with ethyl acetate and water. The ethyl acetate layer was separated and concentrated in vacuum to afford crude product. Crude product was purified by flash chromatography, eluted with hexane/ethyl acetate (1/2, v/v) to furnish the final product as a white semi-solid. (120 mg, 34%) ¹H NMR (400 MHz, MeOD-*d*₄) δ 8.39–8.31 (m, 2H), 8.08 (d, *J* = 8.2 Hz, 2H), 7.55 (d, *J* = 8.2 Hz, 2H), 7.38 (d, *J* = 8.8 Hz, 1H), 4.85–4.70 (m, 3H), 4.50–4.39 (m, 2H), 3.72–3.60 (m, 2H). ¹³C NMR (101 MHz, MeOD-*d*₄) δ 174.4, 168.5, 159.9, 145.8, 133.3, 126.9, 126.7 (*q*, *J*_{C-F} = 5.2 Hz), 126.6, 125.6, 122.9 (*q*, *J*_{C-F} = 273 Hz), 119.4 (*q*, *J*_{C-F} = 32 Hz), 116.4, 113.8, 81.2 (d, *J*_{C-F} = 171 Hz), 74.1, 68.6 (d, *J*_{C-F} = 20.2 Hz), 67.1. HRMS (ESI) calcd for C₁₉H₁₆F₄N₂O₄ [M + H⁺] 413.1119, found 413.1115.

(*E*)-*N'*-Hydroxy-4-((methoxymethoxy)methyl)benzimidamide (25a). Compound **25a** was synthesized according to the general procedure as described for compound **12a-c**. Yield: 79%, a pale white solid. MP: 158–159 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.64 (s, 1H), 7.70–7.62 (m, 2H), 7.37–7.26 (m, 2H), 5.80 (s, 3H), 4.69–4.61 (m, 2H), 4.53 (s, 2H), 3.30 (s, 2H).

(*E*)-*N'*-Hydroxy-4-((2-(methoxymethoxy)ethoxy)methyl)benzimidamide (25b). Compound **25b** was synthesized according to the general procedure as described for compound **12a-c**. Yield: 60%, a white solid. MP: 150–151 °C. ¹H NMR (400 MHz, MeOD-*d*₄) δ 7.62 (d, *J* = 8.1 Hz, 2H), 7.38 (d, *J* = 8.1 Hz, 2H), 4.63 (s, 2H), 4.57 (s, 2H), 3.74–3.63 (m, 4H), 3.34 (s, 3H).

(*E*)-*N'*-Hydroxy-4-(2,4,7,10-tetraoxaundecan-11-yl)benzimidamide (25c). Compound **25c** was synthesized according to the general procedure as described for compound **12a-c**. Yield: 50%, a white solid. MP: 140–142 °C. ¹H NMR (400 MHz, MeOD-*d*₄) δ 7.65 (d, *J* = 8.1 Hz, 2H), 7.41 (d, *J* = 8.1 Hz, 2H), 4.66 (s, 2H), 4.60 (s, 2H), 3.73–3.65 (m, 8H), 3.37 (s, 3H).

(*E*)-4-(2,4,7,9-Tetraoxadecan-5-yl)-*N'*-hydroxybenzimidamide (25d). Compound **25d** was synthesized according to the general procedure as described for compound **12a-c**. Yield: 62%, a yellow solid. MP: 170–172 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.40 (s, 1H), 8.19 (d, *J* = 8.8 Hz, 1H), 8.11 (d, *J* = 8.1 Hz, 2H), 7.50 (d, *J* = 8.1 Hz, 2H), 7.04 (d, *J* = 8.6 Hz, 1H), 4.91–4.87 (m, 1H), 4.75–4.71 (m, 1H), 4.69–4.64 (m, 3H), 3.84–3.72 (m, 2H), 3.40 (s, 3H), 3.31 (s, 3H).

General procedure for the synthesis of 26a–d

To a round-bottom flask equipped with a stir bar was added 4-hydroxy-3-(trifluoromethyl)benzoic acid (1.0 eq.), HOBt (0.2 eq.), TBTU (1.0 eq.), DIPEA (3.0 eq.), and DMF (5.0 mL mmol^{−1}). The reaction mixture was stirred for 0.5 h followed by adding methoxymethyl (MOM) protected amidoxime **25a-d** (1.0 eq.). The reaction mixture was stirred for 1 h at room temperature, then refluxed in a pre-heated 120 °C oil-bath for 4 h and monitored by TLC. After cooling, the reaction mixture was diluted with water and extracted with ethyl acetate. The ethyl acetate layer was washed with 1 M HCl, saturated brine,

and dried over anhydrous MgSO₄. After filtration and concentration under reduced pressure, the crude residue was purified on a silica gel column to give **26a-d**.

4-(3-(4-((Methoxymethoxy)methyl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenol (26a). Compound **26a** was purified by flash chromatography, eluted with hexane/ethyl acetate (2/1, v/v) to afford a white solid. Yield: 50%, white solid, MP: 130–135 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 11.83 (br, 1H), 8.29–8.24 (m, 2H), 8.07 (d, *J* = 8.0 Hz, 2H), 7.55 (d, *J* = 8.0 Hz, 2H), 7.27 (d, *J* = 8.8 Hz, 1H), 4.70 (s, 2H), 4.63 (s, 2H), 3.33 (s, 3H).

4-(3-(4-((2-(Methoxymethoxy)ethoxy)methyl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenol (26b). Compound **26b** was purified by flash chromatography, eluted with hexane/ethyl acetate (3/2, v/v) to afford a white solid. Yield: 45%, white solid, MP: 120–122 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.34 (s, 1H), 8.04–7.93 (m, 3H), 7.53–7.45 (m, 1H), 7.41 (d, *J* = 8.1 Hz, 2H), 6.99 (d, *J* = 8.6 Hz, 1H), 4.73 (s, 2H), 4.62 (s, 2H), 3.86–3.76 (m, 4H), 3.41 (s, 3H).

4-(3-(4-(2,4,7,10-Tetraoxaundecan-11-yl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenol (26c). Compound **26c** was purified by flash chromatography, eluted with hexane/ethyl acetate (3/2, v/v) to afford a white solid. Yield: 60%, MP: 103–105 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.26 (d, *J* = 6.1 Hz, 1H), 7.90 (d, *J* = 7.9 Hz, 2H), 7.80 (d, *J* = 8.6 Hz, 1H), 7.30 (d, *J* = 7.9 Hz, 2H), 6.83 (d, *J* = 8.6 Hz, 1H), 4.61 (s, 2H), 4.54 (s, 2H), 3.85–3.70 (m, 8H), 3.34 (s, 3H), 2.86 (s, 2H).

4-(3-(4-(2,4,7,9-Tetraoxadecan-5-yl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenol (26d). Compound **26d** was purified by flash chromatography, eluted with hexane/ethyl acetate (1/1, v/v) to afford a white semi-solid. Yield: 50%. ¹H NMR (400 MHz, CDCl₃) δ 8.38 (s, 1H), 8.14 (d, *J* = 8.4 Hz, 1H), 8.08 (d, *J* = 8.1 Hz, 2H), 7.48 (d, *J* = 8.1 Hz, 2H), 7.02 (d, *J* = 8.6 Hz, 1H), 4.89–4.84 (m, 1H), 4.73–4.69 (m, 1H), 4.68–4.61 (m, 3H), 3.84–3.69 (m, 3H), 3.38 (s, 3H), 3.29 (s, 3H).

General procedure for the synthesis of 27a–d

To a round-bottom flask equipped with a stir bar was added **26a-d** (1.0 eq.), ethylene ditosylate (2.0 eq.), K₂CO₃ (3.0 eq.), and acetonitrile (10 mL mmol^{−1}). The reaction mixture was stirred at RT for 1 h, then refluxed in a pre-heated 90 °C oil-bath overnight and monitored by TLC. The reaction was diluted with ethyl acetate and water, the ethyl acetate layer was washed with saturated brine and dried over anhydrous MgSO₄. After filtering and concentrated in vacuum, the crude residue was purified on a silica gel column to give **27a-d**.

2-(4-(3-(4-((Methoxymethoxy)methyl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenoxy)ethyl 4-methyl-benzenesulfonate (27a). Compound **27a** was purified by flash chromatography, eluted with hexane/ethyl acetate (3/1, v/v) to afford a white solid. Yield: 30%, MP: 125–127 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.44 (s, 1H), 8.32 (d, *J* = 8.8 Hz, 1H), 8.15 (d, *J* = 8.0 Hz, 2H), 7.81 (d, *J* = 8.0 Hz, 2H), 7.51 (d, *J* = 8.0 Hz, 2H), 7.35 (d, *J* = 8.0 Hz, 2H), 7.09 (d, *J* = 8.8 Hz, 1H), 4.75 (s, 2H), 4.68 (s, 2H), 4.45–4.35 (m, 4H), 3.44 (s, 3H), 2.45 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 174.3, 169.0, 159.2, 145.3, 141.6,

133.5, 132.6, 130.1, 128.2, 128.1, 127.8 (q, J_{C-F} = 5.5 Hz), 127.8, 126.1, 122.8 (q, J_{C-F} = 271 Hz), 120.3 (q, J_{C-F} = 32 Hz), 117.5, 113.5, 96.1, 68.9, 67.3, 66.8, 55.6, 21.8. HRMS (ESI) calcd for $C_{27}H_{25}F_3N_2O_7S$ [$M + H^+$] 579.1407. Found [$M + H^+$] 579.1415.

2-(4-(3-(4-((2-(Methoxymethoxy)ethoxy)methyl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenoxy)ethyl 4-methyl-benzenesulfonate (27b). Compound 27b was purified by flash chromatography, eluted with hexane/ethyl acetate (3/1, v/v) to afford a white solid. Yield: 57%, MP: 115–117 °C. 1H NMR (400 MHz, $CDCl_3$) δ 8.37 (d, J = 1.4 Hz, 1H), 8.28–8.23 (m, 1H), 8.10 (d, J = 8.2 Hz, 2H), 7.78 (d, J = 8.2 Hz, 2H), 7.48 (d, J = 8.1 Hz, 2H), 7.32 (d, J = 8.1 Hz, 2H), 7.05 (d, J = 8.8 Hz, 1H), 4.67 (s, 2H), 4.64 (s, 2H), 4.40–4.33 (m, 4H), 3.78–3.67 (m, 4H), 3.37 (s, 3H), 2.41 (s, 3H). ^{13}C NMR (101 MHz, $CDCl_3$): δ 174.1, 168.7, 159.0, 145.2, 141.8, 133.3, 132.3, 129.9, 127.9, 127.8, 127.7, 127.5 (q, J_{C-F} = 5.5 Hz), 125.8, 122.6 (q, J_{C-F} = 271 Hz), 120.0 (q, J_{C-F} = 32 Hz), 117.1, 113.3, 96.5, 72.7, 69.7, 67.3, 66.8, 66.5, 55.2, 21.6. HRMS (ESI) calcd for $C_{29}H_{29}F_3N_2O_8S$ [$M + H^+$] 623.1669, found 623.1660.

2-(4-(3-(4-(2,4,7,10-Tetraoxaundecan-11-yl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenoxy)ethyl 4-methyl-benzenesulfonate (27c). Compound 27c was purified by flash chromatography, eluted with hexane/ethyl acetate (3/1, v/v) to afford a white semi-solid. Yield: 38%. 1H NMR (400 MHz, $CDCl_3$) δ 8.41 (s, 1H), 8.31 (d, J = 8.7 Hz, 1H), 8.13 (d, J = 8.1 Hz, 2H), 7.80 (d, J = 8.2 Hz, 2H), 7.49 (d, J = 8.1 Hz, 2H), 7.34 (d, J = 8.1 Hz, 2H), 7.08 (d, J = 8.8 Hz, 1H), 4.67 (s, 2H), 4.65 (s, 2H), 4.44–4.35 (m, 4H), 3.77–3.67 (m, 8H), 3.37 (s, 3H), 2.44 (s, 3H). ^{13}C NMR (101 MHz, $CDCl_3$): δ 174.1, 168.8, 159.0, 145.2, 141.9, 133.3, 132.3, 129.9, 127.9, 127.9, 127.6 (q, J_{C-F} = 5.5 Hz), 127.5, 125.8, 122.6 (q, J_{C-F} = 271 Hz), 120.1 (q, J_{C-F} = 32 Hz), 117.2, 113.3, 96.5, 72.7, 70.6, 69.8, 67.2, 66.8, 66.5, 55.2, 21.6. HRMS (ESI) calcd for $C_{31}H_{33}F_3N_2O_9S$ [$M + H^+$] 667.1932, found 667.1925.

(\pm)-2-(4-(3-(4-(2,4,7,9-Tetraoxadecan-5-yl)phenyl)-1,2,4-oxadiazol-5-yl)-2-(trifluoromethyl)phenoxy)ethyl 4-methyl-benzenesulfonate (27d). Compound 27d was purified by flash chromatography, eluted with hexane/ethyl acetate (2/1, v/v) to afford a white semi-solid. Yield: 44%. 1H NMR (400 MHz, $CDCl_3$) δ 8.42 (s, 1H), 8.32 (d, J = 8.7 Hz, 1H), 8.15 (d, J = 8.1 Hz, 2H), 7.81 (d, J = 7.9 Hz, 2H), 7.52 (d, J = 8.2 Hz, 2H), 7.34 (d, J = 8.0 Hz, 2H), 7.09 (d, J = 8.7 Hz, 1H), 4.91–4.86 (m, 1H), 4.75–4.59 (m, 5H), 4.46–4.36 (m, 4H), 3.84–3.70 (m, 2H), 3.38 (s, 3H), 3.29 (s, 3H), 2.44 (s, 3H). ^{13}C NMR (101 MHz, $DMSO-d_6$) δ 174.6, 168.5, 159.5, 145.5, 143.7, 134.5, 132.4, 130.5, 128.3, 128.0, 127.5, 127.1, 127.0 (q, J_{C-F} = 5.5 Hz), 123.1 (q, J_{C-F} = 271 Hz), 125.8, 118.8, 116.4, 115.4, 96.2, 94.9, 76.9, 71.1, 68.8, 67.3, 55.4, 55.1, 21.5. HRMS (ESI) calcd for $C_{30}H_{31}F_3N_2O_9S$ [$M + H^+$] 653.1775, found 653.1770.

General procedure for the radiosynthesis of [^{18}F]10a, [^{18}F]17a, [^{18}F]17b, and (\pm)-[^{18}F]24

[^{18}F]KF (~7.4 GBq) aqueous was added to a vial containing Kryptofix 222 (~10 mg), and dried by azeotropic evaporation with acetonitrile (3 \times 1 mL) under N_2 flow at 110 °C. To the reaction vial was added precursor 27a–d (~2 mg) as a solution

in acetonitrile (300 μ L). The reaction was placed in a 110 °C oil-bath for 15 min. The reaction was removed from the oil-bath, at which time 6 M HCl (150 μ L) was added. The reaction mixture was heated in a 110 °C oil-bath for another 5 min. Then, the reaction was removed from the oil-bath and quenched with 6 M NaOH (150 μ L) and diluted with 2.4 mL of the HPLC mobile phase. The mixture was passed through a Sep-Pak® Alumina N Cartridge (Part No. WAT020510) and injected onto the semi-preparation HPLC column (Agilent SB-C18 250 \times 9.6 mm, 5 μ m, UV = 254 nm, 4.0 mL min $^{-1}$). The HPLC fraction was collected into a water bottle with 60 mL of water and then trapped on a Sep-Pak® C-18 Cartridge (Part No. WAT020515). The activity was washed out with 0.6 mL of ethanol and 5.4 mL of 0.9% saline. After sterile filtration into a glass vial, the radiolabeled product was ready for quality control (QC) analysis and animal studies. An aliquot of sample was injected onto an analytical HPLC column (Agilent Eclipse XDB-C18 250 \times 4.6 mm, 5 μ m) to determine the concentration of tracer. Meanwhile, the tracer was authenticated by co-injecting with the non-radiolabeled standard sample solution.

Semi-preparation HPLC conditions. [^{18}F]10a, [^{18}F]17a, and [^{18}F]17b: Mobile phase, 50% acetonitrile in 0.1 M ammonium formate buffer, pH = 4.2; flow rate, 4.0 mL min $^{-1}$; UV = 254 nM; t_R = 18–21 min. The radiochemical yields were 38%, 40%, and 35%, respectively (decay corrected to EOS).

(\pm)-[^{18}F]24: Mobile phase, 40% acetonitrile in 0.1 M ammonium formate buffer, pH = 4.2; flow rate, 4.0 mL min $^{-1}$; UV = 254 nM; t_R = 25–27 min. The radiochemical yield was 38% (decay corrected to EOS).

QC HPLC conditions. [^{18}F]10a: Mobile phase, 75% acetonitrile in 0.1 M ammonium formate buffer, pH = 4.2; flow rate, 1.5 mL min $^{-1}$; UV = 254 nM; t_R = 5.5 min.

[^{18}F]17a: Mobile phase, 70% acetonitrile in 0.1 M ammonium formate buffer, pH = 4.2; flow rate, 1.0 mL min $^{-1}$; UV = 254 nM; t_R = 5.7 min.

[^{18}F]17b: Mobile phase, 70% acetonitrile in 0.1 M ammonium formate buffer, pH = 4.2; flow rate, 1.0 mL min $^{-1}$; UV = 254 nM; t_R = 5.6 min.

(\pm)-[^{18}F]24: Mobile phase, 63% acetonitrile in 0.1 M ammonium formate buffer, pH = 4.2; flow rate, 1.5 mL min $^{-1}$; UV = 254 nM; t_R = 4.6 min.

In vitro S1PR1–5 binding assay

The binding potencies of the newly synthesized compounds were determined by competition against the binding of [^{32}P] S1P to commercial cell membranes expressing recombinant human S1PRs (1, 2, 3, 4, and 5) according to methods reported previously.^{32,33,35}

MicroPET studies in cynomolgus macaque

Male macaques (9–10 kg) were studied with a microPET Focus 220 scanner (Concorde/CTI/Siemens Microsystems, Knoxville, TN). Animals were maintained in facilities with 12-hour dark and light cycles, given access to food, water, and libitum. Animals were also provided a variety of psychologically enriching tasks to prevent inappropriate deprivation. Animals were

scanned under anesthesia (induced with ketamine and glycopyrrolate and maintained with inhalation isoflurane). Core temperature was kept about 37 °C with a heated water blanket. The head was secured in a head holder with the brain in the center of the field of view. Subsequently, a 2 h dynamic emission scan was acquired after administration of ~0.35 GBq of radiotracer *via* the venous catheter.

PET scans were collected from 0–120 min with the following time frames: 3 × 1 min, 4 × 2 min, 3 × 3 min and 20 × 5 min. Emission data were corrected for dead time, scatter and attenuation and then reconstructed to a final resolution of 2.0 mm full-width half maximum in all 3 dimensions at the center of the field of view. PET and MRI images were co-registered using automated image registration program (AIR). For quantitative analyses, three-dimensional ROI (the global brain) was identified on the MRI and transformed to the reconstructed PET images to obtain time-activity curves. Activity measures were standardized to body weight and the dose of radioactivity injected to yield SUV.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the USA National Institutes of Health including the National Institute of Neurological Disorders and Stroke, and the National Institute on Aging [NS075527 and NS103988], the National Institute of Mental Health [MH092797], and National Institute of Biomedical Imaging and Bioengineering [EB025815].

This work was also partially supported by USA Department of Energy Training Grant titled “Training in Techniques and Translation: Novel Nuclear Medicine Imaging Agents for Oncology and Neurology” [DE-SC0008432]. American Parkinson Disease Association (APDA), the Greater St Louis Chapter of the APDA and the Barnes Jewish Hospital Foundation.

Notes and references

- 1 R. L. Proia and T. Hla, *J. Clin. Invest.*, 2015, **125**, 1379–1387.
- 2 M. Kono and R. L. Proia, *Exp. Cell Res.*, 2015, **333**, 178–182.
- 3 V. A. Blaho and T. Hla, *J. Lipid Res.*, 2014, **55**, 1596–1608.
- 4 M. J. Lee, J. R. Van Brocklyn, S. Thangada, C. H. Liu, A. R. Hand, R. Menzeleev, S. Spiegel and T. Hla, *Science*, 1998, **279**, 1552–1555.
- 5 T. Kohama, A. Olivera, L. Edsall, M. M. Nagiec, R. Dickson and S. Spiegel, *J. Biol. Chem.*, 1998, **273**, 23722–23728.
- 6 T. Hla and T. Maciag, *J. Biol. Chem.*, 1990, **265**, 9308–9313.
- 7 N. Ishizaka, H. Okazaki, K. Kurokawa, M. Kumada and Y. Takuwa, *Biochim. Biophys. Acta*, 1994, **1218**, 173–180.
- 8 C. Wang, J. Mao, S. Redfield, Y. Mo, J. M. Lage and X. Zhou, *Exp. Mol. Pathol.*, 2014, **97**, 259–265.
- 9 M. Maceyka and S. Spiegel, *Nature*, 2014, **510**, 58–67.
- 10 F. D. Lublin and S. C. Reingold, *Neurology*, 1996, **46**, 907–911.
- 11 G. K. Sheridan and K. K. Dev, *Glia*, 2012, **60**, 382–392.
- 12 V. E. Miron, S. K. Ludwin, P. J. Darlington, A. A. Jarjour, B. Soliven, T. E. Kennedy and J. P. Antel, *Am. J. Pathol.*, 2010, **176**, 2682–2694.
- 13 S. J. Jackson, G. Giovannoni and D. Baker, *J. Neuroinflammation*, 2011, **8**, 76.
- 14 V. Brinkmann, *Br. J. Pharmacol.*, 2009, **158**, 1173–1182.
- 15 J. W. Choi, S. E. Gardell, D. R. Herr, R. Rivera, C. W. Lee, K. Noguchi, S. T. Teo, Y. C. Yung, M. Lu, G. Kennedy and J. Chun, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 751–756.
- 16 S. Trkov, M. Stenovec, M. Kreft, M. Potokar, V. Parpura, B. Davletov and R. Zorec, *Glia*, 2012, **60**, 1406–1416.
- 17 H. Noda, H. Takeuchi, T. Mizuno and A. Suzumura, *J. Neuroimmunol.*, 2013, **256**, 13–18.
- 18 E. W. Radue, P. O'Connor, C. H. Polman, R. Hohlfeld, P. Calabresi, K. Selmaj, N. Mueller-Lenke, C. Agoropoulou, F. Holdbrook, A. de Vera, L. Zhang-Auberson, G. Francis, P. Burtin and L. Kappos, *Arch. Neurol.*, 2012, **69**, 1259–1269.
- 19 T. Karuppuachamy, E. h. Behrens, P. González-Cabrera, G. Sarkisyan, L. Gima, J. D. Boyer, G. Bamias, P. Jedlicka, M. Veny, D. Clark, R. Peach, F. Scott, H. Rosen and J. Rivera-Nieves, *Mucosal Immunol.*, 2016, **10**, 162.
- 20 M. van der Giet, M. Tolle and B. Kleuser, *Biol. Chem.*, 2008, **389**, 1381–1390.
- 21 J. E. Hughes, S. Srinivasan, K. R. Lynch, R. L. Proia, P. Ferdek and C. C. Hedrick, *Circ. Res.*, 2008, **102**, 950–958.
- 22 H. Lee, J. Deng, M. Kujawski, C. Yang, Y. Liu, A. Herrmann, M. Kortylewski, D. Horne, G. Somlo, S. Forman, R. Jove and H. Yu, *Nat. Med.*, 2010, **16**, 1421–1428.
- 23 J. Liang, M. Nagahashi, E. Y. Kim, K. B. Harikumar, A. Yamada, W. C. Huang, N. C. Hait, J. C. Allegood, M. M. Price, D. Avni, K. Takabe, T. Kordula, S. Milstien and S. Spiegel, *Cancer Cell*, 2013, **23**, 107–120.
- 24 C. Watson, J. S. Long, C. Orange, C. L. Tannahill, E. Mallon, L. M. McGlynn, S. Pyne, N. J. Pyne and J. Edwards, *Am. J. Pathol.*, 2010, **177**, 2205–2215.
- 25 N. J. Pyne and S. Pyne, *Nat. Rev. Cancer*, 2010, **10**, 489–503.
- 26 S. Ponnusamy, S. P. Selvam, S. Mehrotra, T. Kawamori, A. J. Snider, L. M. Obeid, Y. Shao, R. Sabbadini and B. Ogretmen, *EMBO Mol. Med.*, 2012, **4**, 761–775.
- 27 H. Liu, H. Jin, X. Yue, Z. Luo, C. Liu, A. J. Rosenberg and Z. Tu, *Mol. Imaging Biol.*, 2016, **18**, 724–732.
- 28 H. Jin, H. Yang, H. Liu, Y. Zhang, X. Zhang, A. J. Rosenberg, Y. Liu, S. E. Lapi and Z. Tu, *J. Nucl. Cardiol.*, 2017, **24**, 558–570.
- 29 H. Liu, H. Jin, X. Yue, J. Han, P. Baum, D. R. Abendschein and Z. Tu, *Mol. Imaging*, 2017, **16**, 1–7.
- 30 V. P. Prasad, S. Wagner, P. Keul, S. Hermann, B. Levkau, M. Schäfers and G. Haufe, *Bioorg. Med. Chem.*, 2014, **22**, 5168–5181.

- 31 R. S. Shaikh, S. S. Schilson, S. Wagner, S. Hermann, P. Keul, B. Levkau, M. Schäfers and G. Haufe, *J. Med. Chem.*, 2015, **58**, 3471–3484.
- 32 A. J. Rosenberg, H. Liu, H. Jin, X. Yue, S. Riley, S. J. Brown and Z. Tu, *J. Med. Chem.*, 2016, **59**, 6201–6220.
- 33 Z. Luo, A. J. Rosenberg, H. Liu, J. Han and Z. Tu, *Eur. J. Med. Chem.*, 2018, **150**, 796–808.
- 34 V. W. Pike, *Trends Pharmacol. Sci.*, 2009, **30**, 431–440.
- 35 A. J. Rosenberg, H. Liu and Z. Tu, *Appl. Radiat. Isot.*, 2015, **102**, 5–9.
- 36 J. M. Harris and R. B. Chess, *Nat. Rev. Drug Discovery*, 2003, **2**, 214.
- 37 C. S. Fishburn, *J. Pharm. Sci.*, 2008, **97**, 4167–4183.
- 38 T. Riley and J. Riggs-Sauthier, *Pharm. Technol.*, 2008, **32**, 88, 90–92, 94.
- 39 W. Li, P. Zhan, E. De Clercq, H. Lou and X. Liu, *Prog. Polym. Sci.*, 2013, **38**, 421–444.
- 40 G. Mattheolabakis, C. C. Wong, Y. Sun, C. A. Amella, R. Richards, P. P. Constantinides and B. Rigas, *J. Pharmacol. Exp. Ther.*, 2014, **351**, 61–66.
- 41 H. F. Kung, S. R. Choi, W. Qu, W. Zhang and D. Skovronsky, *J. Med. Chem.*, 2010, **53**, 933–941.
- 42 Z. Zha, S. R. Choi, K. Ploessl, B. P. Lieberman, W. Qu, F. Hefti, M. Mintun, D. Skovronsky and H. F. Kung, *J. Med. Chem.*, 2011, **54**, 8085–8098.
- 43 H. Pajouhesh and G. R. Lenz, *NeuroRx*, 2005, **2**, 541–553.
- 44 D. S. Miller, B. Bauer and A. M. S. Hartz, *Pharmacol. Rev.*, 2008, **60**, 196–209.
- 45 N. Kihara, M. Hashimoto and T. Takata, *Org. Lett.*, 2004, **6**, 1693–1696.
- 46 J. L. Gilmore, J. E. Sheppeck, S. H. Watterson, L. Haque, P. Mukhopadhyay, A. J. Tebben, M. A. Galella, D. R. Shen, M. Yarde, M. E. Cvijic, V. Borowski, K. Gillooly, T. Taylor, K. W. McIntyre, B. Warrack, P. C. Levesque, J. P. Li, G. Cornelius, C. D'Arienzo, A. Marino, P. Balimane, L. Salter-Cid, J. C. Barrish, W. J. Pitts, P. H. Carter, J. Xie and A. J. Dyckman, *J. Med. Chem.*, 2016, **59**, 6248–6264.