Communications



Reductive Aldol Reaction

P. Nuhant, C. Allais, W. R. Roush* _____ ∎∎∎∎−∎∎∎∎

Diisopinocampheylborane-Mediated Reductive Aldol Reactions: Highly Enantio- and Diastereoselective Synthesis of *syn* Aldols from *N*-Acryloylmorpholine

Cutting costs, cutting corners: In an inexpensive and straightforward synthesis of *syn* propionamide aldols, formation the *Z* enolborinate by the hydroboration of 4-acryloylmorpholine with diisopino-campheylborane ($(Ipc)_2BH$) was followed

([/]lpc)₂BH

Et₂O

0 °C, 2 h

by aldol reactions with achiral and chiral aldehydes to provide syn α -methyl- β -hydroxymorpholinecarboxamides with excellent enantio- and diastereoselectivity (see scheme; R = alkyl, alkenyl, aryl, heteroaryl).

Me

58–91%, d.r. >20:1 96–98% *ee*

o^{B([/]Ipc)₂}

Z enolate

RCHO

-78 °C overnight



Reductive Aldol Reaction

Diisopinocampheylborane-Mediated Reductive Aldol Reactions: Highly Enantio- and Diastereoselective Synthesis of *syn* Aldols from *N*-Acryloylmorpholine**

Philippe Nuhant, Christophe Allais, and William R. Roush*

The reductive aldol reaction of α,β -unsaturated carbonyl compounds is an important emerging method for stereocontrolled C–C bond formation.^[1] Numerous studies^[2-6] have focused on reductive aldol reactions of enones and enoates catalyzed by transition-metal complexes. Many such reactions provide excellent levels of enantio- and diastereoselectivity with aromatic aldehyde substrates; however, reductive aldol reactions with aliphatic aldehydes have been generally less selective.^[1,2a,b,d,i,3,4c,i,6] Very effective catalytic enantioselective aldol reactions have also been developed, including reactions with aliphatic aldehydes.^[7]

It is well-established that boron enolates are exceptionally useful intermediates for asymmetric aldol reactions.^[7] We reasoned that the synthetic utility of reductive aldol reactions could be enhanced by utilizing enolborinate intermediates, in view of the tight (B–O bond length 1.4–1.5 Å), closed, structurally well-defined transition states that are invoked to rationalize the enhanced stereochemical control in aldol reactions of enolborinates as compared to those of other metal enolates.^[7] Although examples of borane-mediated 1,4-reductions of enone and enoate Michael acceptors have been reported (including the use of diisopinocampheylborane ((Ipc)₂BH) as the reducing agent),^[8] as well as their subsequent reaction with aldehydes to give *syn* aldols, these processes have not yet been found to deliver the *syn* aldol products with synthetically useful enantioselectivity.^[8bc]

We report herein the development of a highly enantioand diastereoselective boron-mediated reductive aldol reaction that delivers *syn* aldols with exceptionally high levels of stereoselectivity (\geq 96% *ee*; \geq 20:1 d.r.). It was anticipated that the hydroboration of a Michael acceptor **1** with (Ipc)₂BH would proceed via transition state **2** and lead directly to an (*O*)-*Z* enolate (Scheme 1).^[8a-c] We suspected that *Z*–*E* enolborinate equilibration through a reversible 1,3-boratropic shift occurred in prior studies of this process to deliver

[*] Dr. P. Nuhant,^[+] Dr. C. Allais,^[+] Prof. Dr. W. R. Roush Department of Chemistry, The Scripps Research Institute 130 Scripps Way, 3A2, Jupiter, FL 33458 (USA) E-mail: roush@scripps.edu Homepage: http://www.scripps.edu/roush

[⁺] These authors contributed equally.

- [**] We thank the National Institutes of Health (GM038436) for support of this research. We also sincerely thank Prof. Glenn Micalizio and Dr. Daniel Canterbury for fruitful discussions and comments on this work.
- Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201302535.



Scheme 1. Hydroboration of α , β -unsaturated carbonyl compounds with (Ipc)₂BH and subsequent aldol reactions.

a mixture of *syn* and *anti* aldol adducts **6** and **7** from the (O)-Z and (O)-E enolborinate, respectively.^[8c,9] Therefore, two major objectives of this study became 1) the identification of a substrate that would undergo 1,4-reduction to give the (O)-Z enolborinate **3** with high kinetic (if not thermodynamic) control, and 2) the identification of a chiral hydroboration reagent capable of inducing excellent enantioselectivity in the subsequent aldol reaction. We elected to pursue (diisopinocampheyl)enolborinates, which are known to be useful intermediates for enantioselective aldol reactions, although they frequently undergo aldol reactions with only moderate levels of enantioselectivity.^[8c,10]

We selected commercially available (and very inexpensive)^[11] 4-acryloylmorpholine (8) as the substrate for the studies reported herein.^[12] Morpholine amides are a safe alternative^[13] to Weinreb amides but have similar modes of reactivity and comparable ease of manipulation.^[14] We quickly found that excellent results were obtained when the hydroboration of 8 with (¹Ipc)₂BH^[15] was performed in Et₂O at 0°C for 2 h, followed by addition of the aldehyde (0.85 equiv) at -78 °C (Scheme 2). By using this procedure, we obtained the syn- α -methyl- β -hydroxymorpholine amides **11a**-**f** in good to excellent yields (68-91%) and with excellent enantio- and diastereoselectivities (96-98% ee, d.r. > 20:1). The separation of aldols 11 from pinene-derived by-products is trivial owing to the large polarity difference and essentially involved filtration through a short column of silica gel. The enantiofacial selectivity derived from (^lIpc)₂BH in these reactions is the same as in the very well studied allylboration reaction.^[17]

www.angewandte.org

2

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

K These are not the final page numbers!



Scheme 2. Enantioselective synthesis of syn- α -methyl- β -hydroxymorpholine amides 11 from achiral aldehydes. [a] Yield of the isolated aldol 11 after column chromatography. [b] The diastereomeric ratio was determined by ¹H NMR spectroscopic analysis of the crude reaction mixture. [c] The *ee* value and absolute configuration were determined by Mosher ester analysis.^[16] DMtr = dimethoxytrityl.

The very high selectivity observed in these reactions reflects, in part, the essentially exclusive ($\geq 99\%$) formation of the (O)-Z enol diisopinocampheylborinate (Z)-9 (which we characterized by 1D and 2D NMR spectroscopy; see the Supporting Information). Isomerization of (Z)-9 to (E)-9 evidently does not occur to any significant extent owing to A^{1,3} strain between the morpholine unit and the terminal methyl substituent of the enolborinate.^[18] Most remarkable, however, is the exceptional level of enantioselectivity of these reactions, which significantly exceeds that observed in previous studies of enantioselective aldol reactions of (diisopinocampheyl)enolborinates.^[8c,10] The relative and absolute configuration determined for aldols 11 is consistent with transition state 10 being dominant in these reactions. That other aldol reactions [8c, 10] in which the $(Ipc)_2B$ auxiliary is used proceed with significantly lower levels of enantioselectivity implies that at least one heterochirally related transition state is competitive in those cases, but significantly less so in the reactions of (Z)-9 reported herein.^[19]

To test the utility of this reductive aldol procedure in more complex synthetic contexts, we examined the possibility of double asymmetric induction^[20] in aldol reactions of (Z)-9 (generated in situ from acrylamide 8 and (Ipc)₂BH as described for the reactions in Scheme 2) with four chiral aldehydes, **12a**, **12b**,^[21a] **12c**,^[21b] **12d**^[21b] (Scheme 3). The intrinsic diastereofacial preference of these aldehydes was determined to be 1.5:1 (in favor of **13a**), 1:2 (in favor of **13d**),



Scheme 3. Double asymmetric aldol reactions of chiral aldehydes and the chiral *Z* enolborinate generated from **8**. Yields of the isolated aldol adducts after column chromatography are given. Diastereomeric ratios were determined by ¹H NMR spectroscopic analysis of the crude reaction mixture. The absolute and relative configurations of **13a–h** were determined by Mosher ester analysis^[16] and the Rychnovsky acetonide method^[22] (see the Supporting Information). [a] Very slow reaction, incomplete after 48 h at -78 °C. brsm=based on recovered starting material, DMPM=3,4-dimethoxybenzyl, PMB=*p*-methoxybenzyl, TBDPS=*tert*-butyldiphenylsilyl, TBS=*tert*-butyldimethylsilyl.

^{© 2013} Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



3:1 (in favor of 13e), and 1.3:1 (in favor of 13g), respectively, in aldol reactions with the achiral enolborinate generated from 8 and dicyclohexylborane (see the Supporting Information). Remarkably, the double asymmetric aldol reactions of **12a-d** with the chiral Z enolborinate (Z)-9 derived from 8 and either (^lIpc)₂BH or (^dIpc)₂BH proceeded with excellent stereoselectivity (d.r. > 20:1; in each case, the minor diastereomer was not detected by ¹H NMR spectroscopic analysis of the crude reaction mixture) in both the stereochemically matched and mismatched combinations for each aldehyde substrate. The mismatched double asymmetric reaction of 12 c to give 13 f (56% yield, 71% based on recovered 12 c) was very slow and had not reached completion even after 48 h at -78°C; all other reactions reached completion overnight at -78°C. Given the intrinsic facial selectivity of aldehyde 12c (d.r. 3:1; see the Supporting Information), the enantiofacial selectivity of the Z enol diisopinocampheyborinate (Z)-9, expressed in energetic terms, must be at least 1.57 kcalmol⁻¹ to override the intrinsic diastereofacial preference of 12c to the extent of > 20:1. This selectivity corresponds to a reagent enantioselectivity of 96.5 % ee, which is fully consistent with the results in Scheme 1 for reactions of (Z)-9 with achiral aldehydes.

This method for the synthesis of syn- α -methyl- β -hydroxymorpholinecarboxamides **11** and **13** is a highly attractive and highly competitive alternative to existing methods for the enantioselective synthesis of syn aldols.^[1-7,23] It also sheds light on the great potential of boron-mediated reductive aldol reactions, despite the less than stellar history of the use of (diisopinocampheyl)enolborinates in enantioselective aldol transformations of achiral substrates.^[8c,10]

The aldol reactions of (Z)-9 described herein were performed under exceptionally mild and simple conditions, with no added bases. The results summarized in Scheme 2 and 3 demonstrate that standard (e.g., TBDPS, PMB, DMPM) as well as potentially sensitive protecting groups, such as dimethoxytrityl (DMTr; see 11e), are fully compatible with the reaction. The diastereo- and enantioselectivity of this procedure rivals that of the very best technology currently available.^[1-7,23] The morpholine amide unit in the aldol products exhibits ease of manipulation resembling that of Weinreb amides in subsequent steps.^[13,14] Our procedure requires only two steps and begins with the straightforward synthesis of diisopinocampheylborane.^[15] Strikingly, the cost of the raw materials required for the synthesis of enolborinate (Z)-9 (including the synthesis of diisopinocampheylborane) is less than \$0.25 per mmol scale of the aldol reaction (2012 Sigma–Aldrich prices for bulk quantities of reagents).^[11] If the cost, reagent accessibility, selectivity (both enantio- and diastereoselectivity), substrate scope, and generality are considered, as well as the ease of manipulation of the morpholine amide aldol products,^[14] we propose that the reductive aldol procedure described herein is not only the least expensive^[24] but also among the most enantio- and diastereoselective and generally applicable of currently available procedures for the synthesis of syn aldols.

In summary, we have developed a highly enantioselective synthesis of syn- α -methyl- β -hydroxymorpholine amides **11** and **13** from achiral and chiral aldehydes, respectively,

through the hydroboration of 4-acryloylmorpholine (8) with diisopinocampheylborane. This reaction produces the Z (diisopinocampheyl)enolborinate (Z)-9 with excellent selectivity, and intermediate (Z)-9 then undergoes highly enantioselective aldol reactions with achiral aldehydes (96-98% ee, Scheme 2) and equally highly diastereoselective double asymmetric reactions with a range of chiral aldehydes (Scheme 3). The exceptional enantioselectivity of this process distinguishes it from the vast majority of previously reported examples of aldol reactions of (diisopinocampheyl)enolborinates, which generally proceed with lower levels of enantioselectivity. This difference suggests that the transition-state control in the aldol reactions reported herein is more precise than in the previously studied aldol reactions of (diisopinocampheyl)enolborinates.^[8c, 10] The extension of this methodology to other aldol substrates and the synthesis of natural products is currently under investigation and will be reported in due course.

Experimental Section

4-Acryloylmorpholine (**8**; 35 μ L, 0.275 mmol) was added to a suspension of (^{*l*}Ipc)₂BH or (^{*d*}Ipc)₂BH (weighed in a glovebox; 72 mg, 0.25 mmol) or dicyclohexylborane (weighed in a glovebox; 45 mg, 0.25 mmol) in Et₂O (1.0 mL) at 0 °C, and the resulting mixture was stirred for 2 h at 0 °C, during which time it became homogeneous. The mixture was then cooled to -78 °C, the aldehyde (0.213 mmol) was added, and the mixture was stirred overnight at -78 °C. An aqueous buffer solution (pH 7, 0.5 mL), MeOH (0.5 mL), and THF (0.5 mL) were then added, and the reaction mixture was stirred for 6 h at room temperature. The aqueous phase was extracted three times with CH₂Cl₂ (10 mL), and the combined organic extracts were washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. Purification of the crude product by flash chromatography through a short plug of silica gel (1:1 CH₂Cl₂/ethyl acetate) provided the corresponding β-hydroxymorpholine amide **11** or **13**.

Received: March 26, 2013 Published online:

Keywords: aldol reaction \cdot boranes \cdot diastereoselectivity \cdot enantioselectivity \cdot morpholine amides

- a) H.-C. Guo, J.-A. Ma, Angew. Chem. 2006, 118, 362–375; Angew. Chem. Int. Ed. 2006, 45, 354–366; b) H. Nishiyama, T. Shiomi, Top. Curr. Chem. 2007, 279, 105–137; c) S. B. Han, A. Hassan, M. J. Krische, Synthesis 2008, 2669–2679; d) "Metal-Catalyzed Reductive Aldol Coupling": S. A. Garner, S. B. Han, M. J. Krische in Modern Reduction Methods (Eds.: P. Andersson, I. Munslow), Wiley-VCH, Weinheim, 2008, pp. 387–408.
- [2] For rhodium-catalyzed asymmetric reductive aldol reactions, see: a) S. J. Taylor, M. O. Duffey, J. P. Morken, J. Am. Chem. Soc. 2000, 122, 4528-4529; b) A. E. Russell, N. O. Fuller, S. J. Taylor, P. Aurriset, J. P. Morken, Org. Lett. 2004, 6, 2309-2312; c) B. M. Bocknack, L.-C. Wang, M. J. Krische, Proc. Natl. Acad. Sci. USA 2004, 101, 5421-5424; d) N. O. Fuller, J. P. Morken, Synlett 2005, 1459-1461; e) H. Nishiyama, T. Shiomi, Y. Tsuchiya, I. Matsuda, J. Am. Chem. Soc. 2005, 127, 6972-6973; f) C.-K. Jung, M. J. Krische, J. Am. Chem. Soc. 2006, 128, 17051-17056; g) S. B. Han, M. J. Krische, Org. Lett. 2006, 8, 5657-5660; h) J. Ito, T. Shiomi, H. Nishiyama, Adv. Synth. Catal. 2006, 348, 1235-1240; i) T. Shiomi, J. Ito, Y. Yamamoto, H. Nishiyama, Eur. J. Org.

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

Chem. **2006**, 5594–5600; j) T. Shiomi, H. Nishiyama, *Org. Lett.* **2007**, *9*, 1651–1654; k) T. Hashimoto, T. Shiomi, J. Ito, H. Nishiyama, *Tetrahedron* **2007**, *63*, 12883–12887; l) T. Hashimoto, J. Ito, H. Nishiyama, *Tetrahedron* **2008**, *64*, 9408–9412; m) C. Bee, S. B. Han, A. Hassan, H. Iida, M. J. Krische, *J. Am. Chem. Soc.* **2008**, *130*, 2746–2747; n) T. Shiomi, T. Adachi, J. Ito, H. Nishiyama, *Org. Lett.* **2009**, *11*, 1011–1014; o) M. Sugiura, N. Sato, Y. Sonoda, S. Kotani, M. Nakajima, *Chem. Asian J.* **2010**, *5*, 478–481.

- [3] For iridium-catalyzed asymmetric reductive aldol reactions, see: C.-X. Zhao, M. O. Duffey, S. J. Taylor, J. P. Morken, *Org. Lett.* 2001, *3*, 1829–1831.
- [4] For copper-catalyzed asymmetric reductive aldol reactions, see:
 a) H. Lam, G. J. Murray, J. D. Firth, Org. Lett. 2005, 7, 5743–5746; b) H. Lam, P. M. Joensuu, Org. Lett. 2005, 7, 4225–4228;
 c) O. Chuzel, J. Deschamp, C. Chausteur, O. Riant, Org. Lett. 2006, 8, 5943–5946; d) J. Deschamp, O. Chuzel, J. Hannedouche, O. Riant, Angew. Chem. 2006, 118, 1314–1319; Angew. Chem. Int. Ed. 2006, 45, 1292–1297; e) D. Zhao, K. Oisaki, M. Kanai, M. Shibasaki, Tetrahedron Lett. 2006, 47, 1403–1407; f) D. Zhao, K. Oisaki, M. Kanai, M. Shibasaki, J. Am. Chem. Soc. 2006, 128, 14440–14441; g) B. H. Lipshutz, B. Amorelli, J. B. Unger, J. Am. Chem. Soc. 2008, 130, 14378–14379; h) J. Deschamp, O. Riant, Org. Lett. 2009, 11, 1217–1220; i) M. Kato, H. Oki, K. Ogata, S. Fukuzawa, Synlett 2009, 1299–1302; j) J. Ou, W.-T. Wong, P. Chiu, Tetrahedron 2012, 68, 3450–3456.
- [5] For cobalt-catalyzed asymmetric reductive aldol reactions, see: R. J. Lumby, P. M. Joensuu, H. W. Lam, *Tetrahedron* 2008, 64, 7729-7740.
- [6] For the use of a tertiary amine as a hydride donor for asymmetric reductive aldol reactions, see: K. Osakama, M. Sugiura, M. Nakajima, S. Kotani, *Tetrahedron Lett.* 2012, 53, 4199–4201.
- [7] For selected reviews of enantioselective aldol reactions, see: a) C. H. Heathcock in Comprehensive Organic Synthesis, Vol. 2 (Eds.: B. M. Trost, I. Fleming), Pergamon, New York, 1991, pp. 181-238; b) B. M. Kim, S. F. Williams, S. Masamune in Comprehensive Organic Synthesis, Vol. 2 (Eds.: B. M. Trost, I. Fleming), Pergamon, New York, 1991, pp. 239-275; c) C. J. Cowden, I. Paterson, Org. React. 1997, 51, 1-200; d) R. Mahrwald in Modern Aldol Reactions, Vol. 2 (Ed.: R. Mahrwald), Wiley-VCH, Weinheim, 2004; e) S. E. Denmark, S. Fiujimori in Modern Aldol Reactions, Vol. 2 (Ed.: R. Mahrwald), Wiley-VCH, Weinheim, 2004, pp. 229-326; f) M. Shibasaki, S. Matsunaga, N. Kumagai in Modern Aldol Reactions, Vol. 2 (Ed.: R. Mahrwald), Wiley-VCH, Weinheim, 2004, pp. 197-227; g) J. S. Johnson, D. A. Nicewicz in Modern Aldol Reactions, Vol. 2 (Ed.: R. Mahrwald), Wiley-VCH, Weinheim, 2004, pp. 69-103; h) V. Bisai, A. Bisai, V. K. Singh, Tetrahedron 2012. 68. 4541-4580.
- [8] a) D. A. Evans, G. C. Fu, J. Org. Chem. 1990, 55, 5678-5680;
 b) G. P. Boldrini, F. Mancini, E. Tagliavini, C. Trombini, A. Umani-Ronchi, J. Chem. Soc. Chem. Commun. 1990, 1680-1681;
 c) G. P. Boldrini, M. Bortolotti, F. Mancini, E. Tagliavini, C. Trombini, A. Umani-Ronchi, J. Org. Chem. 1991, 56, 5820-5826;
 d) Y. Matsumoto, T. Hayashi, Synlett 1991, 349-350;
 e) A. K. Ghosh, J. Kass, D. D. Anderson, X. Xu, C. Marian, Org. Lett. 2008, 10, 4811-4814;
 f) R. R. Huddleston, D. F. Cauble, M. J. Krische, J. Org. Chem. 2003, 68, 11-14.
- [9] a) S. Masamune, S. Mori, D. van Horn, S. W. Brooks, *Tetrahedron Lett.* **1979**, *20*, 1665–1668; b) J. L. Duffy, T. P. Yoon, D. A. Evans, *Tetrahedron Lett.* **1995**, *36*, 9245–9248; c) A. Abiko, T. Inoue, S. Masamune, *J. Am. Chem. Soc.* **2002**, *124*, 10759–10764.
- [10] a) I. Paterson, J. M. Goodman, M. A. Lister, R. C. Schumann, C. K. McClure, R. D. Norcross, *Tetrahedron* 1990, 46, 4663– 4684; b) P. V. Ramachandran, D. Pratihar, Org. Lett. 2009, 11, 1467–1470. Ramachandran and Pratihar reported the synthesis of syn aldols with d.r. ≥ 94:6 and 90–98% ee in Ipc₂BOTf-

mediated aldol reactions of methyl propionate (Tf = trifluoromethanesulfonyl). However, after repeated attempts, the best selectivity we observed for the *syn* aldol was d.r. 2:1 and 79% *ee* from the reaction of methyl propionate and cinnamaldehyde by their reported procedure.

- [11] Cost of reagents used in the synthesis of (Ipc)₂BH and enolborinate (Z)-9 (2012 Sigma–Aldrich prices): N-acryloyl morpholine (\$168 per 250 mL, or \$0.008 per mmol); (+)-pinene (\$72 per kilogram, or \$0.01 per mmol); (-)-pinene is less expensive than (+)-pinene; borane–dimethyl sulfide (\$550 for 800 mL of a 10.0 M solution, or \$0.07 per mmol).
- [12] The Weinreb amide of acrylic acid failed to undergo the reductive aldol reaction, presumably owing to chelation of boron by the *N*-methoxy group after the 1,4-reduction.
- [13] a) M. M. Jackson, C. Leverett, J. F. Toczko, J. C. Roberts, *J. Org. Chem.* 2002, 67, 5032–5035; b) R. Peters, P. Waldmeier, A. Joncour, *Org. Process Res. Dev.* 2005, 9, 508–512.
- [14] a) R. Martín, P. Romea, C. Tey, F. Urpi, J. Vilarrasa, Synlett 1997, 1414–1416; b) J. M. Concellón, H. Rodríguez-Solla, C. Méjica, E. G. Blanco, Org. Lett. 2007, 9, 2981–2984; c) F. Dhoro, T. E. Kristensen, V. Stockmann, G. P. A. Yap, M. A. Tius, J. Am. Chem. Soc. 2007, 129, 7256–7257; d) J. M. Concellón, H. Rodríguez-Solla, P. Díaz, J. Org. Chem. 2007, 72, 7974–7979; e) K.-W. Lin, C.-H. Tsai, I.-L. Hsieh, T.-H. Yan, Org. Lett. 2008, 10, 1927–1930; f) J. M. Concellón, H. Rodríguez-Solla, V. del Amo, P. Díaz, Synthesis 2009, 2634–2645; g) C. Rye, D. Barker, Synlett 2009, 3315–3319.
- [15] a) H. C. Brown, B. Singaram, J. Org. Chem. 1984, 49, 945–947;
 b) as defined by Yamamoto, (^lIpc)₂BH derives from (-)-α-pinene, whereas (^dIpc)₂BH derives from (+)-α-pinene: Y. Yamamoto, N. Asao, Chem. Rev. 1993, 93, 2207–2293.
- [16] a) J. A. Dale, H. S. Mosher, J. Am. Chem. Soc. 1973, 95, 512–519; b) I. Ohtani, T. Kusumi, Y. Kashman, H. Kakisawa, J. Am. Chem. Soc. 1991, 113, 4092–4096.
- [17] a) P. K. Jadhav, K. S. Bhat, P. T. Perumal, H. C. Brown, J. Org. Chem. 1986, 51, 432–439; b) H. C. Brown, K. S. Bhat, R. S. Randad, J. Org. Chem. 1989, 54, 1570–1576; c) U. S. Racherla, H. C. Brown, J. Org. Chem. 1991, 56, 401–404.
- [18] R. W. Hoffmann, Chem. Rev. 1989, 89, 1841–1860.
- [19] For computational studies of aldol reactions of enolborinate intermediates, see: a) A. Bernardi, A. M. Capelli, C. Gennari, J. M. Goodman, I. Paterson, J. Org. Chem. 1990, 55, 3576–3581;
 b) Y. Li, M. N. Paddon-Row, K. N. Houk, J. Org. Chem. 1990, 55, 481–493; c) F. Bernardi, M. A. Robb, G. Suzzi-Valli, E. Tagliavini, C. Tromboni, A. Umani-Ronchi, J. Org. Chem. 1991, 56, 6472–6475.
- [20] S. Masamune, W. Choy, J. S. Petersen, L. R. Sita, Angew. Chem. 1985, 97, 1–31; Angew. Chem. Int. Ed. Engl. 1985, 24, 1–30.
- [21] a) P. Nuhant, J. Kister, R. Lira, A. Sorg, W. R. Roush, *Tetrahedron* 2011, 67, 6497–6512; b) R. Lira, W. R. Roush, *Org. Lett.* 2007, 9, 4315–4318.
- [22] S. D. Rychnovsky, D. J. Skalitzky, *Tetrahedron Lett.* **1990**, *31*, 945–948.
- [23] a) D. A. Evans, J. Bartroli, T. L. Shih, J. Am. Chem. Soc. 1981, 103, 2127–2129; b) M. T. Crimmins, B. W. King, A. E. Tabet, J. Am. Chem. Soc. 1997, 119, 7883–7884; c) M. T. Crimmins, K. Chaudhary, Org. Lett. 2000, 2, 775–777.
- [24] By comparison, the cost of valine-derived *N*-propionyl oxazolidinone (e.g., Evans aldol reagent)^[23] is \$20 per mmol (2012 Sigma–Aldrich). The current cost of the chiral oxazolidinone (the use of which requires an N-acylation step prior to the aldol reaction) is \$6.80 per mmol. The cost of the parent (*S*)-valinol (the less expensive of the two valinol enantiomers), common to both the Evans and Crimmins aldol methods, is \$1.90 per mmol, but two additional synthetic steps are required to generate the reagents used in the aldol experiments. Virtually all of the catalytic enantioselective methods^[1–7] currently available require

Angew. Chem. Int. Ed. 2013, 52, 1-6

These are not the final page numbers!

www.angewandte.org

Angewandte Communications

expensive transition-metal catalysts and/or expensive chiral ligands (many of which require multistep synthesis if not commercially available). For example, $Rh(cod)_2OTf$ and [{Rh-(cod)Cl}_2] (cod = 1,5-cyclooctadiene), two of the least expensive and most accessible Rh¹ catalyst starting materials used in catalytic enantioselective reductive aldol reactions,^[2] cost \$62 and \$86 per mmol, respectively (a 5% Rh¹ loading is used in

many of the reported examples; therefore, the cost of the Rh^I catalyst is 3-5 for an aldol reaction on a 1 mmol scale). (*R*)-Binap, one of the least expensive widely available chiral phosphine ligands, costs \$80 per mmol; hence, the cost of this ligand when used with a 5 mol% catalyst loading in a catalytic enantioselective reductive aldol reaction is approximately \$5 per mmol scale of the aldol reaction.

6 www.angewandte.org

C 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angew. Chem. Int. Ed. 2013, 52, 1-6

These are not the final page numbers!