## **ARTICLE IN PRESS**

Bioorganic & Medicinal Chemistry Letters xxx (2014) xxx-xxx

Contents lists available at ScienceDirect



**Bioorganic & Medicinal Chemistry Letters** 

journal homepage: www.elsevier.com/locate/bmcl

# Synthesis and biological evaluation of novel trichodermin derivatives as antifungal agents

Min Zheng, Ting-Ting Yao, Xiao-Jun Xu, Jing-Li Cheng, Jin-Hao Zhao\*, Guo-Nian Zhu

College of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310029, China

#### ARTICLE INFO

Article history: Received 10 March 2014 Revised 13 May 2014 Accepted 15 May 2014 Available online xxxx

Keywords: Trichodermin derivatives Synthesis Antifungal activities

## ABSTRACT

To discover more potential antifungal agents, 17 novel trichodermin derivatives were designed and synthesized by modification of **3** and **4a**. The structures of all the synthesized compounds were confirmed by <sup>1</sup>H NMR, ESI-MS and HRMS. Their antifungal activities against *Ustilaginoidea oryzae* and *Pyricularia oryzae* were evaluated. Most of the target compounds showed potent inhibitory activity, in which **4g** showed superior inhibitory effects than **4a** and commercial fungicide prochloraz. Furthermore, **4h** demonstrated comparable inhibitory activity to **4a**. Moreover, **4i** and **4l** exhibited excellent inhibitory activity for *Pyricularia oryzae*. Additionally, compound **9** was found to be more active against all tested fungal strains than **3**, with EC<sub>50</sub> values of 0.47 and 3.71 mg L<sup>-1</sup>, respectively.

© 2014 Elsevier Ltd. All rights reserved.

Fungal diseases have long been one of the major causes of crop losses, and to control the epidemic of these diseases has a vital importance.<sup>1</sup> To date, the commonly treatment method for the epidemic spread of fungal diseases is the application of chemical fungicide. The excessive use of these synthetic chemicals over the years, however, has led to the rapid development of fungal diseases more precarious.<sup>2–4</sup> Therefore, there is an urgent need for the development of novel, effective and environmentally-friendly agents to replace the conventional chemical fungicides.

Natural products have received a considerable attention as shown by the numerous studies published on their broad-spectrum biological activities, such as insecticidal, anti-inflammatory, antifungal, antitumor and antibacterial activities.<sup>5–11</sup> Trichodermin (**1**, Fig. 1), a naturally occurring sesquiterpene antibiotic, was isolated from metabolites of fungi, which had been found to possess excellent antifungal activity.<sup>12,13</sup> Trichodermin was considered as one of the ideal lead compounds for new fungicides because of its unique mechanism of action that could inhibit protein synthesis in eukaryotes.<sup>14–16</sup>

As far as structure-activity relationship is concerned, in our previous studies, series of structure modifications on trichodermin mother nucleus have been carried out, indicating that appropriate substituents introduced to the C-4 and C-8 position could help improve the antifungal activities of the compounds, and the conjugated structure in molecular could contribute to enhancing their

http://dx.doi.org/10.1016/j.bmcl.2014.05.047 0960-894X/© 2014 Elsevier Ltd. All rights reserved. bioactivity.<sup>17,18</sup> Among these compounds, **3** and **4a**, which all contained conjugated structure, exhibited relatively stronger antifungal activities against *Ustilaginoidea oryzae* (*U. ory.*) and *Pyricularia oryzae* (*P. ory.*) than trichodermin (Fig. 1). Therefore, inspired by the above results, we designed and synthesized a series of novel trichodermin derivatives based on the general structures of **3** and **4a** to better understand the structure–activity relationship.

The synthesis of the key intermediates **2** and **3** were readily obtained in one step from starting material trichodermin. The synthetic pathway was depicted in Scheme 1. In our previous manuscript, we disclosed a method to obtain **2** and **3** by reacting **1** with stannic oxide in 1,4-dioxane.<sup>18</sup> Through this method, **2** could be obtained as expected. However, intermediate **3** as by-product resulted in a relatively low yield. To optimize this reaction, several different solvent systems were investigated (Table 1). According to the screening of solvent systems, acetonitrile/H<sub>2</sub>O (10:1, v/v) (Table 1, entry 3) presented a relatively higher yield of both **2** and **3**, which was then introduced to produce these two key intermediates.

In an attempt to explore the electronic effect of the substituent in the benzene ring on the investigated biological activity, the desired compounds **4a–4m** were prepared through treating **2** with substituted cinnamoyl chloride in the presence of triethylamine as acid acceptor and *N*,*N*-dimethylaminopyridine as catalyst, as shown in Scheme 2. In addition, considering that the chirality of drug isomer sometimes has significant effects on their biological activity,<sup>19,20</sup> we designed **4n** (cis-conformer of **4a**) via the route outlined in Scheme 2. On the other hand, we also did some proper structure modification on **3** with instability of aldehyde group taken into account. As outlined in Scheme 3, thiazole amine, which

<sup>\*</sup> Corresponding author. Tel.: +86 (0)571 86971220; fax: +86 (0)571 6430193. *E-mail address: jinhaozhao@zju.edu.cn* (J.-H. Zhao).

## **ARTICLE IN PRESS**

M. Zheng et al./Bioorg. Med. Chem. Lett. xxx (2014) xxx-xxx



Figure 1. Chemical structures of trichodermin (1), compounds 3 and 4a.



Scheme 1. Synthetic route to compounds 2 and 3.

 Table 1

 The optimization of compounds 2 and 3

Entry	Solvent	Yield (%)	
		2	3
1	1,4-Dioxane/H <sub>2</sub> O (10:1, v/v)	50	25
2	1,4-Dioxane /TBHP <sup>a</sup> (10:1, v/v)	32	15
3	Acetonitrile/H <sub>2</sub> O (10:1, v/v)	50	35
4	Acetonitrile/TBHP (10:1, v/v)	45	20
5	Dicloromethane/TBHP (10:1, v/v)	36	22
6	Ethanol (95%)	26	12
7	THF <sup>b</sup>	70	5
8	Xylene	56	2
9	Acetone	30	1

<sup>a</sup> TBHP: *tert*-butyl hydroperoxide.

<sup>b</sup> THF: tetrahydrofuran.

is usually used as antifungal pharmacophore, was reacted with **3** in dry methanol to give compounds **7a** and **7b**. Compound **8** was prepared by treatment of **3** with potassium carbonate, which reacted further with acetyl chloride to yield **9** (to improve its lipophilicity). The structures of all the synthesized compounds were confirmed by <sup>1</sup>H NMR, ESI-MS and HRMS analyses.

In our previous study, the 3D structural information of compound **4a** was confirmed by single-crystal X-ray diffraction analysis.<sup>18</sup> In this Letter, in order to identify the stereostructure

of **3**, we were successful in preparing X-ray quality single crystal of intermediate **3** obtained by slow evaporation of its 95% ethanol solution (CCDC 934209). As shown in Figure 2, the crystal was monoclinic with space group *P212121*, and the cell parameters were a = 7.1994(3) Å, b = 12.9564(5) Å, c = 16.9369(7) Å, V = 1579.85(11) Å<sup>3</sup>, Z = 4. The final R = 0.0398, wR = 0.0987, S = 1.000, the difference density were <0.001 e/Å<sup>3</sup> (max) and -0.145 e/Å<sup>3</sup> (min).

The in vitro antifungal activities of all the target compounds have been evaluated against two representative pathogenic fungi including *U. ory.* (ACCC 36443) and *P. ory.* (ACCC 37631) by the mycelium growth rate method.<sup>21,22</sup> Commercial fungicide prochloraz was used as a comparative control. Each treatment was performed three times, and the effective concentration 50 ( $EC_{50}$ ) defined as the concentration required to inhibit 50% of the fungal growth, was used to describe the inhibitory activity of the compounds. The antifungal activities of the target compounds along with the standard drug for comparison are summarized in Tables 2 and 3.

Initially, we aimed to investigate the influence of substituent in the benzene ring on the antifungal activity. From the data presented in Table 2, we can see that when **4a** was modified by varying the substituents on the benzene ring, the antifungal activities of these target compounds were changed obviously. When *para*-position of the benzene ring of **4a** was substituted with a



Scheme 2. Reagents and conditions: (a) Substituted cinnamoyl chloride, Et<sub>3</sub>N, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, rt, 75–85%; (b) CrO<sub>3</sub>, pyridine, 80 °C, 3 h, 68%; (c) KBH<sub>4</sub>, NH<sub>4</sub>Cl, H<sub>2</sub>O, CH<sub>3</sub>OH, 24 h, rt, 45%; (c) Cinnamoyl chloride, Et<sub>3</sub>N, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, rt, 12 h, 84%.

Please cite this article in press as: Zheng, M.; et al. Bioorg. Med. Chem. Lett. (2014), http://dx.doi.org/10.1016/j.bmcl.2014.05.047



Scheme 3. Reagents and conditions: (a) H<sub>2</sub>NR<sup>2</sup>, CH<sub>3</sub>OH, 45 °C, 12 h, 70–76%; (b) NH<sub>2</sub>OH·HCl, K<sub>2</sub>CO<sub>3</sub>, CH<sub>3</sub>OH, reflux, 5 h, 75%; (c) CH<sub>3</sub>COCl, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, rt, 1 h, 71%.



Figure 2. The single-crystal X-ray crystallography of 3.

 Table 2

 Antifungal activity of 4a-4n against U. ory. and P. ory

Compd	R <sup>1</sup>	(R) or (S)	Antifungal activity (EC <sub>50</sub> , mg $L^{-1}$ )	
			U. ory.	P. ory.
4a	Н	(R)	1.48	0.74
4b	4-CH <sub>3</sub>	(R)	6.21	5.99
4c	2-0CH <sub>3</sub>	(R)	4.23	6.18
4d	$4-OCH_3$	(R)	6.86	6.68
4e	3,4,5-(OCH <sub>3</sub> ) <sub>3</sub>	(R)	11.04	11.70
4f	3-CF <sub>3</sub>	(R)	10.31	12.47
4g	2-F	(R)	0.56	0.53
4h	4-F	(R)	1.47	1.09
4i	2-Cl	(R)	3.12	1.31
4j	2-Br	(R)	3.17	2.70
4k	3-Br	(R)	6.20	3.82
41	4-Br	(R)	11.28	12.62
4m	2-Cl-4-F	(R)	2.45	1.47
4n	Н	(S)	>20	>20
Prochloraz	_	-	0.82	0.96

methyl (compound **4b**) or methoxy (compound **4d**) group, their fungicidal activity generally decreased. However, introducing halogen atoms into the benzene ring of **4a** led to more than one half compounds showed remarkable antifungal activity. To our delight, both of the compounds **4g** and **4h** containing a fluorine atom showed competitive inhibitory activities to **4a**. Especially compound **4g** (2-fluorine) exhibited the better inhibitory activity against all two tested fungi than prochloraz (EC<sub>50</sub> values: 0.56 vs 0.82 mg L<sup>-1</sup>; 0.53 vs 0.96 mg L<sup>-1</sup>). Altering the substitution groups at the *ortho*-position of the benzene ring with F, Cl, Br systematically (**4g**, **4i**, **4j**), we found the antifungal activity against *U. ory.* decreased in that order, and the same trend was observed in the activity against *P. ory.* Using 3,4,5-triethoxy group (**4e**) instead of a methoxy to determine its importance, the results showed that there was remarkable activity decrease against all two tested fungi.

Table 3											
Antifungal	activity	of 7a,	7b, 8	<b>3</b> , <b>9</b> , and	3	against	U.	ory.	and	Р.	ory.

Compd	Antifungal activity ( $EC_{50}$ , mg $L^{-1}$ )		
	U. ory.	P. ory.	
7a	2.54	13.27	
7b	1.52	12.48	
8	3.67	8.91	
9	0.47	3.71	
3	0.80	11.82	
Prochloraz	0.82	0.96	

These results demonstrated that the volume of the substituent was very important for the inhibitory activity of *U. ory.* and *P. ory.* Notably, the in vitro antifungal profile of **4b–4m** is shown to be increased when groups such as –OCH<sub>3</sub>, –F and –Br are present at *ortho* position as is also inferred by Table 2. For example, with the comparisons of compounds **4j** (*o*-Br), **4k** (*m*-Br) and **4l** (*p*-Br), it can be seen that their inhibitory capacities in decreasing order were: **4l**, **4k** and **4j**. Changing the configuration of the **4a** also led to a remarkable destruction of inhibitory activity in both *U. ory.* and *P. ory.* The result indicated that the trans-isomer at the C-8 position was crucial for the antifungal activity.

On the other side, some preliminary structure–activity relationships on the **3** derivatives were also investigated. As shown in **Table 3**, compound **9** showed better inhibitory activity against all two tested fungi than **3**. Notably, compound **9** also exhibited more promising antifungal activity than prochloraz (an  $EC_{50}$  value: 0.47 vs 0.82 mg L<sup>-1</sup>). However, the loss of the antifungal activities were observed for compounds **7a** and **7b** containing a thiazole amine group, revealing that thiazole amine group was not favorable for the antifungal activity. In addition, compound **8** by replacement aldehyde group with oxime group obviously decreased the antifungal activity against *U. ory.* (an  $EC_{50}$  value: 3.67 vs 0.80 mg L<sup>-1</sup>). These above results suggested that improving the lipophilicity of trichodermin derivatives was more beneficial for the bioactivity.

In summary, we have developed a series of trichodermin derivatives based on the general structures of **3** and **4a**. Antifungal activities of these compounds were evaluated. The SARs revealed that the nature of the substituent, the lipophilicity and the chirality of the compounds were very important for the antifungal activity. The biological test results showed that these compounds displayed moderate to high antifungal activity against *U. ory.* and *P. ory.* Among them, **4g** showed better antifungal activity than the lead compound **4a** and prochloraz. Furthermore, **4h** exhibited excellent inhibitory activity compared with **4a** and prochloraz. Moreover, **4i** and **4l** demonstrated competitive inhibitory activity against *P. ory.* to **4a**. Additionally, compound **9** displayed superior activity against all two tested fungi compared with **3**, with EC<sub>50</sub> values of 0.47 and

## **ARTICLE IN PRESS**

 $3.71 \text{ mg L}^{-1}$ , respectively. Although some of them were still less active than **4a** and **3**, they can be used as a good starting point for further optimization.

#### Acknowledgments

This work was supported by the National Key Technology R&D Program of China (2011BAE06B04-10). The authors also thank senior engineer Jianming Gu (Center of Analysis and Measurement of Zhejiang University) for the X-ray crystallographic diffraction analysis of compound **3**.

## Supplementary data

Supplementary data associated with this article can be found, in the online version, at http://dx.doi.org/10.1016/j.bmcl.2014.05. 047.

### **References and notes**

- Jach, G.; Görnhardt, B.; Mundy, J.; Logemann, J.; Pinsdorf, E.; Leah, R.; Schell, J.; Maas, C. Plant J. 1995, 8, 97.
- 2. Russell, P. E. J. Agric. Sci. 1995, 124, 317.
- 3. Zhang, C. Q.; Liu, Y. H.; Zhu, G. N. Eur. J. Plant Pathol. 2010, 126, 509.

- 4. Latorre, B. A.; Spadaro, I.; Rioja, M. E. Crop Prot. 2002, 21, 957.
- 5. Manici, L. M.; Lazzeri, L.; Palmieri, S. J. Agric. Food Chem. **1997**, 45, 2768.
- Martin, B.; Hernandez, S.; Silvarrey, C.; Jacas, J. A.; Cabaleiro, C. Phytopathol. Mediterr. 2005, 44, 169.
- Luo, R.; Wang, J. B.; Zhao, L.; Lu, N.; You, Q. D.; Guo, Q. L.; Li, Z. Y. Bioorg. Med. Chem. Lett. 2014, 24, 1334.
- Lv, J.; Guo, N.; Wen, S. P.; Teng, Y. O.; Ma, M. X.; Yu, P. J. Asian Nat. Prod. Res. 2013, 15, 867.
- 9. Saeed, A.; Rafique, H.; Ashraf, Z. J. Asian Nat. Prod. Res. 2013, 15, 130.
- Qu, H.; Yu, X.; Zhi, X. Y.; Lv, M.; Xu, H. Bioorg. Med. Chem. Lett. 2013, 23, 5552.
   Wang, Y.; Shao, Y. H.; Wang, Y. Y.; Fan, L. L; Yu, X.; Zhi, X. Y.; Yang, C.; Qu, H.; Yao, X. J.; Xu, H. J. Agric. Food Chem. 2012, 60, 8435.
- 12. Cutler, H. G.; LeFiles, J. H. Plant Cell Physiol. 1978, 19, 177.
- 13. Nielsen, K. F.; Gräfenhan, T.; Zafari, D.; Thrane, U. J. Agric. Food Chem. 2005, 53, 8190.
- 14. Wei, C. M.; Hansen, B. S.; Vaughan, M. H.; McLaughlin, C. S. Proc. Natl. Acad. Sci. U.S.A. 1974, 71, 713.
- 15. Stafford, M. E.; McLaughlin, C. S. J. Cell Physiol. 1973, 82, 121.
- 16. Carter, C. J.; Cannon, M.; Smith, K. E. Biochem. J. 1976, 154, 171.
- 17. Cheng, J. L.; Zhou, Y.; Zhao, J. H.; Zhang, C. L.; Lin, F. C. *Chin. Chem. Lett.* 2010, 21, 1037.
- Xu, X. J.; Cheng, J. L.; Zhou, Y.; Zhang, C. L.; Ou, X. M.; Su, W. K.; Zhao, J. H.; Zhu, G. N. Chem. Biodivers. 2013, 10, 600.
- Zhou, G. C.; Weng, Z. B.; Shao, X. X.; Liu, F.; Nie, X.; Liu, J. S.; Wang, D. C.; Wang, C. G.; Guo, K. Bioorg. Med. Chem. Lett. 2013, 23, 6549.
- 20. Xu, H.; He, X. Q. Bioorg. Med. Chem. Lett. 2010, 20, 4503.
- Vicentini, C. B.; Forlani, G.; Manfrini, M.; Romagnoli, C.; Mares, D. J. Agric. Food Chem. 2002, 50, 4839.
- Xiao, Y. S.; Yan, X. J.; Xu, Y. J.; Huang, J. X.; Yuan, H. Z.; Liang, X. M.; Wang, D. Q. Pest Manag. Sci. 2013, 69, 814.

4