View Article Online

MedChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: P. M. Cheuka, N. Lawrence, D. Taylor, S. Wittlin and K. Chibale, *Med. Chem. Commun.*, 2018, DOI: 10.1039/C8MD00382C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the **author guidelines**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the ethical guidelines, outlined in our <u>author and reviewer resource centre</u>, still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/medchemcomm

YAL SOCIETY CHEMISTRY

Journal Name

ARTICLE

Received 00th January 20xx, Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Antiplasmodial imidazopyridazines: structure-activity relationship studies lead to the identification of analogues with improved solubility and hERG profiles[†]

Peter Mubanga Cheuka,^a Nina Lawrence,^b Dale Taylor,^b Sergio Wittlin^{cd} and Kelly Chibale^{abef}*

3,6-Diarylated imidazopyridazines have recently been shown to possess good *in vitro* antiplasmodial and *in vivo* antimalarial activity. However, frontrunner compounds have been associated with poor solubility and a hERG (human *ether-a-go-go*-related gene) inhibition liability raising concerns for potential cardiotoxicity risks. Herein, we report the synthesis and structure-activity relationship studies of new imidazopyridazines aimed at improving aqueous solubility and countering hERG inhibition while maintaining antiplasmodial potency. While we identified new analogues with potent antiplasmodial activity ($IC_{50} = 0.031 \mu$ M against the NF54 drug-sensitive strain, and $IC_{50} = 0.0246 \mu$ M against the K1 multidrug resistant strain), hERG inhibition profile ($IC_{50} = 7.83 - 32.3 \mu$ M) with sub-micromolar antiplasmodial activity (NF54, $IC_{50} = 0.151 - 0.922 \mu$ M) were identified. Similarly, the introduced molecular features also resulted in analogues with moderate to high solubility ($60 - 200 \mu$ M) while also displaying sub-micromolar antiplasmodial potency (NF54, $IC_{50} = 0.136 - 0.99 \mu$ M).

Introduction

According to the World Health Organization (WHO) estimates, malaria was responsible for 445 000 deaths and 216 million cases in 2016.¹ The WHO African region accounted for 90% of all malaria cases and 91% of malaria deaths. Children under the age of 5 years are particularly susceptible to infection, illness and death in areas with intense transmission of malaria. Although the under-5 malaria death rate has declined by 29% worldwide, malaria remains a major killer of children in this age group claiming the life of a child every 2 minutes.¹

Malaria is caused by five species of protozoan parasites of the genus *Plasmodium* which are transmitted to humans through

^{b.} Drug Discovery and Development Centre (H3D), Division of Clinical Pharmacology, Department of Medicine, University of Cape Town, Observatory, Cape Town 7925, South Africa.

^{e.} Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Rondebosch 7701, South Africa.

* Corresponding author: (KC) E-mail: Kelly.Chibale@uct.ac.za. Phone: +27-21-6502553. Fax: +27-21-6505195.

the bites of infected female *Anopheles* mosquitoes.¹ Of the five species known to infect humans, *Plasmodium falciparum* and *Plasmodium vivax* are the most virulent. On the African continent, *P. falciparum* is the most prevalent and causes the most malaria-related deaths worldwide while *P. vivax* dominates most countries outside of sub-Saharan Africa.¹

Once effective antimalarial drugs like chloroquine and sulfadoxine-pyrimethamine have suffered widespread resistance, which has undermined malaria control efforts.^{2–5} Therefore, antimalarial chemotherapy has shifted towards the use of artemisinin combination therapies (ACTs). Regrettably, emerging resistance to artemisinins has been reported recently in 5 countries of the Greater Mekong sub-region, which includes Cambodia, Lao People's Democratic Republic, Myanmar, Thailand and Viet Nam.¹ Therefore, with this threat of resistance to first line treatment options, there is a critical need to intensify research efforts into novel, affordable and structurally diverse antimalarials.

Imidazopyridazines possess a range of biological activities including kinase inhibition,^{6,7} anxiety treatment⁸ and anticancer potential.⁹ These compounds have also been investigated as potential novel antimalarial agents.^{10–13} More recently, 3,6-diarylated imidazopyridazines with potent antiplasmodial activity were identified through a high throughput screening (HTS) of a SoftFocus kinase library.¹⁴ Further medicinal chemistry optimization led to the identification of, among others, the lead compound **1** (figure 1) with high *in vitro* potency against both the NF54 (IC₅₀ = 7.3 nM) and the K1 (IC₅₀ = 6.3 nM) strains of *P. falciparum*.¹⁴ Compound **1** also showed good oral efficacy (98% at 4 × 50 mg/kg) in the *P. berghei*-infected mouse model.¹⁴ However, the compound exhibited poor solubility (< 5 μ M at pH 6.5) and a hERG liability (IC₅₀

^{a.} Department of Chemistry, University of Cape Town, Rondebosch 7701, South Africa.

^c Swiss Tropical and Public Health Institute, Socinstrasse 57, 4002 Basel, Switzerland.

^{d.} University of Basel, 4003 Basel, Switzerland.

^{f.} South African Medical Research Council Drug Discovery and Development Research Unit, University of Cape Town, Rondebosch 7701, South Africa.

^{*}Electronic Supplementary Information (ESI) available: [Synthetic and characterization procedures for selected compounds, methods for *in vitro* antiplasmodial evaluation, solubility determination, cytotoxicity evaluation, hERG inhibition and copies of NMR spectra]. See DOI: 10.1039/x0xx00000x

= 0.9 μ M). Additionally, the compound only exhibited a mean survival days (MSD) time of 7 days.



P. f, NF54 IC_{50} = 7.3 nM P. f, K1 IC_{50} = 6.3 nM In vivo P. berghei (po) at 4 × 50 mg/kg, 98%, 7 MSD hERG IC_{50} = 0.9 μ M

ARTICLE

P. f, NF54 IC₅₀ = 1.1 nM
 In vivo P. berghei (po) at 4
 × 50 mg/kg, 99.8%, > 30
 MSD, 3 out of 3 mice cured

Fig. 1 Previously explored antimalarial imidazopyridazine analogues

In an article published in the same year, Le Manach and colleagues¹⁵ further explored the structure-activity relationship (SAR) of this class of molecules in the hope of addressing the aforementioned shortcomings. Although compound 2 (figure 1), identified from this study, completely cured P. berghei-infected mice at 4 × 50 mg/kg/day, better efficacy at lower doses was not achieved. The lead compound also retained sub-optimal solubility (< 5 μ M at pH 6.5) and some off-target activities such as hERG inhibition. It is noteworthy that this sulfoxide-substituted analogue has no intrinsic hERG potency. But it is rapidly metabolized in vivo to the sulfone analogue which has potent hERG activity ($IC_{50} = 0.4$ $\mu M).^{^{15}}$ In a more recent paper, further expansion of SAR based on modifications to the imidazopyridazine core aimed at improving pharmacokinetics (PK), in vivo efficacy and selectivity over hERG were reported.¹⁶ This led to the identification of the pyrazolopyridine analogue in which the two substituents in compound 2 were retained. Despite being able to completely cure P. berghei-infected mice at 4×50 mg/kg, the scaffold-hoped analogue still exhibited a hERG liability and poor solubility.¹⁶

Recently, there has been a significant proportion of drug withdrawals arising from their unwanted prolongation of the time between the start of the Q wave and the end of the T wave (QT interval) in an electrocardiogram.¹⁷ Such a phenomenon, which can lead to malignant ventricular arrhythmia and sometimes sudden cardiac death, is known to arise from, mainly, the blockade of the potassium ion channel (I_{Kr}) encoded by the hERG gene.¹⁷ There are a number of strategies used to address the hERG liability in drug molecules.¹⁸ Herein, we report the application of some of these strategies in designing imidazopyridazine analogues (figure 2) with potentially reduced in vitro hERG inhibition and improved solubility, which also served to expand the antiplasmodial SAR around this class of molecules. These approaches include subtle modifications (analogues 3 - 5, 13 and 28 - 32); replacing aromatic rings with saturated systems to discourage π - π stacking with aromatic residues in the large cavity of the hERG channel (analogues 19 -21); introducing polar functionalities into the molecule to potentially disrupt lipophilic drug-hERG interactions (analogues 14 -27).

In a quest to address solubility issues associated with the recently studied antimalarial imidazopyridazines, we employed known strategies, which included introducing polar water-solubilizing groups (analogues 6 – 12 and 14 – 27) and disruption of molecular planarity to discourage the solubility-suppressing effect of $\pi - \pi$ stacking (analogues 19 – 21).^{19–21}

While changes were introduced at positions 3 and 6 of the imidazopyridazine core-scaffold, the 4-methylsulfinylphenyl and 3-methylsulfinylphenyl groups were fixed at these positions in most analogues (analogues 3, 5 - 8, and 10 - 27). Le Manach and coworkers have recently employed such a prodrug strategy by synthesizing solubility-enhancing sulfoxides which were shown to be metabolized *in vivo* to the corresponding active sulfones.¹⁵ In two cases, sulfone-containing derivatives, **4** and **9**, were also synthesized for comparison purposes. For derivatives bearing sulfoxide-containing phenyl rings at the point of variation, sulfone counterparts were synthesized for the same reason in some cases (e.g., analogues **3** vs **5**, **14** vs **15**, **16** vs **17**, **24** vs **27**, and **25** vs **26**).

Results and Discussion

Chemistry

Analogues 1, 4 - 12, 14 - 23 and 29 - 32 were synthesized according to scheme 1. The intermediate 2a was obtained by condensation/cyclization of commercially available 6chloropyridazin-3-amine (1a) with bromoacetaldehyde diethylacetal in the presence of hydrobromic acid (HBr).¹⁴ A regioselective electrophilic aromatic iodination using N-iodosuccinimide (NIS) afforded the iodinated intermediate, **3a**, in 81% yield.¹⁴ Sequential Suzuki-Miyaura cross coupling reactions on intermediate 3a delivered the diaryl-imidazopyridazines (1, 4 - 12 and 29 - 32) in 13 - 80% yield.¹⁴ The aminated target molecules, 14 - 23, were realized, in poor to low yields (8 - 30%), via a palladium-mediated Buchwald-Hartwig amination^{22,23} of the chloro-bearing intermediate 4a. The aniline starting materials 5d and 5h, which were used to synthesize analogues 22 and 23 respectively, were not available commercially. These were synthesized in-house according to the synthetic schemes in the electronic supplementary information.

Schemes 2A and 2B were used to synthesize analogues **3**, **13**, **24** – **27** and **28**. Using an excess (2.2 equivalents) of the respective boronic acid, analogues **3** and **28** were obtained by a one pot Suzuki-Miyaura cross-coupling on both reaction sites of the previously synthesized intermediate **3a** (scheme 2A).¹⁴ Analogues **13** and **24** – **27** were obtained via a common bromo-substituted intermediate **4i** according to scheme 2B. Intermediate **4i** was synthesized from a previously synthesized imidazopyridazine intermediate **2a** by a Suzuki-Miyaura cross coupling followed by regioselective electrophilic aromatic substitution in presence of *N*bromosuccinimide (NBS).¹⁴ A Suzuki-Miyaura cross-coupling on intermediate **4i** furnished analogue **13** in 40% yield. The aminated derivatives **24** – **27** were obtained in poor yields (9 – 17%) via Buchwald-Hartwig amination.^{22,23}

View Article Online DOI: 10.1039/C8MD00382C ARTICLE





Scheme 1 General synthetic approach for analogues 1, 4 - 12, 14 - 23 and 29 - 32. Reagents and conditions: (i) BrCH₂CH(OEt)₂, aq HBr, EtOH/H₂O, reflux, 103 °C, 22 h; (ii) NIS, DMF, rt (21 °C), 5 d, 81%; (iii) appropriate boronic acid or boronic acid pinacol ester, Pd(PPh₃)₂Cl₂, 1 M aq K₂CO₃, DMF, 80 °C, 3.5 - 46 h, 26 - 77%; (iv) appropriate boronic acid pinacol ester, Pd(PPh₃)₂Cl₂, 1 M aq K₂CO₃, DMF, 10 °C, 4 - 21 h, 13 - 80%; (v) R₃NH₂, Pd₂(dba)₃, (R)-BINAP for 17, BrettPhos for 14, 22 & 23, XPhos for 15, 16 & 18 - 21, K₂CO₃ for 17, Cs₂CO₃ for 14, 22 & 23, NaOt-Bu for 15, 16 & 18 - 21, toluene/1,4-dioxane for 17, 1,4-dioxane for all other analogues, 100 °C for 20, 120 °C for all other analogues, sealed tube, inert atmosphere (N₂), 6 - 43 h, 8 - 30%.



Scheme 2 General synthetic approach for analogues 3, 13, 28 and 24 – 27. Reagents and conditions: (i) appropriate boronic acid (excess, 2.2 eq), Pd(PPh₃)₂Cl₂, 1 M aq K₂CO₃, DMF, 100 °C, 15 h for 3 and 24 h for 28, 39% for 3 and 62% for 28; (ii) 3-methylsulfinylphenylboronic acid, Pd(PPh₃)₂Cl₂, 1 M aq K₂CO₃, DMF, 100 °C, 3.25 h, 48%; (iii) NBS, DMF, rt (23 °C), 22 h, 59%; (iv) 3-methylsulfinylphenylboronic acid, Pd(PPh₃)₂Cl₂, 1 M aq K₂CO₃, DMF, 100 °C, 20 h, 40%; (v) R₄-NH₂, Pd₂(dba)₃, BrettPhos, Cs₂CO₃, 1,4-dioxane, 120 °C, sealed tube, 5 – 39 h, 9 – 17%.

Biology and solubility

Published on 06 September 2018. Downloaded by Kaohsiung Medical University on 9/7/2018 9:06:14 AM

A total of 31 imidazopyridazine analogues were synthesized and tested in biological and solubility assays. All the synthesized analogues were evaluated for in vitro antiplasmodial activity against the drug sensitive strain (NF54) of the P. falciparum malaria parasites. Analogues exhibiting sub-micromolar (IC₅₀ < 1 μ M) potency were further tested in vitro against the multi drug-resistant strain (K1) of P. falciparum. In parallel, the solubility of all the synthesized analogues was determined. Significantly potent (IC_{50} < 1 µM) analogues were advanced to cytotoxicity testing using the Chinese Hamster Ovary (CHO) cell line (ATCC). The selectivity index (SI) for each compound was calculated as a ratio of cytotoxic IC_{50} value to corresponding antiplasmodial (P. f NF54) IC₅₀ value. The SAR explorations are discussed with reference to IC₅₀ values on the NF54 strain. Representative compounds were selected and screened for hERG activity on the QPatch gigaseal automated patch clamp platform.

In vitro antiplasmodial activity and solubility. The *in vitro* antiplasmodial (IC_{50} values), and solubility values for all the synthesized compounds are shown in tables 1-3.

We set out by fixing a 4-methylsulfinylphenyl group at position 3 while introducing saturation, subtle modifications and water-solubilizing groups at position 6 (table 1). The antiplasmodial potency of all analogues in this series was generally compromised ($IC_{50} = 0.136 - 2.67 \mu$ M) compared to the previously reported¹⁴ lead compound **1** ($IC_{50} = 0.0034 \mu$ M). All analogues evaluated for antiplasmodial activity against both the NF54 and K1 parasites were equipotent on these two strains, suggesting the absence of cross-resistance. This is in conformity with previously reported imidazopyridazine derivatives.¹⁵ *Para* substitution with either a sulfone or sulfoxide group at 6-phenyl ring proved detrimental to potency as reflected in analogues **3** ($IC_{50} = 0.673 \mu$ M) and **5** ($IC_{50} = 0.974 \mu$ M). As anticipated, the sulfoxide-substituted analogue **3**

exhibited superior solubility (200 µM) compared to the sulfonesubstituted analogue 5 (5 µM). Amongst the analogues substituted with a fluoro and a basic side chain (6, 7 and 8, table 1), 8 proved the most potent (IC₅₀ = 0.136 μ M) signifying the importance of the para-fluoro and meta-diethylaminomethyl substitution pattern to potency. It is noteworthy that compound 8 was also the most potent in the entire series of analogues in table 1. Gratifyingly, we also managed to significantly improve solubility in such basic aminesubstituted analogues (180 - 200 µM). Although potency was, generally, compromised for the aminated imidazopyridazines, most of them still retained sub-micromolar potency. Notable amongst these is 3-methylsulfonylaniline-substituted analogue $\mathbf{14}$ (IC_{50} = 0.274 μ M) which was 3-fold more potent than the reduced form 15 $(IC_{50} = 0.820 \mu M)$. However, the para-substituted versions 16 $(IC_{50} =$ 0.552 $\mu M)$ and 17 (IC_{50} = 0.423 $\mu M)$ were equipotent. In this series, we also designed analogues with saturated cyclic groups (19 - 21) which we hypothesized could help discourage π - π stacking thereby improving aqueous solubility.²⁴ Strikingly, the sulfone group in analogue 21 (IC₅₀ = 0.498 μ M) was important in retaining submicromolar potency - analogue 20 lacking this group exhibited 5fold less potency (IC₅₀ = 2.67 μ M). All the aminated analogues, generally exhibited improved solubility with sulfone-substituted analogues showing inferior solubility relative to their sulfoxide counterparts.

Journal Name

Table 1 In vitro antiplasmodial activity against P. falciparum (NF54 and K1) and solubility of pyridazine-substituted analogues

R⁵ N N 3

		SOMe		
Compound code	B ⁵	<i>Ρ. f</i> IC ₅₀ , μM ^{c,d}		Solubility, μM,
Compound code	ĸ	NF54	K1	pH 6.5
3	MeOS	0.673	0.499	200
5	MeO ₂ S	0.974	0.82	5
6	Et ₂ N	1.88	ND^e	180
7	Et ₂ N	1.67	ND^e	195
8	Et ₂ N	0.136	0.114	200
22	Et ₂ N	1.62	ND ^e	200
23	HO Et ₂ N	0.922	0.774	195
14	MeO ₂ S	0.274	0.236	20
15	MeOS A	0.820	0.854	190
16	MeOS	0.552	0.488	195
17	MeO ₂ S	0.423	0.236	< 5
18	CN L Star	0.468	0.436	25
19	0 0 - N N - N	0.524	0.492	10
20	Me N N H	2.67	ND ^e	200
21	MeO ₂ S N	0.498	0.439	70

21 $h \to h^{-1}$ 0.498 0.439 70 ^cMean from n values of ≥ 2 independent experiments with multidrug resistant (K1) and sensitive (NF54) strains of *P. falciparum*. The majority of the individual values differed less than 2-fold (maximum 2-fold).

^dArtesunate [$IC_{50} = 2.2 \text{ ng/mL}$ (NF54), 0.93 ng/mL (K1)] and chloroquine [$IC_{50} = 4.3 \text{ ng/mL}$ (NF54), 83 ng/mL (K1)] were used as reference drugs.

^eND, not determined.

MedChemComm Accepted Manuscript

DOI: 10.1039/C8MD00382C

Inspired by previous studies¹⁴ which found *meta*substitution at the phenyl ring on position 6 optimal for potency, we kept this group fixed as 3-methylsulfinylphenyl group while exploring other changes at position 3 of the core-scaffold. The antiplasmodial and solubility results are shown in table 2. Apart from compound **13**, which exhibited promising activity against both the NF54 (IC₅₀ = 0.151 μ M) and K1 (IC₅₀ = 0.0866 μ M) strains, generally, this series represented modifications which compromised potency the most (IC₅₀ = 0.99 – 7.55 μ M). However, the introduced water-solubilizing groups imparted moderate to high solubility (35 – 200 μ M) in these analogues. An increase in dihedral angle which compromises $\pi - \pi$ stacking following *ortho* substitution of bi-aryl systems could explain the notable difference in solubility between the *ortho*-fluorinated analogue **10** (200 μ M) and the *meta*fluorinated regioisomer **11** (35 μ M).²⁴ As previously observed, sulfoxide-containing aminated analogues **26** (150 μ M) and **27** (165 μ M) were more soluble than the corresponding sulfone analogues **25** (40 μ M) and **24** (60 μ M).

Table 2 In vitro antiplasmodial activity against P. falciparum (NF54 and K1) and solubility of imidazo-substituted analogues

	-6	<i>Ρ. f</i> IC ₅₀ , μM ^{c,d}		Solubility, μM,	
Compound code	R' -	NF54	К1	pH 6.5	
10	NEt ₂	3.2	ND ^e	200	
11	ANT F	1.05	ND ^e	35	
12		0.99	0.558	200	
13	SOMe	0.151	0.0866	200	
24	HN SO ₂ Me	3.62	ND^e	60	
25	HN SO ₂ Me	7.55	ND^e	40	
26	HN SOMe	4.71	ND^e	150	
27		3.81	ND^e	165	

^cMean from n values of ≥ 2 independent experiments with multidrug resistant (K1) and sensitive (NF54) strains of *P. falciparum*. The majority of the individual values differed less than 2-fold (maximum 2-fold).

^dArtesunate [IC₅₀ = 2.2 ng/mL (NF54), 0.93 ng/mL (K1)] and chloroquine [IC₅₀ = 4.3 ng/mL (NF54), 83 ng/mL (K1)] were used as reference drugs. ^eND, not determined.

In another strategy to address the hERG liability, we designed analogues 4, 28 - 32 (table 3), 3, 5 and 13 (table 1) with subtle modifications, a strategy that has also been used by other drug discovery programmes to diminish hERG activity.¹⁸ These included peripheral positional changes of the sulfone and sulfoxide groups on the two phenyl rings as well as switching sulfones with sulfoxide groups. The sulfone version of compound 7 in table 1, compound 9 (table 3), was also included in this series of

compounds. Interestingly, compound 28, a regioisomer of the previously reported¹⁴ lead compound **1** was highly potent (IC_{50} = 0.031 μ M) indicating that the para-to-meta positional change of the right-hand-side sulfone was tolerated albeit solubility (< 5 µM) was still an issue. However, this isomer is still 9-fold less potent than the lead compound 1 (IC₅₀ = 0.0034 μ M) indicating a slight decrease in potency accompanies this positional change. While maintaining the sulfone at the *meta* position on the right-hand-side phenyl ring, para substitution on the left-hand-side phenyl ring, irrespective of the nature of substituent as shown in analogues 29 (IC₅₀ = 0.396 μ M) and **30** (IC₅₀ = 0.264 μ M) was detrimental to potency. Similarly, keeping the sulfoxide fixed at the meta position of the right-handside phenyl ring while introducing the sulfone and sulfoxide groups at the para position of the left-hand-side phenyl ring proved detrimental to potency (see analogues **31**, $IC_{50} = 1.68 \mu M$ and **32**, $IC_{50} = 0.870 \,\mu$ M). It is also noteworthy that the sulfoxide-substituted analogue 32 was twice as potent as the sulfone-substituted counterpart **31**. Furthermore, analogue **4** was 7-fold less potent than its previously reported¹⁵ regioisomer further confirming that para-substitution on the left-hand-side phenyl ring is detrimental to activity. Save for analogue 9, whose excellent solubility (200 µM) could be attributed to the water-solubilizing basic side chain, all analogues in this series having at least one sulfone group were poorly soluble (< 5 - 10 μ M). As expected, analogue 32 with sulfoxide-substituted phenyl rings, exhibited excellent solubility (200 µM).

Table 3 In vitro antiplasmodial activity against P. falciparum (NF54 and K1) and solubility for discreetly-modified analogues

Compound code	Structure	<i>P. f</i> IC₅	₀ , μM ^{c,d}	Solubility, μM,	
compound code	Structure	NF54	K1	рН 6.5	
4	MeOS N.N.	0.278	0.207	10	
9	NEt ₂ F SO ₂ Me	0.822	ND ^e	200	
1	SO ₂ Me	0.0034	0.00225	10	
28		0.031	0.0246	< 5	
29	MeO ₂ S	0.396	0.314	< 5	
30	MeOS N.N. SO ₂ Me	0.264	0.189	< 5	
31	MeO ₂ S	1.68	ND ^e	< 5	

MedChemComm Accepted Manuscript

Compound code	Structure	<i>P. f</i> IC ₅₀ , μM ^{c,d}		Solubility, μM,
compound code		NF54	K1	pH 6.5
32	Meos NN Some	0.870	0.872	200

^cMean from n values of \geq 2 independent experiments with multidrug resistant (K1) and sensitive (NF54) strains of *P. falciparum*. The majority of the individual values differed less than 2-fold (maximum 2-fold).

^dArtesunate [IC₅₀ = 2.2 ng/mL (NF54), 0.93 ng/mL (K1)] and chloroquine [IC₅₀ = 4.3 ng/mL (NF54), 83 ng/mL (K1)] were used as reference drugs. ^eND, not determined.

In vitro mammalian cytotoxicity and hERG inhibition. Analogues with *in vitro* antiplasmodial potency $IC_{50} < 1 \mu$ M were evaluated for *in vitro* cytotoxicity against the mammalian CHO cell line. Analogues screened for hERG inhibition were, generally, selected based on demonstrated sub-micromolar antiplasmodial activity. However, for derivatives representing discreet peripheral changes (e.g., **3** - **5**, **13**, **28** - **30** and **32**) with sub-micromolar antiplasmodial activity, only representative compounds were tested for hERG inhibition. Generally, most analogues were non-cytotoxic with selectivity indices in the range 72 - > 874 (table 4). Strikingly, compounds **8**, **9**, **12** and **23**, containing the basic side chains, were particularly cytotoxic with lower selectivity (SI = 10 - 30) (table 4).

The hERG inhibition results for selected analogues are summarized in table 4. As previously reported,¹⁴ compound **1** was potent on hERG (IC₅₀ = 3.61 μ M). Its isomers, **28** (IC₅₀ = 2.35 μ M) and **29** (IC₅₀ = 4.0 μ M), still retained hERG potency. When sulfones

in compound 28 were switched to sulfoxides as shown in compound 13 (IC₅₀ = 8.45 μ M), hERG potency was found to diminish by almost 4-fold. Moreover, positional change of the left-hand-side sulfoxide group in 13 to para in analogue 32 (IC_{50} = 21.1 $\mu M)$ significantly lowers hERG activity. When compared to the lead compound 1, para sulfoxide substitution on the left-hand-side phenyl ring, as shown in analogue 4 (IC₅₀ = 17.9 μ M), reduced hERG potency by 5fold. Analogues with basic side chains, 8 (IC₅₀ = 0.59 μ M), 9 (IC₅₀ = 5.2 μ M) and **23** (IC₅₀ = 7.83 μ M) were potent on the hERG channel. This is not unexpected since most ligands that block the hERG channel contain a basic nitrogen,18 which can get protonated at physiological pH thereby facilitating hERG binding through cation - π interactions between the protonated nitrogen and the π system of the aromatic residues in the channel cavity.¹⁸ For the aniline derivatives, 14 - 17, meta sulfonylation seems to enhance hERG activity (14, IC_{50} = 2.36 μ M). Strikingly, switching the sulfone in 14 with the sulfoxide diminishes hERG activity by about 8-fold as demonstrated in analogue 15 (IC₅₀ = 18.2 μ M). The regioisomer of 15, compound 16 (IC₅₀ = 32.3 μ M), which is para substituted showed further decrease in hERG inhibition. Interestingly, a positional change from meta to para (14 to 17) also led to a 11-fold reduction in hERG activity. We also hypothesized that the introduced saturated rings in 19 and 21 would reduce hERG binding by disrupting $\pi - \pi$ interactions between phenyl rings in our molecules and those embedded in the channel cavity.¹⁸ Interestingly, both analogues 19 (IC₅₀ = 28.8 μ M) and 21 (IC₅₀ = 25.9 µM) indeed displayed significantly weaker hERG inhibition activity.

Table 4 In vitro mammalian cytotoxicity and hERG inhibition profiling for selected analogues

Compound code	Structure	Cytotoxicity ^f CHO cells		_ hERG IC₌օ. սM (SD) ^ց
		IC ₅₀ , μM (SD)	SI ^h	
1	SO ₂ Me	> 234	> 69,000	3.61 (0.623)
28	SO ₂ Me	11.7 (0.89)	377	2.35 (0.46)
13		31.1 (0.2)	206	8.45 (0.8)

This journal is © The Royal Society of Chemistry 20xx

Journal Name

Compound code	Structure _	Cytotoxicity ^f CHO cells		
Compound code		IC ₅₀ , μM (SD)	SI ^h	
29	MeO ₂ S	181 (5.2)	457	4 (0.83)
30	MeOS	18.9 (1.5)	72	ND^e
32	Meos North Some	106 (21)	122	21.1 (2.8)
3	Meos Some	> 253	> 376	ND ^e
5	MeO ₂ S	> 243	> 249	ND ^e
4		> 243	> 874	17.9 (5.56)
8		4.08 (0.96)	30	0.59 (0.02)
23		9.41 (3.0)	10	7.83 (1.21)
14	MeO2S	123 (0.4)	449	2.36 (0.21)
15	MeOS N N N N N N N N N N N N N N N N N N N	ND ^e	ND^e	18.2 (8.45)
16	Meos	> 244	> 442	32.3 (1.54)
17	MeO ₂ S MeO ₂ S MeO ₂ S N N N N N Some	> 234	> 553	26.1 (6.30)

J. Name., 2013, 00, 1-3 | 9

0.1039/C8MD00382C Journal Name

				DOI: 10.10 J
Compound code	Structure	Cytotoxicity ^f CHO cells		
compound code	Structure	IC ₅₀ , μΜ (SD)	SI ^h	
18		277 (18)	592	7.84 (0.76)
21	MeO2S N N N N N N N N N N N N N N N N N N N	> 231	> 464	25.9 (1.83)
19		> 247	> 471	28.8 (3.64)
9	NEt ₂ F SO ₂ Me	9.44 (2.1)	11	5.2 (0.44)
12		12 (2.7)	12	ND ^e

^eND, not determined.

Published on 06 September 2018. Downloaded by Kaohsiung Medical University on 9/7/2018 9:06:14 AM

^fMean from n = 3 independent experiments; SD, standard deviation; Emetine was used as a reference drug (IC₅₀ = 0.033 ± 0.006 μM). ^hSI = selectivity index = IC₅₀ (CHO)/IC₅₀ (*P*. *f* NF54).

SI = Selectivity index = 1050 (CHO)/1050 (F.) inter-

 g Mean from n = 3 independent experiments; Verapamil was used as a positive control (IC₅₀ = 0.560 ± 0.0961 μ M).

| SOMe

Conclusion

In our efforts to improve solubility and counter hERG inhibition, we identified a regioisomer of lead **1**, compound **28** (tables 3 and 4) with potent *in vitro* antiplasmodial activity ($IC_{50} = 0.031 \mu$ M) and high selectivity (SI = 377) against the CHO cell line. However, hERG inhibition ($IC_{50} = 2.35 \mu$ M) and poor solubility (< 5 μ M) remain issues. Interestingly, the sulfoxide version, compound **13** (tables 2 and 4), exhibited excellent solubility (200 μ M) with reduced hERG inhibition ($IC_{50} = 8.45 \mu$ M). Additionally, we successfully managed to substantially reduce hERG activity in molecules **4**, **13**, **15** - **19**, **21**, **23** and **32** ($IC_{50} = 7.83 - 32.3 \mu$ M, table 4). Similarly, the introduced molecular features also resulted in analogues **3**, **6** - **10**, **12**, **13**, **15**, **16**, **20** - **24**, **26**, **27** and **32** (tables 1 - 3) with moderate to high solubility (60 - 200 μ M).

Experimental procedures

Chemistry

All reagents and chemicals used in all reactions were purchased from various commercial sources and used without further purification. All the solvents used in the reactions were anhydrous save for ethanol which was absolute (99.9%). Reactions were monitored by a combination of analytical thin layer chromatography (TLC) and liquid chromatography mass spectrometry (LC-MS). The TLC plates were sourced from Merck (TLC Silica gel 60 F_{254} aluminium-backed) and were developed in a 100 mL beaker covered with either a watch glass or aluminium foil. The plates were visualized under ultraviolet light (UV 254 and 366 nm). An Agilent LC-MS instrument with the following components was used for compound purity checks and reaction monitoring: Agilent 1260[®] Infinity Binary Pump, Agilent 1260[®] Infinity Diode Array Detector, Agilent 1290[®] Infinity Column Compartment, Agilent 1260[®] Infinity Autosampler, Agilent 6120[®] Quadrupole LC-MS, and Peak Scientific[®] Genius 1050 Nitrogen Generator. An X-bridge[®] (C18, 2.5 μ m, 3.0 mm (ID) x 50 mm length) column maintained at 35 °C or 40 °C was used. The chromatographic mobile phase was composed of 10 mM aqueous ammonium acetate (NH₄Ac) spiked with 0.4% acetic acid while the organic phase was composed of 10 mM NH₄Ac in methanol spiked with 0.4% acetic acid. The mass spectra were acquired using electrospray ionisation (ESI) and/or atmospheric pressure chemical ionization (APCI) in the positive ionisation mode unless otherwise stated.

Biotage grade silica gel was employed for column chromatographic purifications on the Biotage Isolera One Flash Chromatography system. Additionally, Analtech Uniplate preparative TLC (prep-TLC) plates (silica gel GF, 20 × 20 cm, 2000 microns) were used for prep-TLC purifications. The mobile phase solvents were AR grade and were used without further distillation. All yields reported are isolated and correspond to individual synthetic steps in a scheme.

¹H-NMR and ¹³C-NMR spectra were acquired on either Bruker AV 400 (¹H 400.0, ¹³C 101 MHz) or Varian Mercury 300 (¹H 300.1 MHz) spectrometers.

All final compounds were subjected to purity check experiments using LC-MS to ensure acceptable purity (\geq 95%). Melting points were measured using the Automatic Stuart Melting

Point Apparatus SMP40 and are uncorrected. Purification and characterization for some selected intermediates and analogues are reported in electronic supplementary information.

6-Chloroimidazo[1,2-*b*]*pyridazine* (**2***a*). Bromoacetaldehyde diethylacetal (12.0 mL, 78 mmol) was added to a suspension of 3-amino-6-chloropyridazine (**1a**) (5.00 g, 39 mmol) in absolute ethanol (99.9%) (78 mL) and deionized water (50 mL). A 48% aqueous solution of HBr (4.4 mL, 39 mmol) was added after which the suspension cleared. The reaction mixture was refluxed at 103 °C for 22 hours. The resulting brown solution was extracted with ethyl acetate (EtOAc) (100 mL × 3). The organic layers were combined after which the solvent was removed *in vacuo* to obtain **2a** as a brown solid which was used in the next step without further purification (LC-MS, ESI/APCI⁺: *m/z* [M + H]⁺ = 154.0, calculated exact mass = 153.0094, t_r = 1.5 min).

6-Chloro-3-iodoimidazo[1,2-b]pyridazine (3a). A reddish solution of 6-chloroimidazo[1,2-b]pyridazine (2a) (4.60 g, 30 mmol) in 30 mL of N,N-dimethylformamide (DMF) was purged with nitrogen for 30 minutes. NIS (7.41 g, 33 mmol) was added followed by the addition of another 45 mL of DMF. The reaction mixture was then left to magnetically stir at room temperature (21 °C) for 5 days (initially a suspension, the reaction mixture cleared after 2 days of stirring). DMF was then removed in vacuo. The resulting brown residue was taken up in 200 mL dichloromethane (DCM) and washed with deionized water (100 mL \times 3), a saturated aqueous solution of a mixture of sodium metabisulfite (Na₂S₂O₅) and sodium bisulfite (NaHSO₃) (100 mL \times 2). The solvent from the organic layer was removed in vacuo to afford 3a as a brown solid (6.82 g, 81%); ¹H-NMR δ_{H} (300 MHz; CDCl₃) 7.92 (d, J = 9.4 Hz, 1H), 7.88 (s, 1H), 7.14 (d, J = 9.4 Hz, 1H); LC-MS, ESI/APCI⁺: m/z [M + H]⁺ = 279.8, calculated exact mass = 278.9060, t_r = 3.0 min.

General procedure for synthesis of chloro-substituted intermediates (4a – g). A suspension of 6-chloro-3-iodoimidazo[1,2-*b*]pyridazine (3a) (1.0 eq), an appropriate boronic acid or boronic acid pinacol ester (1.1 eq), and Pd(PPh_3)₂Cl₂ (0.05 eq) in DMF (3 mL/mmol of 3a) was purged with nitrogen for 20 minutes. A 1 M aqueous solution of K_2CO_3 (1.05 eq) was then added after which the reaction mixture was heated to 80 °C and left to magnetically stir at that temperature for 3.5 – 46 hours. The reaction mixture was diluted with DCM, washed with deionized water (8 ×), saturated aqueous solutions of NaHCO₃ (3 ×), NH₄Cl (3 ×) and NaCl (1 ×). For intermediates 4e – g, containing the basic side chain, washing with aq NH₄Cl was avoided. After drying the organic layer (MgSO₄), the solvent was removed *in vacuo* and the resulting residue subjected to automated column chromatography on silica to give the chlorosubstituted derivatives in 26 – 77% yield.

6-Chloro-3-(4-(methylsulfinyl)phenyl)imidazo[1,2-

$$\begin{split} b] pyridazine ~~(\textbf{4a}). & \text{Purified by flash chromatography (0 - 3% CH_3OH/DCM). Yellow solid (0.587 g, 77%); 1H-NMR $$_H$ (300 MHz; CDCl_3) 8.24 (d, J = 8.2 Hz, 2H), 8.18 (s, 1H), 8.11 (d, J = 9.4 Hz, 1H), 7.82 (d, J = 8.0 Hz, 2H), 7.22 (d, J = 9.4 Hz, 1H) 2.81 (s, 3H); LC-MS, ESI/APCl^*: m/z [M + H]^+ = 292.0, calculated exact mass = 291.0233, t_r = 3.0 min. \end{split}$$

General procedure for synthesis of diarylated imidazopyridazines (1, 4 – 12 and 29 - 32). A suspension of the relevant chloro-substituted intermediate (4a – g) (1.0 eq), appropriate boronic acid or boronic acid pinacol ester (1.1 eq) and Pd(PPh₃)₂Cl₂ (0.05 eq) in DMF (3 mL/mmol of 4a – g) was purged with nitrogen for 20 minutes. A 1 M aqueous solution of K₂CO₃ (1.05 eq) was then added after which the reaction mixture was heated to 100 °C and magnetically stirred at that temperature for 4 – 21 hours. The reaction mixture was then diluted with DCM, washed with deionized water (8 ×), saturated aqueous solutions of NaHCO₃ (3 ×), NH₄Cl (3 ×) and NaCl (1 ×). For analogues **6** – **12**, with basic side chains, washing with NH₄Cl (aq) was avoided. The organic layer was dried (MgSO₄ for **1** and **6** - **12**; Na₂SO₄ for **4**, **5** and **29** – **32**) after which the solvent was removed *in vacuo* and the resulting crude mixture subjected to further purification procedures.

DOI: 10.1039/C8MD00382C

ARTICLE

6-(4-(Methylsulfinyl)phenyl)-3-(4-

(*methylsulfonyl*)*phenyl*)*imidazo*[1,2-*b*]*pyridazine* (4). Purified by prep-TLC (developed in 4% CH₃OH/DCM) and trituration in ethyl acetate. Yellow solid (0.0683 g, 57%), mp 236.6 – 239.0 °C (from ethyl acetate); R_f (CH₃OH : CH₂Cl₂ 4 : 96) 0.18; ¹H-NMR $\delta_{\rm H}$ (400 MHz; CDCl₃) 8.41 (d, J = 8.8 Hz, 2H), 8.27 (s, 1H), 8.24 (d, J = 9.5 Hz, 1H), 8.19 (d, J = 8.7 Hz, 2H), 8.13 (d, J = 8.8 Hz, 2H), 7.88 (d, J = 8.7 Hz, 2H), 7.70 (d, J = 9.5 Hz, 1H), 3.15 (s, 3H), 2.84 (s, 3H); ¹³C-NMR $\delta_{\rm C}$ (101 MHz; CDCl₃) 151.23, 148.38, 140.09, 139.43, 137.92, 134.52, 133.74, 128.08 (3C), 127.97 (2C), 127.05 (2C), 126.65, 124.49 (2C), 116.66, 44.58, 44.00; LC-MS, APCl⁺: m/z [M + H]⁺ = 412.0, calculated exact mass = 411.0711, purity: 98.8%, t_r = 3.4 min.

N-Ethyl-N-(3-fluoro-4-(3-(4-

(methylsulfinyl)phenyl)imidazo[1,2-b]pyridazin-6-

yl)benzyl)ethanamine (6). Purified by flash chromatography (0 – 4% CH₃OH/DCM) and trituration in diethyl ether. Yellow solid (0.128 g, 59%), mp 110.7 – 113.0 °C (from diethyl ether); R_f (CH₃OH : CH₂Cl₂ 6 : 94) 0.32; ¹H-NMR $\delta_{\rm H}$ (300 MHz; CDCl₃) 8.35 (d, J = 8.3 Hz, 2H), 8.20 (s, 1H), 8.11 (d, J = 9.0 Hz, 1H), 7.94 – 7.75 (m, 3H), 7.62 (d, J = 9.2 Hz, 1H), 7.46 – 7.34 (m, 2H), 3.76 (s, 2H), 2.81 (s, 3H), 2.70 (br s, 4H), 1.17 (t, J = 6.8 Hz, 6H); ¹³C-NMR $\delta_{\rm c}$ (100.6 MHz; CDCl₃) 162.1, 159.4, 148.8, 144.8, 134.1, 131.6, 130.5, 127.3 (4C), 125.8 (2C), 124.0 (3C), 119.0, 117.4, 56.6, 46.9 (2C), 44.0, 11.2 (2C); LC-MS, ESI/APCl⁺: m/z [M + H]⁺ = 437.1, calculated exact mass = 436.1733, purity: 99.6%, t_r = 2.4 min.

3-(3-(Methylsulfonyl)phenyl)-6-(4-

(methylsulfonyl)phenyl)imidazo[1,2-b]pyridazine (**29**). Purified by prep-TLC (developed in 4% CH₃OH/DCM). Yellow solid (0.0223 g, 20%), mp 264.0 – 267.5 °C (from 7% MeOH in DCM); R_f (CH₃OH : CH₂Cl₂ 4 : 96) 0.38; ¹H-NMR $\delta_{\rm H}$ (400 MHz; DMSO-d₆) 9.12 – 9.08 (m, 1H), 8.56 (s, 1H), 8.53 (d, *J* = 8.0 Hz, 1H), 8.50 (d, *J* = 8.4 Hz, 2H), 8.45 (d, *J* = 9.6 Hz, 1H), 8.13 (d, *J* = 8.5 Hz, 2H), 8.09 (d, *J* = 9.6 Hz, 1H), 8.00 – 7.96 (m, 1H), 7.86 (t, *J* = 7.9 Hz, 1H), 3.33 (m, 6H); ¹³C-NMR $\delta_{\rm C}$ (101 MHz; DMSO-d₆) 150.20, 142.63, 142.04, 140.26, 140.07, 135.36, 131.32, 130.51, 129.90, 128.49 (2C), 128.17 (2C), 127.33, 126.10, 126.38, 124.60, 116.99, 44.08, 43.90; LC-MS, APCI⁺: *m/z* [M + H]⁺ = 428.0, calculated exact mass = 427.0660, purity: 95.8%, t_r = 3.5 min.

General procedure for synthesis of aminated analogues (15, 16, 18 and 21). A suspension of 6-chloro-3-(4-(methylsulfinyl)phenyl)imidazo[1,2-*b*]pyridazine (4a) (1.0 eq), an appropriate amine (1.1 eq), $Pd_2(dba)_3$ (0.08 eq), XPhos (0.12 eq) and sodium *tert*-butoxide (2.0 eq) in anhydrous 1,4-dioxane (6.0 mL/mmol of 4a) in a sealed tube was flashed with nitrogen for 20 minutes. The reaction mixture was then heated to 120 °C and stirred for 6 - 20 hours. The solvent was removed *in vacuo* after which the resulting residue was taken up in DCM, washed with saturated aqueous solutions of NaHCO₃ (2 ×) and NaCl (2 ×). The organic layer was dried (MgSO₄) (Na₂SO₄ for 16), where after the solvent was removed *in vacuo* and obtained residue further purified.

N-(3-(Methylsulfinyl)phenyl)-3-(4-

(*methylsulfinyl)phenyl)imidazo*[1,2-b]pyridazin-6-amine (15). Purified by prep-TLC (developed twice in 6% CH₃OH/DCM). Brown solid (0.0175 g, 18%); R_f (CH₃OH : CH₂Cl₂ 6 : 94) 0.19; ¹H-NMR δ_H (400 MHz; DMSO-d₆) 9.81 (s, 1H), 8.28 (d, J = 8.5 Hz, 2H), 8.11 (s,

DOI: 10.1039/C8MD00382C

Journal Name

1H), 8.05 (s, 1H), 8.03 (d, J = 9.7 Hz, 1H), 7.85 (d, J = 8.5 Hz, 2H), 7.77 (dd, J = 8.1, 1.3 Hz, 1H), 7.57 (t, J = 7.9 Hz, 1H), 7.31 (d, J = 7.7 Hz, 1H), 7.02 (d, J = 9.7 Hz, 1H), 2.82 (s, 3H), 2.73 (s, 3H); ¹³C-NMR $\delta_{\rm C}$ (101 MHz; DMSO-d₆) 151.09, 147.73, 145.44, 141.43, 138.21, 132.23, 131.62, 130.32, 127.41 (2C), 127.34 (2C), 124.69 (2C), 121.14, 117.01, 113.89, 113.50, 43.83, 43.73; LC-MS, APCI⁺: m/z [M + H]⁺ = 411.1, calculated exact mass = 410.0871, purity: 98.9%, t_r = 3.8 min.

3-(4-(Methylsulfinyl)phenyl)-N-(pyrazin-2-yl)imidazo[1,2b]pyridazin-6-amine (**18**). Purified by prep-TLC (developed in 5% CH₃OH/DCM). Light yellow solid (0.0242 g, 29%); R_f (CH₃OH : CH₂Cl₂ 6 : 94) 0.3; ¹H-NMR δ_H (400 MHz; DMSO-d₆) 10.43 (s, 1H), 9.23 (d, J = 1.3 Hz, 1H), 8.38 (dd, J = 2.6, 1.5 Hz, 1H), 8.35 (d, J = 8.5 Hz, 2H), 8.26 (d, J = 2.6 Hz, 1H), 8.16 – 8.09 (m, 2H), 7.83 (d, J = 8.5 Hz, 2H), 7.43 (d, J = 9.7 Hz, 1H), 2.82 (s, 3H); ¹³C-NMR δ_C (101 MHz; DMSOd₆) 150.20, 149.95, 145.70, 142.75, 138.50, 137.91, 135.82, 132.81, 131.46, 127.50, 127.35 (2C), 127.09, 124.45 (2C), 113.48, 43.68; LC-MS, APCI⁺: m/z [M + H]⁺ = 351.1, calculated exact mass = 350.0950, purity: 96.5%, t_r = 3.7 min.

General procedure for synthesis of diarylated imidazopyridazines 3 and 28. A suspension of 6-chloro-3iodoimidazo[1,2-*b*]pyridazine (3a) (1.0 eq), an appropriate boronic acid (2.2 eq), and Pd(PPh_3)_2Cl₂ (0.10 eq) in anhydrous DMF (2.6 mL) was purged with nitrogen for 30 minutes. A 1 M aqueous solution of K₂CO₃ (2.1 eq) was then added. The reaction mixture was heated to 100 °C and left to magnetically stir at this temperature for 24 hours (28) and 15 hours (3). The reaction mixture was then diluted with DCM (60 mL), washed with deionized water (40 mL × 7), saturated aqueous solutions of NaHCO₃ (40 mL × 3), NH₄Cl (40 mL × 3), and NaCl (40 mL × 1). The organic layer was dried (Na₂SO₄), after which the solvent was removed *in vacuo*. The resulting crude mixture was subjected to prep-TLC purification to get the title compounds.

3,6-Bis(4-(methylsulfinyl)phenyl)imidazo[1,2-b]pyridazine

(3). Purified by prep-TLC (developed twice in 5% CH₃OH/DCM). Crystallized in diethyl ether. Yellow solid (0.0665 g, 39%); mp 180.4 – 182.3 °C (from diethyl ether); R_f (CH₃OH : CH₂Cl₂ 4 : 96) 0.21; ¹H-NMR δ_H (400 MHz; DMSO-d₆) 8.48 (d, J = 8.8 Hz, 2H), 8.45 (s, 1H), 8.39 (d, J = 9.6 Hz, 1H), 8.35 (d, J = 8.7 Hz, 2H), 7.99 (d, J = 9.6 Hz, 1H), 7.91 (d, J = 8.6 Hz, 2H), 7.89 (d, J = 8.8 Hz, 2H), 2.84 (s, 3H), 2.83 (s, 3H); ¹³C-NMR δ_c (101 MHz; DMSO-d₆) 151.04, 148.96, 145.89, 140.15, 137.75, 135.09, 131.10, 128.43 (2C), 127.23 (3C), 127.13, 124.91 (2C), 124.68 (2C), 116.98, 43.67 (2C); LC-MS, APCI⁺: m/z [M + H]⁺ = 396.1, calculated exact mass = 395.0762, purity: 98.5%, t_r = 3.7 min.

6-(3-(Methylsulfinyl)phenyl)imidazo[1,2-b]pyridazine (4h). A suspension of 6-chloroimidazo[1,2-b]pyridazine (2a) (1.50 g, 9.8 mmol), 3-methylsulfinylphenylboronic acid (1.78 g, 9.8 mmol), and Pd(PPh₃)₂Cl₂ (0.344 g, 0.49 mmol) in anhydrous DMF (30 mL) was purged with nitrogen for 30 minutes. A 1 M aqueous solution of K₂CO₃ (1.42 g, 10.29 mmol) was then added. The reaction mixture was heated to 100 °C and magnetically stirred at this temperature for 3.25 hours. The black reaction mixture was then diluted with DCM (150 mL), washed with deionized water (75 mL × 8), saturated aqueous solutions of NaHCO₃ (75 mL × 3), NH₄Cl (75 mL × 1) and NaCl (50 mL \times 5). After drying (MgSO₄) the organic layer, the solvent was removed in vacuo and the resulting crude mixture subjected to flash column chromatography (0 - 5% CH₃OH/DCM) to afford the phenylated intermediate, 4h which was further crystallized in diethyl ether to obtain a yellow solid (1.20 g, 48%); ¹H-NMR δ_{H} (400 MHz; CDCl₃) 8.36 (td, J = 1.8, 0.6 Hz, 1H), 8.25 (d, J = 9.6 Hz, 1H), 8.15 (dt, J = 7.7, 1.7 Hz, 1H), 8.10 (d, J = 1.3 Hz, 1H), 7.89 (d, J = 1.4

3-Bromo-6-(3-(methylsulfinyl)phenyl)imidazo[1,2-(4i). of 6-(3b]pyridazine А brown solution (methylsulfinyl)phenyl)imidazo[1,2-b]pyridazine (4h) (1.20 g, 4.7 mmol) in anhydrous DMF (4 mL) was purged with nitrogen for 30 minutes. NBS (0.912 g, 5.1 mmol) was added where after the reaction mixture was stirred at room temperature (23 °C) for 22 hours. The reaction mixture, containing a yellow precipitate at this point, was taken up in 100 mL DCM and washed with deionized water (75 mL × 8), saturated aqueous solutions of NaHCO₃ (75 mL × 2), NH₄Cl (75 mL × 2) and NaCl (75 mL × 1). The organic layer was dried (MgSO₄) after which the solvent was removed in vacuo. The obtained oily residue was crystallized and triturated with ethyl acetate for 30 minutes. The solid was filtered and washed with ethyl acetate to give **4i** as a yellow solid (0.939 g, 59%); ¹H-NMR $\delta_{\rm H}$ (300 MHz; CDCl₃) 8.44 - 8.35 (m, 2H), 8.29 - 8.23 (m, 1H), 7.93 (s, 1H), 7.90 – 7.73 (m, 3H), 2.85 (s, 3H); LC-MS, ESI^+ : $m/z [M + H]^+ =$ 336.0 & 338.0, calculated exact mass = 334.9728, t_r = 3.9 min.

3,6-Bis(3-(methylsulfinyl)phenyl)imidazo[1,2-b]pyridazine (13). suspension of 3-bromo-6-(3-(methylsulfinyl)phenyl)imidazo[1,2-b]pyridazine (4i) (0.100 g, 0.30 mmol), 3-methylsulfinylphenylboronic acid (0.0602 g, 0.33 mmol), and $Pd(PPh_3)_2Cl_2$ (0.0105 g, 0.015 mmol) in anhydrous DMF (2.0 mL) was purged with nitrogen for 30 minutes. A 1 M aqueous solution of K₂CO₃ (0.0435 g, 0.32 mmol) was then added. The reaction mixture was heated to 100 °C and magnetically stirred at this temperature for 20 hours. The reaction mixture was then diluted with DCM (60 mL), washed with deionized water (40 mL \times 6), saturated aqueous solutions of NaHCO₃ (40 mL × 3), NH₄Cl (40 mL × 3) and NaCl (40 mL \times 1). The organic layer was dried (MgSO₄), where after the solvent was removed in vacuo. The resulting crude mixture was subjected to prep-TLC (developed twice in 4% CH₃OH/DCM) to get 13 as an oil. Crystallization with diethyl ether gave 13 as a yellow solid (0.0481 g, 40%); R_f (CH₃OH : CH₂Cl₂ 4 : 96) 0.17; ¹H-NMR δ_{H} (400 MHz; DMSO-d₆) 8.80 (m, 1H), 8.48 – 8.42 (m, 2H), 8.42 – 8.35 (m, 2H), 8.33 (dt, J = 7.6, 1.6 Hz, 1H), 8.03 (d, J = 9.5 Hz, 1H), 7.94 - 7.88 (m, 1H), 7.82 (t, J = 7.7 Hz, 1H), 7.76 (t, J = 7.7 Hz, 1H), 7.73 – 7.69 (m, 1H), 2.88 (s, 3H), 2.86 (s, 3H) 13 C-NMR δ_{C} (101 MHz; DMSO-d₆) 150.75, 148.34, 147.67, 140.06, 136.44, 134.81, 130.65, 130.14, 129.89, 129.73, 128.72, 127.15, 127.08, 125.63, 123.28, 122.60, 121.19, 116.75, 43.97, 43.82; LC-MS, APCI⁺: $m/z [M + H]^{+} = 396.1$, calculated exact mass = 395.0762, purity: 99.8%, t_r = 3.8 min.

General procedure for synthesis of aminated analogues (24 suspension 27). А of 3-bromo-6-(3-(methylsulfinyl)phenyl)imidazo[1,2-b]pyridazine (4i) (1.0 eq), an appropriate amine (1.3 eq), Pd₂(dba)₃ (0.2 eq), BrettPhos (0.12 eq) and Cs₂CO₃ (2.0 eq) in anhydrous 1,4-dioxane (2 mL) in a sealed tube was flashed with nitrogen for 20 minutes. The reaction mixture was then heated to 120 °C and stirred for 5 - 39 hours. The solvent was removed in vacuo after which the resulting black residue was taken up in DCM (60 mL), washed with saturated aqueous solutions of NaHCO₃ (40 mL × 3) and NaCl (40 mL × 1). The organic layer was dried (Na2SO4), after which the solvent was removed in vacuo and the crude mixture further purified.

6-(3-(Methylsulfinyl)phenyl)-N-(4-

(methylsulfonyl)phenyl)imidazo[1,2-b]pyridazin-3-amine (24). Purified by prep-TLC (developed thrice in 2.5% CH₃OH/DCM). Crystallized in diethyl ether. Brown solid (0.0121 g, 9%); mp 105.0 – 107.2 °C (from diethyl ether); R_f (CH₃OH : CH₂Cl₂ 4 : 96) 0.23; ¹H-

Journal Name

NMR $\delta_{\rm H}$ (400 MHz; DMSO-d₆) 9.13 (s, 1H), 8.32 – 8.25 (m, 2H), 8.18 (d, *J* = 7.7 Hz, 1H), 7.91 – 7.80 (m, 3H), 7.78 – 7.68 (m, 3H), 6.99 (d, *J* = 8.4 Hz, 2H), 3.11 (s, 3H), 2.80 (s, 3H); ¹³C-NMR $\delta_{\rm C}$ (101 MHz; DMSO-d₆) 150.57, 149.91, 148.01, 136.65, 135.99, 130.46, 130.33, 129.64, 129.31 (2C), 127.66, 127.04, 126.95, 125.62, 122.61, 115.86, 114.13 (2C), 44.75, 43.63; LC-MS, APCI⁺: *m/z* [M + H]⁺ = 427.1, calculated exact mass = 426.0820, purity: 97.7%, t_r = 3.4 min.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We gratefully acknowledge the support of the University of Cape Town, South African Medical Research Council and the South African Research Chairs Initiative of the Department of Science and Technology administered through the South African National Research Foundation (K.C.). At Swiss TPH, we thank Christoph Fischli, Christin Gumpp, Sibylle Sax and Christian Scheurer for assistance in performing the antimalarial assays.

Notes and references

- (1) World Health Organization World Malaria Report; 2017.
- White, N. J. Antimalarial Drug Resistance. J. Clin. Invest. 2004, 113, 1084–1092.
- Sibley, C. H.; Hyde, J. E.; Sims, P. F.; Plowe, C. V.; Kublin, J.
 G.; Mberu, E. K.; Cowman, A. F.; Winstanley, P. A.; Watkins, W. M.; Nzila, A. M. Pyrimethamine–sulfadoxine Resistance in Plasmodium Falciparum: What Next? *Trends Parasitol.* 2001, *17*, 570–571.
- Terlouw, D. J.; Nahlen, B. L.; Courval, J. M.; Kariuki, S. K.; Rosenberg, O. S.; Oloo, A. J.; Kolczak, M. S.; Hawley, W. A.; Lal, A. A.; Kuile, F. O. T. Sulfadoxine-Pyrimethamine in Treatment of Malaria in Western Kenya: Increasing Resistance and Underdosing. *Antimicrob. Agents Chemother.* 2003, 47, 2929–2932.
- Warhurst, D. C. Resistance to Antifolates in Plasmodium Falciparum, the Causative Agent of Tropical Malaria. *Sci. Prog.* 2002, *85*, 89–111.
- Kusakabe, K.-I.; Yoshida, H.; Nozu, K.; Hashizume, H.; Tadano, G.; Sato, J.; Tamura, Y.; Mitsuoka, Y. Imidazo[1,2b]Pyridazine and Pyrazolo[1,5-a]Pyrimidine Derivatives and Their Use as Protein Kinase Inhibitors. WO2011/013729A1, 2011.
- Xu, Y.; Brenning, B. G.; Kultgen, S. G.; Liu, X.; Saunders, M.; Ho, K.-K. Preparation of Imidazo[1,2-b]Pyridazine and Pyrazolo[1,5- a]Pyrimidine Derivatives as Protein Kinase Inhibitors. US20120058997, 2012.
- Moran, D. B.; Powell, D. W.; Albright, J. D. Imidazo[1,2b]Pyridazines. US4569934A, 1986.
- (9) Matsumoto, S.; Miyamoto, N.; Hirayama, T.; Oki, H.; Okada, K.; Tawada, M.; Iwata, H.; Nakamura, K.; Yamasaki, S.; Miki, H.; Hori, A.; Imamura, S. Structure-Based Design, Synthesis, and Evaluation of Imidazo[1,2-b]Pyridazine and Imidazo[1,2-a]Pyridine Derivatives as Novel Dual c-Met and VEGFR2 Kinase Inhibitors. *Bioorg. Med. Chem.* **2013**, *21*,

7686-7698.

- (10) Osborne, S.; Chapman, T.; Large, J.; Bouloc, N.; Wallace, C. Fused Heterocyclic Compounds for Use in the Treatment of Malaria. WO 2011101640, 2011.
- (11) Lemercier, G.; Fernandez-Montalvan, A.; Shaw, J. P.; Kugelstadt, D.; Bomke, J.; Domostoj, M.; Schwarz, M. K.; Scheer, A.; Kappes, B.; Leroy, D. Identification and Characterization of Novel Small Molecules as Potent Inhibitors of the Plasmodial Calcium-Dependent Protein Kinase 1. *Biochemistry* **2009**, *48*, 6379–6389.
- (12) Sahu, N. K.; Kohli, D. V. Structural Insight for Imidazopyridazines as Malarial Kinase PfPK7 Inhibitors Using QSAR Techniques. *Med. Chem. (Los. Angeles).* 2012, *8*, 636–648.
- McNamara, C. W.; Lee, M. C. S.; Lim, C. S.; Lim, S. H.; Roland, J.; Nagle, A.; Simon, O.; Yeung, B. K. S.; Chatterjee, A. K.; McCormack, S. L.; Manary, M. J.; Zeeman, A.-M.; Dechering, K. J.; Kumar, T. R. S.; Henrich, P. P.; Gagaring, K.; Ibanez, M.; Kato, N.; Kuhen, K. L.; Fischli, C.; Rottmann, M.; Plouffe, D. M.; Bursulaya, B.; Meister, S.; Rameh, L.; Trappe, J.; Haasen, D.; Timmerman, M.; Sauerwein, R. W.; Suwanarusk, R.; Russell, B.; Renia, L.; Nosten, F.; Tully, D. C.; Kocken, C. H. M.; Glynne, R. J.; Bodenreider, C.; Fidock, D. A.; Diagana, T. T.; Winzeler, E. A. Targeting Plasmodium PI(4)K to Eliminate Malaria. *Nature* 2013, *504*, 248–253.
- (14) Le Manach, C.; Gonzàlez Cabrera, D.; Douelle, F.; Nchinda, A. T.; Younis, Y.; Taylor, D.; Wiesner, L.; White, K. L.; Ryan, E.; March, C.; Duffy, S.; Avery, V. M.; Waterson, D.; Witty, M. J.; Wittlin, S.; Charman, S. A.; Street, L. J.; Chibale, K. Medicinal Chemistry Optimization of Antiplasmodial Imidazopyridazine Hits from High Throughput Screening of a SoftFocus Kinase Library: Part 1. J. Med. Chem. 2014, 57, 2789–2798.
- Le Manach, C.; Paquet, T.; Gonzàlez Cabrera, D.; Younis, Y.; Taylor, D.; Wiesner, L.; Lawrence, N.; Schwager, S.; Waterson, D.; Witty, M. J.; Wittlin, S.; Street, L. J.; Chibale, K. Medicinal Chemistry Optimization of Antiplasmodial Imidazopyridazine Hits from High Throughput Screening of a SoftFocus Kinase Library: Part 2. J. Med. Chem. 2014, 57, 8839–8848.
- (16) Le Manach, C.; Paquet, T.; Brunschwig, C.; Njoroge, M.; Han, Z.; Gonzàlez Cabrera, D.; Bashyam, S.; Dhinakaran, R.; Taylor, D.; Reader, J.; Botha, M.; Churchyard, A.; Lauterbach, S.; Coetzer, T. L.; Birkholtz, L.-M.; Meister, S.; Winzeler, E. A.; Waterson, D.; Witty, M. J.; Wittlin, S.; Jiménez-Díaz, M.-B.; Santos Martínez, M.; Ferrer, S.; Angulo-Barturen, I.; Street, L. J.; Chibale, K. A Novel Pyrazolopyridine with in Vivo Activity in Plasmodium Berghei - and Plasmodium Falciparum - Infected Mouse Models from Structure–Activity Relationship Studies around the Core of Recently Identified Antimalarial Imidazopyridazines. J. Med. Chem. 2015, 58, 8713–8722.
- (17) Du, L.; Li, M.; You, Q. The Interactions Between HERG Potassium Channel and Blockers. *Curr. Top. Med. Chem.* **2009**, *9*, 330–338.
- Jamieson, C.; Moir, E. M.; Rankovic, Z.; Wishart, G.Medicinal Chemistry of HERG Optimizations: Highlights and

DOI: 10.1039/C8MD00382C

ARTICLE

Hang-Ups. J. Med. Chem. 2006, 49, 5029-5046.

- (19) Zhang, L.; Zhu, H.; Mathiowetz, A.; Gao, H. Deep Understanding of Structure–solubility Relationship for a Diverse Set of Organic Compounds Using Matched Molecular Pairs. *Bioorg. Med. Chem.* **2011**, *19*, 5763–5770.
- (20) Leach, A. G.; Jones, H. D.; Cosgrove, D. A.; Kenny, P. W.; Ruston, L.; MacFaul, P.; Wood, J. M.; Colclough, N.; Law, B. Matched Molecular Pairs as a Guide in the Optimization of Pharmaceutical Properties; a Study of Aqueous Solubility, Plasma Protein Binding and Oral Exposure. J. Med. Chem. 2006, 49, 6672–6682.
- (21) Forster, S.; Buckton, G.; Beezer, A. E. The Importance of Chain Length on the Wettability and Solubility of Organic Homologs. Int. J. Pharm. **1991**, 72, 29–34.
- (22) Wolfe, J. P.; Wagaw, S.; Buchwald, S. L. An Improved Catalyst System for Aromatic Carbon–Nitrogen Bond Formation: The Possible Involvement of Bis(Phosphine) Palladium Complexes as Key Intermediates. J. Am. Chem. Soc. 1996, 118, 7215–7216.

Graphical

- (23) Driver, M. S.; Hartwig, J. F. A Second-Generation Catalyst for Aryl Halide Amination: Mixed Secondary Amines from Aryl Halides and Primary Amines Catalyzed by (DPPF)PdCl2. J. Am. Chem. Soc. 1996, 118, 7217–7218.
- (24) Ishikawa, M.; Hashimoto, Y. Improvement in Aqueous Solubility in Small Molecule Drug Discovery Programs by Disruption of Molecular Planarity and Symmetry. *J. Med. Chem.* **2011**, *54*, 1539–1554.



abstract