

A Convergent Synthesis of (+)-Cryptophycin B, a Potent Antitumor Macrolide from *Nostoc* sp. Cyanobacteria


Arun K. Ghosh* and Alexander Bischoff

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607

arunghos@uic.edu

Received March 14, 2000

ABSTRACT



An efficient and highly stereoselective synthesis of cryptophycin B (2), a potent cytotoxic agent, is described. The ester-derived titanium-enolate-mediated *syn*-aldol reaction was employed to generate the stereocenters C(5) and C(6). The route is convergent and provides a convenient access to the synthesis of structural variants of cryptophycin B as well as members of its family.

Macrocyclic marine natural products continue to be a rich source for diverse antitumor agents with significant clinical potential.¹ Of particular interest, the cryptophycins, a group of depsipeptides isolated recently, have exhibited remarkably potent antitumor properties. The first cryptophycin (1) was isolated from terrestrial *Nostoc* sp. ATCC 53789 by researchers at Merck and was found to be very active against *Cryptococcus*, a fungus that often infects persons with immunodeficiencies.² In 1994, Moore et al. isolated a host of cyclic depsipeptides including cryptophycin A (1) and B (2) from *Nostoc* sp. GSV 224 and established their absolute stereochemistry (Figure 1).³ Cryptophycins A and B exhibited cytotoxic IC₅₀ values of 5 and 7 pg/mL, respectively, against KB cells. In addition, the compounds were equally effective against drug-sensitive and drug-resistant tumor cells.^{3,4} Arenastatin A (3), another member of the cryptophycin family, was isolated from the Okinawan marine sponge

Dysidea arenaria.^{5,6} It has displayed a cytotoxic IC₅₀ value of 5 pg/mL against KB cells.⁶

Cryptophycin A and other members of the family have been shown to promote depletion of microtubules.⁴ Furthermore, a number of these compounds effectively inhibited *in vitro* tubulin polymerization.⁷ The significant clinical potential of the cryptophycins and their relatively low natural abundance has attracted immense interest in their synthesis and structural modification. Several total syntheses and synthetic approaches to cryptophycins and arenastatin A have been described in recent years.^{8,9} As part of our interest in the structure–function studies of cryptophycins, we sought a flexible, enantioselective synthesis of cryptophycin B. Herein we report a convergent and stereocontrolled total synthesis of cryptophycin B.

As outlined in Figure 1, we planned to assemble cryptophycin B in a convergent manner from the protected

(1) Harris, C. R.; Danishefsky, S. J. *J. Org. Chem.* **1999**, *64*, 8434.
(2) (a) Schwarz, R. E.; Hirsch, C. F.; Sesin, D. F.; Flor, J. E.; Chartrain, M.; Fromtling, R. E.; Harris, G. H.; Salvatore, M. J.; Liesch, J. M.; Yudin, K. *J. Ind. Microbiol.* **1990**, *5*, 113. (b) Schwarz, R. E.; Hirsch, C. F.; Sesin, D. F. U.S. Patent 4,946,835, Aug 7, 1990.
(3) Trimurtulu, G.; Ohtani, I.; Patterson, G. M. L.; Moore, R. E.; Corbett, T. H.; Valeriote, F. A.; Demchik, L. *J. Am. Chem. Soc.* **1994**, *116*, 4729.

(4) Smith, C. D.; Zhang, X.; Mooberry, S. L.; Patterson, G. M. L.; Moore, R. E. *Cancer Res.* **1994**, *54*, 3779.
(5) Kobayashi, M.; Aoki, S.; Ohayabu, N.; Kurosu, M.; Wang, W.; Kitagawa, I. *Tetrahedron Lett.* **1994**, *35*, 7969.
(6) Koiso, Y.; Morita, K.; Kobayashi, M.; Wang, W.; Ohayabu, N.; Iwasaki, S. *Chem.-Biol. Interact.* **1996**, *102*, 183.
(7) Kerksiek, M. R.; Mejillano, R. E.; Schwartz, R. E.; Georg, G. I.; Himes, R. H. *FEBS Lett.* **1995**, *377*, 59.





Figure 1.

octadienoic acid **4**, D-tyrosine ester **5**, and protected β -amino acid derivative **6**. The fragments were planned to be connected by Yamaguchi esterification and cycloamidation reactions. Introduction of the sensitive epoxide functionality was planned at the final stage of the synthesis. The key elements of our synthesis involved an ester-derived titanium-enolate-mediated *syn*-aldol reaction to set the stereocenters at C(5) and C(6) of fragment **4**^{10a,b}. Thus, acylation of *trans*-4-phenyl-3-butenic acid and (+)-(1*R*,2*S*)-1-(*N*-tosylamino)-2-indanol with DCC and DMAP afforded the ester **7** in 98% yield.^{10c} As shown in Scheme 1, exposure of **7** to TiCl₄ and *N,N*-diisopropylethylamine in CH₂Cl₂ at 23 °C generated the corresponding titanium enolate. Subsequent reaction of the enolate with (benzyloxy)propionaldehyde at -78 °C furnished the aldol **8** as a single diastereomer in 98% yield after silica gel chromatography. Reduction of **8** by lithium aluminum hydride in THF at 0 °C for 1 h afforded the diol in 92% yield. The primary hydroxyl group of the



^a (a) TiCl₄, *i*Pr₂NEt, 0 to 23 °C, 1 h then BnO(CH₂)₂CHO, -78 °C, 20 min (98%); (b) LAH, THF, 0 °C, 1 h (92%); (c) PhLi, THF, -78 °C, 30 min, TsCl, -20 °C, 30 min, then LAH, 0 °C, 20 min (96%); (d) TIPSOTf, 2,6-lutidine (99%); (e) K₂CO₃, BBr₃, CH₂Cl₂, 0 °C, (83%); (f) PCC, MS 4 Å, 23 °C, 10 min (98%); (g) NaH, triethylphosphonoacetate, THF, 0 °C, 30 min (92%); (h) aq LiOH, EtOH (1:1), 23 °C, 2 h (94%); (i) **5**, DCC, DMAP, 23 °C, 12 h (79%); (j) TBAF, THF, 23 °C (99%).

resulting diol was selectively converted to the corresponding methyl group in an one-pot, two-step sequence. Thus, treatment of the diol with phenyllithium and *p*-toluenesulfonyl chloride at -78 °C to -20 °C followed by reduction of the resulting tosylate with lithium aluminum hydride provided the alcohol **9** in 96% yield.

Protection of the secondary alcohol using triisopropylsilyl triflate and 2,6-lutidine in CH₂Cl₂ gave **10** in 99% yield.¹¹ Selective removal of the benzyl protecting group was effectively achieved by treatment of **10** with boron tribromide in CH₂Cl₂ in the presence of potassium carbonate to provide the corresponding alcohol in 83% yield.¹² Initial attempts with other conditions such as TMSCl/*n*Bu₄N⁺I⁻, FeCl₃, or Na/NH₃ were unsuccessful.¹³ Oxidation of the resulting alcohol with PCC in CH₂Cl₂ followed by Horner–Emmons olefination of the aldehyde with sodium hydride and triethyl phosphonoacetate provided the α,β -unsaturated ester **11** in 92% yield. Ester hydrolysis with aqueous lithium hydroxide in a mixture (1:1) of EtOH and H₂O gave the octadienoic acid fragment **4** in 94% yield. Coupling of this acid with

(8) (a) Norman, B. H.; Hemscheidt, T.; Schultz, R. M.; Andis, S. L. *J. Org. Chem.* **1998**, *63*, 5288. (b) Gardinier, K. M.; Leahy, J. W. *J. Org. Chem.* **1997**, *62*, 7098. (c) Salamonczyk, G. M.; Han, K.; Guo, Z.-W.; Sih, C. J. *J. Org. Chem.* **1996**, *61*, 6893. (d) Rej, R.; Nguyen, D.; Go, B.; Fortin, S.; Lavallée, J.-F. *J. Org. Chem.* **1996**, *61*, 6289. (e) Barrow, R. A.; Hemscheidt, T.; Liang, J.; Paik, S.; Moore, R. E.; Tius, M. A. *J. Am. Chem. Soc.* **1995**, *117*, 2479. (f) Dhokte, U. P.; Khau, V. V.; Hutchison, D. R.; Martinelli, M. J. *Tetrahedron Lett.* **1998**, *39*, 8771. (g) White, J. D.; Hong, J.; Robarge, L. A. *J. Org. Chem.* **1999**, *64*, 6206. (h) Kobayashi, M.; Wang, W.; Ohyabu, N.; Kurosu, M.; Kitagawa, I. *Chem. Pharm. Bull.* **1995**, *43*, 1598. (i) Kobayashi, M.; Wang, W.; Ohyabu, N.; Kurosu, M.; Kitagawa, I. *Chem. Pharm. Bull.* **1994**, *42*, 2394.

(9) (a) Liang, L.; Hoard, D. W.; Khau, V. V.; Martinelli, M. J.; Moher, E. D.; Moore, R. E.; Tius, M. A. *J. Org. Chem.* **1999**, *64*, 1459. (b) Furuyama, M.; Shimizu, I. *Tetrahedron: Asymmetry* **1998**, *9*, 1351. (c) Ali, S. M.; Georg, G. I. *Tetrahedron Lett.* **1997**, *38*, 1703.

(10) (a) Ghosh, A. K.; Fidanze, S.; Senanayake, C. H. *Synthesis* **1998**, 937. (b) Ghosh, A. K.; Fidanze, S.; Onishi, M.; Hussain, K. A. *Tetrahedron Lett.* **1997**, *38*, 7171. (c) Ghosh, A. K.; Onishi, M. *J. Am. Chem. Soc.* **1996**, *118*, 2527.


(11) Rücker C. *Chem. Rev.* **1995**, *95*, 1009 and references cited therein.

(12) Treatment of **10** with boron tribromide in CH₂Cl₂ afforded the corresponding alcohol in only 32% yield. The major side product was the desilylation product.

(13) (a) Vedejs, E.; Buchanan, R. A.; Watanabe, Y. *J. Am. Chem. Soc.* **1989**, *111*, 8430. (b) Oikawa, Y.; Tanaka, T.; Yonemitsu, O. *Tetrahedron Lett.* **1986**, *27*, 3647. (c) Park, M. H.; Takeda, R.; Nakanishi, K. *Tetrahedron Lett.* **1987**, *28*, 3823.

O-methyl-*D*-tyrosine *tert*-butyl ester¹⁴ in the presence of DCC and DMAP in CH₂Cl₂ resulted in the amide **12** in 79% yield.¹⁵ Exposure of **12** with *n*Bu₄N⁺F⁻ in the presence of a trace of water in THF furnished the alcohol **13** in near quantitative yield.¹⁶

The synthesis of β -amino acid derivative **6** was carried out in two steps as shown in Scheme 2. Esterification of



^a (a) DCC, benzyl (*S*)-(-)-2-hydroxyisocaproate, DMAP, CH₂Cl₂, 23 °C, 12 h (91%); (b) H₂, 10% Pd-C, EtOAc, 5 h (95%); (c) 2,4,6-(Cl₃)PhCOCl, *i*Pr₂NEt, DMAP, alcohol **13**, 23 °C; (d) CF₃CO₂H, 23 °C, 2 h; (e) dimethyldioxirane, CH₂Cl₂, -30 to 23 °C, 12 h (87%, 3:1 mixture).

optically active *N*-Boc-3-amino-2-methylpropionic acid **14**¹⁷ and benzyl (*S*)-(-)-2-hydroxyisocaproate¹⁸ with dicyclohexylcarbodiimide and DMAP in CH₂Cl₂ at 23 °C for 12 h afforded the diester **15** in 91% yield. Catalytic hydrogenation of diester **15** over 10% Pd/C in ethyl acetate under a hydrogen-filled balloon for 5 h furnished the acid **6** in 95% yield. The acid **6** was subjected to esterification by reaction with 2,4,6-trichlorobenzoyl chloride, *N,N*-diisopropylethylamine, and alcohol **13** in the presence of DMAP under Yamaguchi conditions to provide diester **17** in 84% yield.¹⁹ Exposure of **17** to trifluoroacetic acid at 23 °C for 2 h resulted in the removal of both the *tert*-butyl ester and the Boc-

protecting group to furnish the requisite amino acid precursor for the macrocyclization. Thus, subjection of the resulting amino acid to Yamaguchi conditions with 2,4,6-trichlorobenzoyl chloride, *N,N*-diisopropylethylamine, and DMAP at 23 °C for 12 h furnished the cycloamide **18** ($[\alpha]_D^{23} +36.2$, *c* 0.72, MeOH; lit.³ $[\alpha]_D +36.7$, *c* 1.93, MeOH) in 74% yield (from **17**). Epoxidation of **18** with dimethyldioxirane in CH₂-Cl₂ at -30 °C to 23 °C for 12 h provided a 3:1 mixture of epoxides in 87% yield. The major epoxide was separated by reverse-phase HPLC on a C-18 column²⁰ (eluent, 3:1 MeOH/H₂O) to provide the synthetic cryptophycin B (**2**, $[\alpha]_D^{23} +20.6$, *c* 0.24, MeOH; lit.³ $[\alpha]_D +20.4$, *c* 0.54, MeOH). Spectral data (500 MHz ¹H NMR, IR, and ¹³C NMR) for the synthetic cryptophycin B are in full agreement with that reported by Moore et al. for the natural cryptophycin B.³

In summary, a stereocontrolled synthesis of cryptophycin B has been accomplished in 14 steps from **7** in 22% overall yield. The present synthetic route is easily amenable to the synthesis of other members of the cryptophycin family. Further studies of cryptophycins are ongoing in our laboratory.

Acknowledgment. Financial support for this work was provided by the National Institutes of Health (GM 55600). We thank Mr. Alexander Ho for preliminary experimental assistance.

Supporting Information Available: ¹H NMR and ¹³C NMR spectra for compounds **2** and **4–18**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL000058I

(14) *O*-Methyl-*D*-tyrosine *tert*-butyl ester was prepared in a three-step sequence involving (1) treatment of commercially available *N*-Cbz-*D*-tyrosine with Me₂SO₄ and NaOH in aqueous EtOH at reflux; (2) reaction of the resulting *O*-methyltyrosine with 2,4,6-trichlorobenzoyl chloride, *i*Pr₂NEt, DMAP, and *t*BuOH; and (3) removal of Cbz group by hydrogenation over 10% Pd-C in MeOH (55% overall).

(15) Hassner A.; Alexanian V. *Tetrahedron Lett.* **1978**, *46*, 4475.

(16) Reaction rate was accelerated in the presence of a trace amount of water.

(17) Amino acid **14** was prepared by tosylation of commercially available methyl (*R*)-(-)-3-hydroxy-2-methylpropionate with TsCl and NEt₃ in CH₂-Cl₂ for 1 h. Displacement of tosylate with NaN₃ in DMSO at 23 °C for 6 h. Hydrogenation of the resulting azide over 10% Pd/C in the presence of Boc₂O in EtOAc followed by saponification with aqueous LiOH at 23 °C for 30 min (90% overall).

(18) Benzyl (*S*)-(-)-2-hydroxyisocaproate was prepared from commercially available (*S*)-(-)-2-hydroxyisocaproate by treatment with CsCO₃ in MeOH/H₂O (6:1) for 30 min followed by reaction with BnBr in DMF at 0 °C (91%).

(19) Inanaga J.; Hirata K.; Saeki H.; Katsuki T.; Yamaguchi M. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 1989.

(20) The major epoxide was cleanly separated by an isocratic reverse phase HPLC (see ref 8g for similar separation) on a YMC-Pack ODS-AQ 5S 120 Å column (4.6 × 250 mm) with flow rate 1 mL/min and UV detection at 254 nm. Retention times for **2**: 31.58 min; and diastereomer: 37.20 min.