

Enantioselective Michael Reaction of Malonates and α,β -Unsaturated Aldehydes Using a *trans*-4-Hydroxyproline Derived Organocatalyst

Hisayuki Sato, Fumiaki Nagashima, and Takeshi Oriyama*

Department of Chemistry, Faculty of Science, Ibaraki University, 2-1-1 Bunkyo, Mito 310-8512

(Received January 20, 2010; CL-100064; E-mail: tor@mx.ibaraki.ac.jp)

Asymmetric Michael addition of malonates to various α,β -unsaturated aldehydes using an organocatalyst derived from *trans*-4-hydroxyproline in MeOH proceeds smoothly to afford the corresponding Michael adducts in high yields with high to excellent enantioselectivities.

Organocatalytic asymmetric Michael addition¹ has widely been used for the stereocontrolled formation of carbon–carbon and carbon–heteroatom bonds. α,β -Unsaturated compounds are known as versatile Michael acceptors that provide important synthetic intermediates with various nucleophiles via Michael reaction. The organocatalytic asymmetric Michael addition of malonates to α,β -unsaturated aldehydes has been reported in recent years.² For example, Jørgensen et al. demonstrated that *O*-TMS diarylprolinol derived from (*S*)-proline was an effective organocatalyst.³ Zlotin et al. have reported that *O*-TMS diphenylprolinol modified with an ionic liquid moiety can be used four times without any decrease in activity and enantioselectivity.⁴ However, these reactions are very slow (1–4 d). Ma et al. have mentioned asymmetric Michael reaction catalyzed by *O*-TMS-protected diphenylprolinol and acetic acid in water.⁵ In this case, the reaction reached completion in less than 24 h. However, more than 20 mol % of additive was needed for the reaction to be completed in reasonable time with high enantioselectivity. Therefore, the design and synthesis of more promising diarylprolinol silyl ethers⁶ is a significant requirement for the organocatalytic asymmetric Michael reaction.

On the other hand, we reported the solvent-free organocatalytic asymmetric Michael addition of thiols to α,β -unsaturated aldehydes using an organocatalyst derived from *trans*-4-hydroxyproline in 2007.⁷ This reaction proceeded smoothly without any organic solvent to give the corresponding chiral sulfides in almost enantiomerically pure form (up to 99% ee). We speculated that organocatalysts derived from *trans*-4-hydroxyproline may be extended to the enantioselective asymmetric Michael addition of malonates to α,β -unsaturated aldehydes. Herein, we disclose our fruitful results of these investigations.

First, we examined the reaction of cinnamaldehyde (0.45 mmol) with diethyl malonate (0.3 mmol) in MeOH as a model combination to optimize the organocatalyst (Figure 1 and Table 1). Organocatalyst **1** was found to be the most effective for this reaction (Entry 1). While a higher enantioselectivity was obtained with organocatalyst **3**, the chemical yield of Michael adduct was much lower compared to organocatalyst **1** (Entry 3). In the case of using organocatalyst **3**, a longer reaction time was required (Entry 4). In the presence of organocatalyst **4**, which has a free hydroxy group, yield and enantioselectivity of the Michael adduct were reduced (Entry 5). The reaction catalyzed by organocatalyst **5** also afforded the corresponding Michael




Figure 1. Organocatalysts examined in this study.

Table 1. Catalyst screening for the asymmetric Michael addition of diethyl malonate to cinnamaldehyde^a

$\text{EtO}_2\text{C}-\text{CH}_2-\text{CO}_2\text{Et} + \text{Ph}-\text{CH}=\text{CH}-\text{CHO} \xrightarrow[\text{MeOH / r.t. / 24 h}]{10 \text{ mol \% organocatalyst}} \text{EtO}_2\text{C}-\text{CH}(\text{Ph})-\text{CH}(\text{Ph})-\text{CHO}$			
Entry	Organocatalyst	Yield/% ^b	Ee/% ^c
1	1	94	97
2	2	90	94
3	3	62	92
4 ^d	3	80	94
5	4	53	62
6	5	75	98
7 ^e	1	98	97
8 ^f	1	74	96
9 ^g	1	67	97

^aUnless otherwise specified, the reactions were performed using cinnamaldehyde (0.45 mmol), diethyl malonate (0.3 mmol), and organocatalyst (0.03 mmol) in MeOH (0.5 mL).

^bIsolated yields. ^cEe was determined by HPLC analysis using a chiral column after oxidation to the corresponding methyl ester.³ ^dThe reaction was carried out for 48 h. ^eThe reaction was performed in MeOH (0.3 mL) for 3 h. ^fThe reaction was carried out with 5 mol % catalyst for 12 h. ^gThe reaction was carried out at 0 °C for 3 h.

adducts with 98% ee (Entry 6). When the reaction was carried out at 1.0 M, the chemical yield of the Michael adduct increased and the reaction was completed in 3 h (Entry 7). Decreasing the catalyst loading from 10 to 5 mol % resulted in a lower product yield (Entry 8). A reaction carried out at 0 °C did not improve the enantioselectivity (Entry 9).

Next, we investigated the effect of solvents in the presence of organocatalyst **1** (Table 2). As shown in Table 2, polar protic solvents, like MeOH and EtOH were effective and led to high enantioselectivities (Entries 1 and 2), whereas DMF, CH₃CN, CH₂Cl₂, and hexane were not effective from the viewpoint of

Table 2. Effect of solvents^a

Entry	Solvent	Yield/% ^b	Ee/% ^c
1	MeOH	98	97
2	EtOH	70	97
3	DMF	8	96
4	CH ₃ CN	9	90
5	CH ₂ Cl ₂	4	95
6	hexane	8	94
7	neat	46	97

^aUnless otherwise specified, the reactions were performed using cinnamaldehyde (0.45 mmol), diethyl malonate (0.3 mmol), and organocatalyst **1** (0.03 mmol) in MeOH (0.3 mL).

^bIsolated yields. ^cEe was determined by HPLC analysis using a chiral column after oxidation to the corresponding methyl ester.³

chemical yields (Entries 3–6). A similar solvent effect was observed in the Jørgensen's paper.³ Interestingly, the desired Michael adduct was obtained in 46% yield with 97% ee under solvent-free conditions (Entry 7).

A variety of reactions were conducted under optimized conditions in the presence of 10 mol % of organocatalyst **1** in MeOH at room temperature to establish the scope and limitations of the present Michael reaction (Table 3).⁸ Using dimethyl and dibenzyl malonates as Michael donors with cinnamaldehyde caused reactions to occur readily (Entries 1 and 3). The reaction of aromatic α,β -unsaturated aldehydes substituted with electron-donating (OMe) and electron-withdrawing groups (NO₂) afforded the corresponding Michael adduct in good to high yields with high enantioselectivities (Entries 4–11). It is notable that ortho substituted aromatic α,β -unsaturated aldehydes were found to be more effective in the enantioselectivity (Entries 5 vs. 7 and 9 vs. 11). This result might arise from steric hindrance between enals and two silyloxy groups of the organocatalyst molecule. Furthermore, heteroaromatic and aliphatic α,β -unsaturated aldehydes performed well in the presence of 10 mol % of organocatalyst **1**. Unfortunately, a longer reaction time was required for these aldehydes and the Michael adducts were obtained in moderate yields with good enantioselectivities (Entries 12 and 13).

In summary, we have developed a highly enantioselective Michael addition of malonates to various α,β -unsaturated aldehydes in the presence of an organocatalyst derived from *trans*-4-hydroxyproline in MeOH. In contrast to known methods, the reaction proceeds smoothly in a shorter reaction time to afford the corresponding Michael adducts with excellent enantioselectivities (up to 98% ee). Further studies on the development of environmentally benign reactions using organocatalysts derived from *trans*-4-hydroxyproline are currently in progress in our laboratory.

This work was partially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Table 3. Asymmetric Michael addition of malonates to various enals^a

Entry	R ¹	R ²	Time/h	Yield/% ^b	Ee/% ^c
1	Me	Ph	3	97	94
2	Et	Ph	3	98	97
3	Bn	Ph	3	87	94
4	Me	<i>o</i> -MeOC ₆ H ₄	6	86	87
5	Et	<i>o</i> -MeOC ₆ H ₄	6	73	95
6	Bn	<i>o</i> -MeOC ₆ H ₄	6	85	92
7	Et	<i>p</i> -MeOC ₆ H ₄	6	87	91
8	Me	<i>o</i> -NO ₂ C ₆ H ₄	6	88	95
9	Et	<i>o</i> -NO ₂ C ₆ H ₄	6	87	98
10	Bn	<i>o</i> -NO ₂ C ₆ H ₄	6	77	95
11	Et	<i>p</i> -NO ₂ C ₆ H ₄	6	76	92
12	Me	2-Furyl	13	30	82
13	Bn	Me	48	42	79

^aAll reactions were performed at a 0.3 mmol scale. ^bIsolated yields. ^cEe was determined by HPLC analysis using a chiral column after oxidation to the corresponding methyl ester.³

References and Notes

- For recent reviews, see: a) T. Ooi, T. Miki, M. Taniguchi, M. Shiraishi, M. Takeuchi, K. Maruoka, *Angew. Chem., Int. Ed.* **2003**, *42*, 3796. b) Y. Yamamoto, N. Momiyama, H. Yamamoto, *J. Am. Chem. Soc.* **2004**, *126*, 5962. c) M. Marigo, J. Franzén, T. B. Poulsen, W. Zhuang, K. A. Jørgensen, *J. Am. Chem. Soc.* **2005**, *127*, 6964. d) F. Wu, R. Hong, J. Khan, X. Liu, L. Deng, *Angew. Chem., Int. Ed.* **2006**, *45*, 4301. e) Y. K. Chen, M. Yoshida, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2006**, *128*, 9328. f) D. Almaši, D. A. Alonso, C. Nájera, *Tetrahedron: Asymmetry* **2007**, *18*, 299. g) H. Pellissier, *Tetrahedron* **2007**, *63*, 9267. h) S. Mukherjee, J. W. Yang, S. Hoffmann, B. List, *Chem. Rev.* **2007**, *107*, 5471. i) D. Enders, C. Grondal, M. R. M. Hüttel, *Angew. Chem., Int. Ed.* **2007**, *46*, 1570. j) G. Guillena, D. J. Ramón, M. Yus, *Tetrahedron: Asymmetry* **2007**, *18*, 693. k) D. Enders, K. Lüttgen, A. A. Narine, *Synthesis* **2007**, 959. l) H. Pellissier, *Tetrahedron* **2007**, *63*, 9267. m) S. B. Tsogoeva, *Eur. J. Org. Chem.* **2007**, 1701. n) J. L. Vicario, D. Badia, L. Carrillo, *Synthesis* **2007**, 2065. o) P. Melchiorre, M. Marigo, A. Carlone, G. Bartoli, *Angew. Chem., Int. Ed.* **2008**, *47*, 6138. p) B. Tan, X. Zeng, Y. Lu, P. J. Chua, G. Zhong, *Org. Lett.* **2009**, *11*, 1927. q) Y. Liu, B. Sun, B. Wang, M. Wakem, L. Deng, *J. Am. Chem. Soc.* **2009**, *131*, 418. r) Y.-F. Sheng, Q. Gu, A.-J. Zhang, S.-L. You, *J. Org. Chem.* **2009**, *74*, 6899. s) S. Chandrasekhar, K. Mallikarjuna, G. Pavankumarreddy, K. V. Rao, B. Jagadeesh, *Chem. Commun.* **2009**, 4985. t) S. Syu, T.-T. Kao, W. Lin, *Tetrahedron* **2010**, *66*, 891.
- a) C. Palomo, A. Landa, A. Mielgo, M. Oiarbide, A. Puente, S. Vera, *Angew. Chem., Int. Ed.* **2007**, *46*, 8431. b) G.-L. Zhao, A. Córdova, *Tetrahedron Lett.* **2007**, *48*, 5976. c) Y. Wang, P. Li, X. Liang, J. Ye, *Adv. Synth. Catal.* **2008**, *350*, 1383. d) I. Fleischer, A. Pfaltz, *Chem.—Eur. J.* **2010**, *16*, 95.

- 3 S. Brandau, A. Landa, J. Franzén, M. Marigo, K. A. Jørgensen, *Angew. Chem., Int. Ed.* **2006**, *45*, 4305.
- 4 O. V. Maltsev, A. S. Kucherenko, S. G. Zlotin, *Eur. J. Org. Chem.* **2009**, 5134.
- 5 A. Ma, S. Zhu, D. Ma, *Tetrahedron Lett.* **2008**, *49*, 3075.
- 6 For recent reviews on diarylprolinol silyl ether-catalyzed asymmetric reactions, see: a) Y. Hayashi, H. Gotoh, T. Hayashi, M. Shoji, *Angew. Chem., Int. Ed.* **2005**, *44*, 4212. b) M. Marigo, T. Schulte, J. Franzén, K. A. Jørgensen, *J. Am. Chem. Soc.* **2005**, *127*, 15710. c) T. Govender, L. Hojabri, F. M. Moghaddam, P. I. Arvidsson, *Tetrahedron: Asymmetry* **2006**, *17*, 1763. d) Y. Hayashi, T. Okano, T. Itoh, T. Urushima, H. Ishikawa, T. Uchimaru, *Angew. Chem., Int. Ed.* **2008**, *47*, 9053. e) A. Carlone, M. Marigo, C. North, A. Landa, K. A. Jørgensen, *Chem. Commun.* **2006**, 4928. f) A. Mielgo, C. Palomo, *Chem. Asian J.* **2008**, *3*, 922. g) B.-C. Hong, R. Y. Nimje, A. A. Sadani, J.-H. Liao, *Org. Lett.* **2008**, *10*, 2345. h) S. Belot, A. Massaro, A. Tenti, A. Mordini, A. Alexakis, *Org. Lett.* **2008**, *10*, 4557. i) Q. Zhu, Y. Lu, *Org. Lett.* **2008**, *10*, 4803. j) Y. Hayashi, M. Toyoshima, H. Gotoh, H. Ishikawa, *Org. Lett.* **2009**, *11*, 45.
- k) G. Luo, S. Zhang, W. Duan, W. Wang, *Tetrahedron Lett.* **2009**, *50*, 2946. l) M. Rueping, A. Kuenkel, F. Tato, J. W. Bats, *Angew. Chem., Int. Ed.* **2009**, *48*, 3699. m) A. Landa, M. Maestro, C. Palomo, A. Puente, S. Vera, M. Oiarbide, C. Palomo, *Chem.—Eur. J.* **2009**, *15*, 1562. n) H. Ishikawa, T. Suzuki, Y. Hayashi, *Angew. Chem., Int. Ed.* **2009**, *48*, 1304.
- 7 T. Ishino, T. Oriyama, *Chem. Lett.* **2007**, *36*, 550.
- 8 A general experimental procedure is as follows: Diethyl malonate (45.5 μL , 0.3 mmol) was added to a solution of organocatalyst **1** (13.7 mg, 0.03 mmol) and cinnamaldehyde (57 μL , 0.45 mmol) in MeOH (0.3 mL) at room temperature. After being stirred for 3 h, the resultant mixture was purified by silica gel thin-layer chromatography (AcOEt:hexane = 1:3) to provide (*R*)-2-(3-oxo-1-phenylpropyl)malonic acid diethyl ester in 98% yield. The product gave satisfactory NMR and IR spectra. The product was not so stable. Ee was determined by HPLC analysis using a chiral column after oxidation to the corresponding methyl ester (see: Ref. 3). HPLC conditions: Daicel Chiralpak AD, *i*-PrOH:hexane = 1:4, flow rate = 0.5 mL min⁻¹, *t*_R = 13.0 min (major), 19.7 min (minor), 97% ee.