

Communication

Ligand-Controlled Palladium-Catalyzed Regiodivergent Suzuki-Miyaura Cross-Coupling of Allylboronates and Aryl Halides

Yang Yang, and Stephen L. Buchwald

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/ja405950c • Publication Date (Web): 09 Jul 2013 Downloaded from http://pubs.acs.org on July 10, 2013

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Ligand-Controlled Palladium-Catalyzed Regiodivergent Suzuki-Miyaura Cross-Coupling of Allylboronates and Aryl Halides

Yang Yang and Stephen L. Buchwald*

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States *Supporting Information Placeholder*

ABSTRACT: An orthogonal set of catalyst systems has been developed for the Suzuki-Miyaura coupling of 3,3-disubstituted and 3-monosubstituted allylboronates with (hetero)aryl halides. These methods allow for the highly selective preparation of either the α - or the γ isomeric coupling product.

The broad spectrum of intriguing molecular architectures and biological activities of prenylated and reverse-prenylated natural products¹ has spurred extensive efforts toward the development of efficient methods to prepare these compounds. In this context, the selective syntheses of both of these isomers based on a regiodivergent coupling methodology using a common prenyl-metal species and an aryl halide is very appealing (Scheme 1).² Among various types of organometallic reagents, organoboron compounds are most frequently used due to their air and moisture stability, functional group compatibility, ready availability and

Scheme 1. Prenylated Natural Products and Proposed Regioselective Allylation of Aryl Halides.



low toxicity.³ However, the successful development of a coupling process involving 3-substituted allylboronates has been hampered by regioselectivity issues (Scheme 2). In principle, a linear σ allylpalladium complex (5) could isomerize to the corresponding branched σ -allylpalladium species (6) through a π -allyl intermediate (7),⁴ thus producing mixtures of isomers (3 and 4) following reductive elimination. Moreover, transmetallation of a prenylboron reagent (2) with oxidative addition complex 1 could proceed through either an S_E2 or S_E2' pathway, which may also contribute to the poor selectivity often

Scheme 2. Proposed Mechanism.



observed.⁵ Previously, Szabó⁶ and Miyaura⁷ have reported good regioselectivity for the formation of the branched product with 3-monosubstituted allylboron reagents. However, a general and practical γ -selective coupling of 3,3-disubstituted allylboronates to prepare branched products that bear a sterically demanding quaternary center (4), and in particular, a method for *tert*-prenylation, remains to be developed.⁸ To date, Organ's well-tailored NHC-based catalyst remains the only a-selective system to access the linear product (3).⁹ However, this protocol necessitated the use of strong aqueous base at high temperature (5 M aq. KOH in refluxing THF for 24 h), thereby limiting the functional group tolerance. More importantly, a unifying regiodivergent method providing rapid access to both the α - and the γ -regioisomers employing a set of catalysts that are structurally similar but furnish orthogonal selectivities continues to be a daunting challenge. In addition, despite the fact that heterocycles are ubiquitous structural motifs in biologically active compounds, the regioselective allylation of heteroaryl halides has rarely been studied.¹⁰

To overcome these challenges, we reasoned that the choice of ligand would influence the transmetallation mechanism, the rate of $\sigma - \pi - \sigma$ interconversion and the rate of reductive elimination, and thus represent the key to achieving high regioselectivity. Over the past decade, our research group has been engaged in the design, development and utilization of bulky biarylphosphine ligands that have proven effective for a broad range of palladium-catalyzed cross-coupling reactions.^{11,12} In general, the synthesis of these ligands is simple, flexible and allows rational tuning of their steric and electronic properties.¹³ Taken together, these features could greatly facilitate the development of highly α - and γ -selective allylation catalysts that are broadly applicable to a diverse array of functionalized aryl halides.

Table 1. Ligand Evaluation.^a

n-Bu 8	∠Br Me + Me	te 1-1.5 → α Bpin — TH 9 40'	mol% [(ally!)PdCI]₂ n-B 4-6 mol% L HF, K₃PO₄ (aq.), °C or 70 °C, 12 h n	He He He He He He He He He He He He He H
entry	L	α/γ	yield of 10a	yield of 10b
1	L1	<1:99	<1%	55%
2	L2	1:99	<1%	42%
3	L3	<1:99	<1%	99%
4	L4	17:83	16%	80%
5	L5	90:10	63%	7%
6	L6	98:2	$68\% (84\%)^b$	<1%
7	L7	88:12	53%	7%
8	L8	46:64	6%	7%

^{*a*}Reactions with **L1-L4** were carried out with 1 mol% [(allyl)PdCl]₂ and 4 mol% **L** at 40 °C; reactions with **L5-L8** were carried out with 1.5 mol% [(allyl)PdCl]₂ and 6 mol% **L** at 70 °C. Yields were determined by ¹H NMR spectroscopy of the crude reaction mixture. ^{*b*}Reaction was performed in MeCN.



We commenced our study by examining palladium catalysts derived from dialkylbiarylphosphine ligands (Table 1). Although catalysts generated from SPhos (L1) and RuPhos (L2) provided good regioselectivity for the branched product (10b), only moderate conversion of the aryl bromide was observed using 2 mol% Pd. However, the yield of 10b could be increased dramatically using a catalyst derived from L3. Interestingly, replacing the methoxy group on the bottom naph-

thyl ring of L3 with an isoproxy group resulted in inferior selectivity (L4). At this stage, we hypothesized that the use of more sterically demanding ligands would destabilize and/or inhibit the formation of the branched σ -allylpalladium species 6, thus favoring the formation of the linear product (10a). Indeed, the regioselectivity could be reversed when bulkier di-tertbutylbiarylphosphine ligands were employed. In particular, the catalyst generated from *t*-BuXPhos (L6) was found to be highly selective for the production of α -isomer, but further increasing the size of the t-BuXPhos biaryl backbone (L7 and L8) led to less selective catalysts. After extensive optimization, biphasic MeCN/aq. K₃PO₄ was identified as the optimal reaction media for the α -selective coupling. Thus, under the optimized conditions, the *t*-BuXPhos-based catalyst afforded the linear product in 83% yield with excellent selectivity.

Utilizing both protocols, we examined the substrate scope with respect to the aryl halide component (Table 2). A wide variety of aryl bromides, bearing electrondonating or electron withdrawing substituents (**11a-111**), could be effectively converted to the linear or the branched product with high level of regioselectivity. In addition to aryl bromides, aryl chlorides and triflates were also compatible substrates for this transformation (**11f** and **111**). While various heteroaryl halides could be transformed (Table 2 (B)), we were unable, in most cases, to develop a set of regiodivergent conditions for these substrates.¹⁴ Still the examples shown represent the most general ones for transforming a variety of difficult heterocyclic substrates.

Next, we sought to extend the scope of allylboronate substrates that could be employed in the Suzuki-Miyaura coupling (Table 3). In general, unsymmetrical 3,3-disubstituted allylboronates (12a-12c) could be coupled with excellent regioselectivity. Starting from an E/Z mixture of farnesylboronate¹⁵ (12c), either the α - or the γ -isomer could also be accessed under these conditions. Similarly, 3-monosubstituted allylboronates (12d and 12e) represented suitable coupling partners with both conditions. It is noteworthy that an α selective coupling protocol for 3-monosubstituted allylboronates has not been previously described. Tethering the terminal methyl group together with a $(CH_2)_3$ spacer (12f) imposed no detrimental effects on regioselectivity. In addition, a cyclic allylboronate (12g) could be successfully applied in this reaction, maintaining both good yields and selectivities. Finally, 12g reacted smoothly to provide 14g with high diastereocontrol using the γ -selective system.

Although these coupling processes are highly regioselective, we observed scrambling of the olefin configuration when unsymmetrical 3,3-disubstitued allylboronates were subjected to system A (Table 3,

1





^{*a*}Yields are of isolated product, see Supporting Information for details; α/γ ratio and yields in parentheses were determined by ¹H NMR spectroscopy of the crude reaction mixture using 1,3,5-trimethoxybenzene as an internal standard. ^{*b*}L5 was used instead of L6. ^{*c*}60 °C.

Table 3. Substrate Scope of Allylboronates.^a



^{*a*}Yields are of isolated product, see Supporting Information for details; *α*/γ ratio was determined by ¹H NMR spectroscopy of the crude reaction mixture; *E*/*Z* ratio was determined by GC analysis. ^{*b*}60 °C. ^{*c*}Stereochemistry was confirmed by NOESY NMR spectroscopy.

entry 1-2). For example, geranylboronate (12a) and nerylboronate¹⁶ (12b) furnished the linear product with different degrees of isomerization of olefin geometry (13a and 13b). Interestingly, with catalyst system B, 12b delivered higher selectivity for the branched isomer than 12a (14a and 14b). Taken together, these results suggest the intermediacy of a post-transmetallation π -allylpalladium species (7);¹⁷ presumably, the two different (allyl)Pd(Ar)(L) species generated from the transmetallation of (Ar)Pd(X)(L)with 12a and 12b do not fully equilibrate via $\sigma - \pi - \sigma$ interconversion prior to reductive elimination. To gain further insight into the origin of regioselectivity, tertiary allylboronate 15^{18} was subjected to the coupling conditions. With the catalyst derived from L3, reaction of boronates 15 or 9 with 8 afforded drastically different results with regard to regioisomer distribution (Table 4, entry 1-2). Again, these results imply that σ -allyl palladium complexes 5 and 6 do not reach equilibrium via a π -allylpalladium intermediate before reductive elimination occurs.¹⁹ In contrast, using L6-based catalyst, both 15 and 9 provided excellent selectivity for the linear product (entry 3-4). Therefore, if transmetallation of (Ar)Pd(X)(L6) with 9 and 15 furnishes two different σ -allylpalladium complexes (5 or 6), the rate of $\sigma - \pi - \sigma$ interconversion is faster relative to reductive

elimination to form **10b**. Alternatively, the transmetallation of ArPd(X)(**L6**) with **9** and **15** proceeds via two different pathways, S_E2 and S_E2', respectively, to furnish the same σ -allylpalladium complex (**5**), which does not undergo σ - π - σ isomerization at a rate that is competitive with reductive elimination to form **10a**.

Table 4. Mechanistic Insights.



^{*a*}Reaction was carried out in THF/aq. K₃PO₄ at 40 °C for 12 h. ^{*b*}Reaction was carried out in MeCN/aq. K₃PO₄ at 70 °C for 12 h; yield and α/γ ratio were determined by ¹H NMR spectroscopy of the crude reaction mixture using 1,3,5-trimethoxybenzene as an internal standard.

In summary, we have developed a complementary set of palladium-catalyzed regiodivergent protocols for the Suzuki-Miyaura coupling of 3-substituted allylboronates with aryl halides that allow for the rapid construction of allylated arene architectures. This method features excellent regioselectivity, enhanced operational simplicity and a broad scope of aryl halides. Notably, a number of allylated heteroaromatic compounds prepared by the current method would be difficult or tedious to synthesize by other means. Further mechanistic investigations aimed at determining the origin of this regiochemical dichotomy and broadening the application of this method in the context of natural product synthesis are ongoing topics in our laboratory.

ASSOCIATED CONTENT

Supporting Information. Experimental procedures, characterization and spectral data. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

*sbuchwal@mit.edu

ACKNOWLEDGMENT

We acknowledge the National Institutes of Health for financial support of this work (Grant GM46059). We are grateful to Drs. Nathan T. Jui and Jean-Baptiste Langlois for insightful discussions. We thank Dr. J. Robb DeBergh for help with the preparation of this manuscript.

REFERENCES

(1) For recent reviews, see: (a) Williams, R. M.; Stocking, E. M.; Sanz-Cervera, J. F. *Top. Curr. Chem.* **2000**, *209*, 97–173. (b) Li, S. *Nat. Prod. Rep.* **2010**, *27*, 57–78.

(2) For a recent review on prenylation methods, see: Lindel, T.; Marsch, N.; Adla, S. K. *Top. Curr. Chem.* **2012**, *309*, 67–130.

(3) For reviews, see: (a) *Boronic Acids*; Hall, D. G., Ed.; Wiley-VCH: Weinheim, Germany, 2005. (b) *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed.; de Meijere, A., Diederich, F., Eds.; Wiley-VCH: Weinheim, Germany, 2004.

(4) For selected reviews on the $\eta^3 - \eta^1 - \eta^3$ interconversion of allylpalladium intermediate, see: (a) Trost, B. M.; Van Vranken, D. L. *Chem. Rev.* **1996**, *96*, 395–422. (b) Pregosin, P. S.; Salzmann, R. *Coord. Chem. Rev.* **1996**, *155*, 35–68.

(5) For a recent study on the Suzuki-Miyaura coupling of allylboronates where regioselectivity was dictated by the substitution pattern of substrates, see: Glasspoole, B. W.; Ghozati, K.; Moir, J. M.; Crudden, C. M. *Chem. Commun.* **2012**, *48*, 1230–1232.

(6) Sebelius, S.; Olsson, V. J.; Wallner, O. A.; Szabó, K. J. J. Am. Chem. Soc. 2006, 128, 8150–8151.

(7) (a) Yamamoto, Y.; Takada, S.; Miyaura, N. *Chem. Lett.* **2006**, *35*, 704–705. (b) Yamamoto, Y.; Takada, S.; Miyaura, N. *Chem. Lett.* **2006**, *35*, 1368–1369. (c) Yamamoto, Y.; Takada, S.; Miyaura, N. *Organometallics* **2009**, *28*, 152–160.

(8) An exception is the C2-*tert*-prenylation of indoles: Schkeryantz, J. M.; Woo, J. C. G.; Danishefsky, S. J. *J. Am. Chem. Soc.* **1995**, *117*, 7025–7026. For a recent example, see: Kuttruff, C. A.; Zipse, H.; Trauner, D. *Angew. Chem. Int. Ed.* **2011**, *50*, 1402–1405.

(9) Farmer, J. L.; Hunter, H. N.; Organ, M. G. J. Am. Chem. Soc. **2012**, *134*, 17470–17473.

(10) For palladium-catalyzed coupling of other allylmetals, see:
Si: (a) Hatanaka, Y.; Ebina, Y.; Hiyama, T. J. Am. Chem. Soc. 1991, 113, 7076–7077. (b) Hatanaka, Y.; Goda, K.; Hiyama, T. Tetrahedron Lett. 1994, 35, 1279–1282. (c) Hatanaka, Y.; Goda, K.; Hiyama, T. Tetrahedron Lett. 1994, 35, 6511–6514. (d) Denmark, S. E.; Werner, N. S. J. Am. Chem. Soc. 2010, 132, 3612–3620. Sn: (e) Trost, B. M.; Keinan, E. Tetrahedron Lett. 1980, 21, 2595–2598. (f) Echavarren, A. M.; Stille, J. K. J. Am. Chem. Soc. 1987, 109, 5478–5486. (g) Farina, V.; Krishnan, B. J. Am. Chem. Soc. 1987, 109, 5478–5486. (g) Farina, V.; Krishnan, B. J. Am. Chem. Soc. 1991, 113, 9585–9595. (h) Obora, Y.; Tsuji, Y.; Kobayashi, M.; Kawamura, T. J. Org. Chem. 1995, 60, 4647–4649. Zn: (i) Xu, Z.; Negishi, E.-i. Org. Lett. 2008, 10, 4311–4314. (j) Wang, C.; Tobrman, T.; Xu, Z.; Negishi, E.-i. Org. Lett. 2009, 11, 4092–4095.

(11) For recent reviews, see: (a) Surry, D. S.; Buchwald, S. L. *Chem. Sci.* **2011**, *2*, 57–68. (b) Martin, R.; Buchwald, S. L. *Acc. Chem. Res.* **2008**, *41*, 4695–4698.

(12) For recent work on sp²-sp² Suzuki-Miyaura coupling, see: (a) Barder, T. E.; Walker, S. D.; Martinelli, J. R.; Buchwald, S. L. J. Am. Chem. Soc. **2005**, 127, 4685–4696. (b) Billingsley, K. L.; Buchwald, S. L. J. Am. Chem. Soc. **2007**, 129, 3358–3366. (c) Kinzel, T.; Zhang, Y.; Buchwald, S. L. J. Am. Chem. Soc. **2010**, 132, 14073–14075.

(13) Kaye, S.; Fox, J. M.; Hicks, M. A.; Buchwald, S. L. Adv. Syn. Catal. 2001, 343, 789–794.

(14) See Supporting Information for details.

(15) Raducan, M.; Alam, R.; Szabó, K. Angew. Chem. Int. Ed.
 2012, 51, 13050–13053. Zhang, P.; Roundtree, I. A.; Morken, J. P. Org. Lett. 2012, 14, 1416–1419.

(16) For the syntheses of **12a** and **12b**, see: Zhang, P.; Roundtree, I. A.; Morken, J. P. *Org. Lett.* **2012**, *14*, 1416–1419.

(17) To rule out the possibility that 3-substituted allylboronates are configurationally unstable under these reaction conditions, 4-*tert*-butyl-1-bromobenzene was treated with 1.5 equiv of **12b** under conditions A and B and the reaction was stopped at partial conversion. Only **12b** was recovered and no **12a** was observed. Further, the E/Z ratio of allylated products did not change over the course of the reaction, indicating the coupling products were configurationally stable under these conditions. For a review on the configurational stability of allylmetals, see: Hoffmann, R. W. Angew. Chem. Int. Ed. **1982**, 21, 555–566.

(18) For the synthesis of **15**, see: Clary, J. W.; Rettenmaier, T. J.; Snelling, R.; Bryks, W.; Banwell, J.; Wipke, W. T.; Singaram, B. *J. Org. Chem.* **2011**, *76*, 9602–9610. (19) We found that 9 and 15 are not interconvertible under the current reaction conditions.

