

Article

Metal free, Regio and Stereo- Controlled Hydrochlorination and Hydrobromination of Ynones and Ynamides

Xiaojun Zeng, Zhichao Lu, Shiwen Liu, Gerald B. Hammond, and Bo Xu

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.7b02257 • Publication Date (Web): 23 Nov 2017

Downloaded from <http://pubs.acs.org> on November 23, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

Metal free, Regio and Stereo- Controlled Hydrochlorination and Hydrobromination of Ynones and Ynamides

Xiaojun Zeng,^[a] Zhichao Lu,^[b] Shiwen Liu,^[a] Gerald B. Hammond^{[b], *} and Bo Xu^{[a], *}

^[a]College of Chemistry, Chemical Engineering and Biotechnology, Donghua University, 2999 North Renmin Lu, Shanghai 201620, China; e-mail: bo.xu@dhu.edu.cn

^[b]Department of Chemistry, University of Louisville, Louisville, KY 40292, USA; e-mail: gb.hammond@louisville.edu


ABSTRACT: We developed an atom-economical and metal-free method for the regio- and stereo-selective hydrohalogenation of ynones and ynamides using easy to handle DMPU/HX (X = Br or Cl) reagents. The reaction operates under mild conditions and a range of functional groups is well tolerated. We propose that the hydrohalogenation of ynones gives the *anti*-addition products *via* a concerted multi-molecular Ad_{E3} mechanism and that the hydrohalogenation of ynamides produces the *syn*-addition products *via* a cationic keteniminium intermediate.

INTRODUCTION

Vinylhalides are important targets in medicine and materials, and they are also important substrates for cross-coupling reactions.¹ More specifically, halogenated vinyl ketones and halogenated N-vinyl amides (enamides) are especially important targets and synthetic building blocks because they are highly functional sources of vinyl halides.² The hydrohalogenation of alkynes has provided a uniform approach towards vinyl halides,^{1b, 3} and the most straightforward and atom-economical synthesis of halogenated vinyl ketones and haloethenamides has been the *regio*- and *stereo*-selective hydrohalogenation of ynones and ynamides, respectively (Scheme 1, top). For example, the hydrohalogenation of ynones using HBr/water,⁴ AlBr₃/H₂O or LiBr/TFA,^{4c} HCl/water⁵ SnCl₄/H₂O⁵⁻⁶ (Scheme 1, a-e) have been reported, but these methods give poor chemical yields due to side reactions such as hydration. For the hydrohalogenation of ynamides, Hsung and coworkers reported an stereoselective synthesis of 1-haloenamides using *in situ* generated HX from MgX₂ in a wet solvent (Scheme 1f).⁷ Iwasawa and coworkers reported the hydrochlorination of terminal ynamides using HX generated *in situ* from TMSX and water (Scheme 1g).⁸ Sahoo and coworkers demonstrated a one-pot construction of haloenamides using a Ph₃P/H₂O/CX₄ system as the halide source (Scheme 1h).⁹ Recently, Shin and coworkers reported a novel benzoic acid-catalyzed hydrohalogenation of ynamides using N-oxides/di-halogenated ethane system (Scheme 1d)¹⁰ whereas Thibaudeau and coworkers reported a *syn*-addition of HF to the ynamides.¹¹ Although the above mentioned examples underscore the importance of halogenation protocols, the reported methods are either too narrow in scope, or suffer from complex reaction conditions, low chemical yields and/or poor atom-


economy. What is needed is a simple, unified, chemo- and stereo-selective atom-economical hydrohalogenation method for a wide range of ynones and ynamides.

literature systems for halogenation of ynones **1**



- a) HBr/H₂O
- b) HX/H₂O
- c) AlBr₃/H₂O
- d) SnCl₄/H₂O
- e) LiBr/TFA


literature systems for halogenation of ynamides **4**



- f) MgX₂/H₂O, DCM, 60 °C
- g) 1M TMS-X/H₂O
- h) CCl₄ (or CBr₄, CHI₃), PPh₃, H₂O
- i)

- 1) low atom economy; 2) poor yields (hydration by products etc.);
- 3) sometimes low *stereo*-selectivity

This work - an unified hydrohalogenation protocol



Scheme 1. Halogenation of ynones and ynamides.


Recently, we reported a metal-free, hydrogen bonding cluster-enabled addition of sulfonic acids to haloalkynes.¹² We now postulate that a metal-free and atom-economic hydrochlorination or hydrobromination of ynones **1** and ynamides **4** is possible using HX. However, hydrogen halides such as hydrogen chloride and hydrogen bromide are hazardous gases at room temperature. Although many HX reagents (e.g., Et₂O/HCl, H₂O/HCl, isopropanol/HCl,

HOAc/HBr, H₂O/HBr) are commercially available, they exist either in low concentration (e.g. Et₂O/HCl) and/or in protic media (H₂O, isopropanol, HOAc) where the protic solvent hampers the nucleophilicity of HX and also competes with HX in nucleophilic addition. This effect may explain the low efficiency when protic solutions of HX are used (Scheme 1). In this regard, an HX solution in a neutral aprotic medium would be optimal to prevent competitive solvolysis reaction and to achieve high reactivity. Due to its high hydrogen bond basicity, yet low Brønsted basicity,^{13, 14} DMPU (1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone) can form a stable and highly concentrated complex with hydrogen chloride (HCl) and hydrogen bromide (HBr); we have used these complexes in nano-gold catalyzed or non-catalyzed hydrohalogenation of alkenes and alkynes.¹⁵ We are now glad to report an atom-economical and metal-free method for the regio- and stereoselective hydrohalogenation of ynones and ynamides using easy to handle DMPU/HX (X = Br or Cl) reagents.

RESULTS AND DISCUSSION

We carried out the hydrochlorination of **1a** as our model reaction (Table 1). We were delighted to find that DMPU/HCl gave better yields, higher *anti*-selectivity and fewer by-products compared to commercial HCl reagents (Table 1, entries 1-4). Mixing DMPU/HCl with **1a** in DCM from 0 °C to room temperature produced the desired product, chloro-unsaturated ketone **2**, in 98% yield (Table 1, entry 4). Screening of other solvents indicated that weakly coordinating solvents like DCM, DCE and toluene were better than more coordinating solvents such as CH₃CN, THF and DMF (Table 1, entries 5-9).


Table 1. Optimization of the hydrochlorination of alkynyl ketone.^a



entry	HCl reagent	solvent	2 / % ^b (2a : 2a')
1	Et ₂ O/HCl	DCM	70 (6:1)
2	1,4-dioxane/HCl	DCM	81 (7:1)
3	Isopropanol/HCl	DCM	92 (3:1)
4	DMPU/HCl	DCM	98 (12:1)
5	DMPU/HCl	MeCN	89 (6:1)
6	DMPU/HCl	Toluene	88 (7:1)
7	DMPU/HCl	THF	83 (5:1)
8	DMPU/HCl	DMF	78 (2:1)
9	DMPU/HCl	DCE	97 (8:1)


^a Reaction conditions: **1a** (0.2 mmol), HCl reagent (0.4 equiv) in solvent (0.5 mL), 0 °C to rt for 8 h. ^b Determined by GC-MS analysis.

With the optimized conditions in hand, we explored the substrate scope and functional group tolerance for the hydrochlorination of alkynyl ketones (Table 2, top). We evaluated the effect of different [R¹, R²] combinations of ynones **1** (Table 2). Combinations of [R¹ = aryl, R² = alkyl] (Table 2, **2a-l**), [R¹ = aryl, R² = aryl] (Table 2, **2n**), [R¹ = alkyl, R² = aryl] (Table 2, **2o**) worked very well; good chemical yields and high *anti*-selectivity were observed. The bulky R² groups (e.g. *t*-Bu, Cy-) did not affect chemical yields (Table 2, **2b**, **2c**). Equally, the substitution pattern (*meta*, *para*) or the electronic properties of the substituents (electron deficient or rich) on R¹ or R² played only a small role; good yields were obtained regardless (Table 2, **2e-2m**, **2o**). Various functional groups such as esters (Table 2, **2e**, **2k**), ether (Table 2, **2i**), halides (Table 2, **2f**, **2h**), nitrile (Table 2, **2l**), nitro (Table 2, **2o**), and sulfonate (Table 2, **2j**) were well tolerated. Similar conditions worked very well for the hydrobromination of ynones when DMPU/HBr was used (Table 2, bottom). In these cases, we observed a very similar reactivity pattern and functional group tolerance (Table 2, **3a-3j**). We also tested the hydrohalogenation of di-alkyl substituted ynones. Unfortunately we got mostly hydration products and other by-products. We also attempted to conduct hydroiodination using DMPU-HI reagent, but the preparation of DMPU-HI was not successful (formation of I₂ when mixing HI and DMPU).

Table 2. Hydrohalogenation of ynones.^a

^aReaction conditions: ynones **1** (0.2 mmol), DMPU/HX (0.4 mmol) in DCM (0.5 mL), 0 °C to rt for 8 h; trace *syn*-addition products were also observed in most case, but they can be easily removed by chromatography.


Ynamides are readily available compounds that have found wide applications in organic synthesis.¹⁶ We were glad to find that our protocol can also be used in the hydrohalogenation of ynamides **4** (Table 3). We first explored the substrate scope and functional group tolerance in the hydrobromination of ynamides **4** (Table 3, top). Despite the strong acidity of the DMPU/HBr complex and the fact that ynamides are acid sensitive compounds, the substrate scope was broad: both aliphatic and aromatic substituted ynamides produced *syn*-addition products exclusively in good to excellent yields (Table 3): combinations of [EWG = Ts, R² = Bn], [EWG = Ts, R² = *n*-heptyl], [EWG = Ms, R² = methyl], [EWG = Ms, R² = phenyl] all gave good yields (Table 3, **5a-d**). Also, the substitution pattern (*para*-, *ortho*- or *meta*-) on the aryl ring of R¹ only had a small effect on the efficiency of the reaction, high yields were obtained regardless (Table 3, **5e-5i**). It should be noted that acid sensitive functional groups such as nitrile and methyl ester survived our conditions (Table 1, **5j-5k**).

Table 3. Hydrohalogenation of ynamides.^a


^aReaction conditions: ynamide **4** (0.2 mmol), DMPU/HX (0.4 mmol) in DCM (0.5 mL), 0 °C to rt for 8 h.

Next, we investigated the hydrochlorination of ynamides **4** using DMPU/HCl as the chlorination reagent (Table 3, bottom). This protocol also had wide substrate scope and good functional group tolerance (Table 3 bottom). Ynamides **4** with combinations of [EWG = Ts/Ms/carbamate, R² = alkyl, aryl, allyl, homoallyl] delivered **6** in good yields as well as in excellent regio- and stereo-control (Table 3, **6a-g**). Interestingly, acid sensitive functional groups such as alkene, allyl and carbamate survived the reaction conditions. We also investigated the effects of the substitution pattern in R¹ on the efficiency of reactions. For ynamides with various R¹ groups (R¹ = naphthyl, phenyl rings with electron-donating groups or electron-withdrawing groups, heteroaryls such as thiophene), high yields of **6** were obtained regardless (Table 3, **6h-o**). Remarkably, ynamides featuring a complex cholesterol scaffold furnished the desired chlorinated product in 82% yield (Table 3, **6o**).

Our methodology can be used in larger scale synthesis without complications. The hydrohalogenation of ynone **1a** (5 mmol scale) gave chlorinated product **2a** and brominated product **3a** in 85% and 73% yields, respectively (Scheme 2a). Ynamide **4a** (1.0 gram scale) was successfully converted to **6a** in 93% yield (Scheme 2b).



Scheme 2. Gram scale hydrohalogenations.




Scheme 3. Divergent syntheses from halogenated products.

Next, we utilized these halogenated alkene products in transition metal catalyzed cross-coupling reactions (Scheme 3). Suzuki coupling of halogenated vinyl ketones **2** and **3** with phenyl boronic acid gave good yields of the coupling products (Scheme 3a). Similarly, the Pd catalyzed Suzuki coupling of chlorinated enamide **6a** with phenyl boronic acid (Scheme 3b), and the Sonogashira coupling of brominated enamide **5c** with ethynyltrimethylsilane (Scheme 3c) gave good yields of tri-substituted enamides **8a** and **9a**, respectively. On the other hand, copper catalysts such as CuCN was not a good catalyst for coupling of **5c**.

Our proposed mechanism is shown in Scheme 4. For the hydrohalogenation of ynamides (Scheme 4a), the high acidity of DMPU/HX facilitates the rate-determining proton transfer step, which produces the key intermediate--keteniminium **A**.¹⁷ Because the nitrogen atom polarizes the triple bond, the regioselectivity of this protonation is mainly determined by the charge distribution on C1 and C2 of ynamide **4**. Due to the resonance contribution of the lone pair on the nitrogen atom, the NBO charge density of C2 is significantly higher than C1, so DMPU/HX should pro-

tonate C2 preferentially (Scheme 4a). This pathway eventually leads to the α -regioselectivity observed. Also, DFT geometry optimization of **A** indicates that the keteniminium intermediate possesses a linear geometry, with its upper face being sterically hindered by the R¹ group, thus favoring the nucleophile (bromide/chloride) *syn* approach (formation of the *syn*-addition product).^{1b}



Scheme 4. Proposed mechanism. LUMO of keteniminium was calculated at B3LYP/6-311+G(2df,2p) level of theory.

For the hydrohalogenation of the electron deficient ynone **1**, the direct protonation of **1** is more difficult, so we proposed a different mechanism (Scheme 4b). The exclusive formation of the Z-isomer (*anti*-addition) suggested the possibility of an Ad_E3 mechanism¹⁸ because the bromide concentration in the reaction medium is not low.¹⁹ This concerted three-molecule process should furnish the *anti*-addition product. The *regio*-selectivity found can be rationalized by the fact that the NBO charge density of C1 is significantly higher than C2 because of the resonance participation of the carbonyl group (Scheme 4b). This pathway should eventually lead to the β -regioselectivity found.

CONCLUSION

In conclusion, we have developed an atom-economical and metal-free method for the regio- and stereo-selective hydrohalogenation of ynones and ynamides using easily handled DMPU/HX (X = Br or Cl) reagents. The reaction operates under mild conditions and a range of functional groups are well tolerated. Further applications of these new halogenation reagents are currently underway in our laboratory and will be reported in due course.

EXPERIMENTAL SECTION

¹H NMR (400 MHz or 500 MHz) and ¹³C NMR (100 MHz or 125 MHz) spectra were recorded on a Bruker NMR apparatus. The chemical shifts are reported in δ (ppm) values (¹H and ¹³C NMR relative to CHCl₃, δ 7.26 ppm for ¹H NMR and δ 77.0 ppm for ¹³C NMR). Or alternatively, ¹H NMR chemical shifts were referenced to tetramethylsilane signal (0 ppm). Multiplicities are recorded by s (singlet), d (doublet), t (triplet), q (quartet), p (pentet), h (hextet), m (multiplet) and br (broad). Coupling constants (J), are reported in Hertz (Hz). GC analyses were performed using a Shimadzu GC-2010 ultra gas chromatography-mass spectrometry instrument equipped with a Shimadzu AOC-20S autosampler. Commercial reagents and solvents were obtained from the commercial providers and used without further purification. The products were purified using a commercial flash chromatography system or a regular glass column. TLC was developed on silica gel 60 F254 glass plates. The DMPU/HX reagents were prepared using our published procedures¹⁵.

Methyl 3-(3-oxooct-1-yn-1-yl) benzoate (1e).²⁰ White solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.23 (s, 1H), 8.10 (d, *J* = 6.6 Hz, 1H), 7.73 (d, *J* = 7.2 Hz, 1H), 7.47 (s, 1H), 3.93 (s, 3H), 2.66 (t, *J* = 7.2 Hz, 2H), 1.91 – 1.67 (m, 2H), 1.35 (d, *J* = 3.3 Hz, 4H), 0.91 (t, *J* = 7.5 Hz, 3H). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 187.9, 165.8, 136.9, 133.9, 131.3, 130.8, 128.7, 120.6, 88.8, 88.2, 52.5, 45.5, 31.1, 23.8, 22.3, 13.8. HRMS (ESI-FT-ICR) calcd. for C₁₆H₁₉O₃ [M+H]⁺: 259.1334, found: 259.1327.

4-(4-Fluoro-3-methylphenyl)but-3-yn-2-one (1h). Yellow solid. Mp 42–43°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.61 – 7.33 (m, 2H), 7.00 (t, *J* = 8.8 Hz, 1H), 2.42 (s, 3H), 2.26 (s, 3H). ¹⁹F NMR (376 MHz, Chloroform-*d*) δ -109.52 – -112.87 (m). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 184.4, 162.7 (d, *J* = 252.4 Hz), 136.5, 132.7 (d, *J* = 8.9 Hz), 125.9 (d, *J* = 18.2 Hz), 115.8, 115.6, 89.7, 87.9, 32.6 (d, *J* = 5.3 Hz), 14.3. HRMS (ESI-FT-ICR) calcd. for C₁₂H₁₀FO [M+H]⁺: 177.0710, found: 177.0703.

3-(3-Oxobut-1-yn-1-yl)phenyl methanesulfonate (1j).²⁰ White solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.57 – 7.34 (m, 4H), 3.17 (s, 3H), 2.44 (s, 3H). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 184.1, 148.8, 131.8, 130.4, 126.2, 124.6, 122.0, 88.9, 87.6, 37.7, 32.7. HRMS (ESI-FT-ICR) calcd. for C₁₁H₁₁SO₄ [M+H]⁺: 239.0378, found: 239.0371.

Methyl 3-(3-oxobut-1-yn-1-yl)benzoate (1k).²⁰ White solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.22 (s, 1H), 8.12 – 8.00 (m, 1H), 7.79 – 7.64 (m, 1H), 7.46 – 7.42 (m, 1H), 3.92 (s, 3H), 2.44 (s, 3H). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 184.2, 165.8, 136.8, 134.0, 131.5, 130.8, 128.8, 120.4, 88.6, 88.5, 52.4, 32.7. HRMS (ESI-FT-ICR) calcd. for C₁₂H₁₁O₃ [M+H]⁺: 203.0708, found: 203.0720.

4-(3-Oxobut-1-yn-1-yl)benzonitrile (1l).²⁰ White solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.66 – 7.63 (m, 4H), 2.46 (s, 3H). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 183.9, 133.2, 132.2, 124.7, 117.8, 113.9, 90.5, 86.8, 32.7. HRMS

(ESI-FT-ICR) calcd. for C₁₁H₈NO[M+H]⁺: 170.0606, found: 170.0600.

4-(4-Acetylphenyl)but-3-yn-2-one (1m).²⁰ White solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.93 (d, *J* = 8.2 Hz, 2H), 7.62 (d, *J* = 8.2 Hz, 2H), 2.59 (s, 3H), 2.44 (s, 3H). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 196.9, 184.2, 138.0, 133.0, 128.3, 124.5, 89.9, 88.3, 32.7, 32., 26.64. HRMS (ESI-FT-ICR) calcd. for C₁₂H₁₁O₂ [M+H]⁺: 187.0759, found: 187.0753.

2. Synthesis of ynamides 4

General procedure. To a mixture of alkynyl bromide (3.0 mmol), 1,10-phenanthroline (0.6 mmol, 108 mg), CuSO₄·5H₂O (0.3 mmol, 149 mg) and K₂CO₃ (6.0 mmol, 828 mg) in dry toluene (15.0 mL), amide (3.6 mol) was added. The reaction mixture was stirred at 70 °C under the nitrogen atmosphere. The progress of the reaction was monitored periodically by TLC. After the completion of the reaction, the reaction mixture was filtered through a pad of celite and the solvent was removed in vacuum. The resulting residue was purified by chromatography on silica gel to give the desired products.

N-(2-(cyclohex-1-en-1-yl)ethyl)-4-methyl-N-(phenylethynyl)benzenesulfonamide (4f). Pale yellow oil. ¹H NMR (500 MHz, Chloroform-*d*) δ 7.86 (d, *J* = 8.3 Hz, 2H), 7.46 – 7.34 (m, 4H), 7.35 – 7.26 (m, 3H), 5.48 (s, 1H), 3.65 – 3.34 (m, 2H), 2.47 (s, 3H), 2.33 (t, *J* = 7.6 Hz, 2H), 1.99–1.94 (m, 4H), 1.73 – 1.59 (m, 2H), 1.57–1.52 (m, 2H). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 144.5, 134.8, 133.3, 131.3, 129.7, 128.3, 127.7, 124.2, 123.0, 82.5, 70.8, 50.3, 36.4, 28.2, 25.3, 22.8, 22.2, 21.7. HRMS (ESI-FT-ICR) calcd. for C₂₃H₂₆NO₂S [M+H]⁺: 380.1684, found: 380.1678.

(3*S*,10*R*,13*R*,17*R*)-10,13-dimethyl-17-((*R*)-6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1*H*-cyclopenta[*a*]phenanthren-3-yl 6-((*N*,4-dimethylphenyl)sulfonamido)hex-5-ynoate (4o). White solid. Mp 94–95°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.76 (d, *J* = 8.3 Hz, 2H), 7.34 (d, *J* = 7.9 Hz, 2H), 5.35 (d, *J* = 4.9 Hz, 1H), 4.62 – 4.60 (m, 1H), 2.99 (s, 3H), 2.43 (s, 3H), 2.35 – 2.31 (m, 6H), 2.07 – 1.05 (m, 28H), 1.00 (s, 3H), 0.89 (d, *J* = 6.5 Hz, 3H), 0.85 (d, *J* = 1.8 Hz, 3H), 0.83 (d, *J* = 1.8 Hz, 3H), 0.66 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 172.5, 144.5, 139.6, 133.1, 129.7, 127.8, 127.7, 122.6, 75.6, 74.0, 67.4, 56.7, 56.1, 50.0, 42.3, 39.7, 39.5, 38.1, 37.0, 36.6, 36.2, 35.8, 33.4, 31.9, 31.8, 28.2, 28.0, 27.8, 24.2, 23.8, 22.8, 22.5, 21.7, 21.0, 19.3, 18.7, 17.9, 11.8. HRMS (ESI⁺) calcd. for C₄₁H₆₁NO₄S [M+H]⁺: 664.4400, found: 664.4385.

3. Hydrochlorination of ynones 1

To a solution of ynone **1** (0.2 mmol) in DCM (0.5 mL) was added DMPU/HCl (43 wt/wt%) (34 mg, 0.4 mmol) at 0 °C, then the mixture was warmed to room temperature and was stirred for 8 h. After completion, the solvent was evaporated under the reduced pressure and the residue was purified by silica gel column chromatography (hexanes/EtOAc as elute solvent) to afford the corresponding products **2**.

(*Z*)-4-chloro-4-phenylbut-3-en-2-one (2a).^{3e} Purified by flash column chromatography (ethyl acetate / hexanes = 1:20) give the product as colorless oil, 30.2 mg 84% yield. ¹H NMR (400

MHz, Chloroform-*d*) δ 7.86 – 7.55 (m, 2H), 7.58 – 7.31 (m, 4H), 6.77 (s, 1H), 2.47 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.7, 143.1, 137.3, 130.7, 128.6, 127.3, 124.7, 31.9.

(Z)-1-chloro-4,4-dimethyl-1-phenylpent-1-en-3-one (2b).²¹ Purified by flash column chromatography (ethyl acetate / hexanes = 1:20) give the product as yellow oil, 40.8 mg, 92% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.77 – 7.56 (m, 2H), 7.59 – 7.25 (m, 3H), 7.06 (s, 1H), 1.22 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 203.6, 143.2, 137.8, 130.4, 128.6, 127.2, 119.9, 44.5, 26.3. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{13}\text{H}_{16}\text{ClO}[\text{M}+\text{H}]^+$: 223.0890, found: 223.0885.

(Z)-3-chloro-1-cyclohexyl-3-phenylprop-2-en-1-one (2c).²² Purified by flash column chromatography (ethyl acetate / hexanes = 1:20) give the product as yellow oil, 42.1 mg, 85% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.90 – 7.51 (m, 2H), 7.41–7.36 (m, 3H), 6.84 (s, 1H), 2.58 – 2.54 (m, 1H), 1.92 – 1.90 (m, 2H), 1.80 – 1.78 (m, 2H), 1.67 – 1.65 (m, 1H), 1.50 – 1.05 (m, 5H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 201.5, 142.6, 137.55, 130.47, 128.57, 127.24, 122.66, 51.63, 28.35, 25.85, 25.65. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{15}\text{H}_{18}\text{ClO}[\text{M}+\text{H}]^+$: 249.1046, found: 249.1039.

(Z)-1-chloro-1-phenyloct-1-en-3-one (2d). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow oil, 41.5 mg, 88% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.67 – 7.75 (m, 2H), 7.50 – 7.30 (m, 3H), 6.78 (s, 1H), 2.68 (t, J = 7.4 Hz, 2H), 1.67– 1.63 (m, 2H), 1.33 – 1.30 (m, 4H), 0.90 (t, J = 6.8 Hz, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 199.01, 142.31, 137.44, 130.53, 128.59, 127.26, 123.86, 44.44, 31.38, 23.68, 22.48, 13.92. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{14}\text{H}_{18}\text{ClO}[\text{M}+\text{H}]^+$: 237.1046, found: 237.1040.

Methyl (Z)-3-(1-chloro-3-oxooct-1-en-1-yl)benzoate (2e). Purified by flash column chromatography (ethyl acetate / hexanes = 1/20), yellow oil. NMR (400 MHz, Chloroform-*d*) δ 8.31 (s, 1H), 8.07 (d, J = 7.1 Hz, 1H), 7.85 (d, J = 7.1 Hz, 1H), 7.48 (t, J = 7.8 Hz, 1H), 6.83 (s, 1H), 3.93 (s, 3H), 2.68 (t, J = 7.3 Hz, 2H), 1.87 – 1.59 (m, 2H), 1.35–1.29 (m, 4H), 0.88 (t, J = 6.6 Hz, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 198.8, 166.2, 140.9, 137.8, 131.5, 131.4, 130.7, 128.8, 128.2, 124.8, 52.4, 44.5, 31.4, 23.6, 22.5, 13.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{16}\text{H}_{20}\text{ClO}_3[\text{M}+\text{H}]^+$: 295.1101, found: 295.1094.

(Z)-4-chloro-4-(4-chlorophenyl)but-3-en-2-one (2f). Purified by flash column chromatography (ethyl acetate/hexane = 1:20) give the product as pale yellow oil, 35 mg, 82% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.60 (d, J = 8.6 Hz, 2H), 7.37 (d, J = 8.6 Hz, 2H), 6.73 (s, 1H), 2.46 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.3, 141.6, 136.9, 135.7, 128.9, 128.5, 124.9, 31.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{10}\text{H}_9\text{Cl}_2\text{O}[\text{M}+\text{H}]^+$: 215.0030, found: 215.0024.

(Z)-4-(4-bromophenyl)-4-chlorobut-3-en-2-one (2g). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow solid, 44.8 mg, 88% yield. Mp 42–43°C ^1H NMR (400 MHz, Chloroform-*d*) δ 7.53 (s, 4H), 6.74 (s, 1H), 2.45 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.7, 141.7, 136.1, 131.9, 128.7, 125.2, 124.9, 31.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{10}\text{H}_7\text{ClBrO}[\text{M}+\text{H}]^+$: 256.9369, found: 256.9362.

(Z)-4-chloro-4-(4-fluoro-3-methylphenyl)but-3-en-2-one (2h). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow oil, 38.1 mg, 90% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.48 – 7.42 (m, 2H), 7.01 (t, J = 8.8 Hz, 1H), 6.69 (s, 1H), 2.44 (s, 3H), 2.28 (s, 3H). ^{19}F NMR (376 MHz, Chloroform-*d*) δ -101.3 – -135.0 (m, 1F). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.4, 162.7 (d, J = 250.8 Hz), 142.1, 133.1 (d, J = 3.8 Hz), 130.7 (d, J = 6.1 Hz), 126.7 (d, J = 8.7 Hz), 125.4 (d, J = 17.9 Hz), 124.3, 115.3 (d, J = 23.1 Hz), 31.9, 14.6 (d, J = 3.3 Hz). HRMS (ESI-FT-ICR) calcd. for $\text{C}_{11}\text{H}_{11}\text{ClFO}[\text{M}+\text{H}]^+$: 213.0482, found: 213.0477.

(Z)-4-chloro-4-(3-methoxyphenyl)but-3-en-2-one (2i). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow oil, 31.9 mg, 76% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.29 (t, J = 7.9 Hz, 1H), 7.23 (d, J = 7.8 Hz, 1H), 7.18 (s, 1H), 7.00 – 6.91 (m, 1H), 6.75 (s, 1H), 3.82 (s, 3H), 2.45 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.6, 159.6, 142.7, 138.7, 129.6, 124.9, 119.6, 116.3, 112.9, 55.4, 31.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{11}\text{H}_{12}\text{ClO}_2[\text{M}+\text{H}]^+$: 211.0526, found: 211.0519.

(Z)-3-(1-chloro-3-oxobut-1-en-1-yl)phenyl methanesulfonate (2j). Purified by flash column chromatography (ethyl acetate / hexanes = 1:10), pale red oil, 46.5 mg, 85% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.63 (d, J = 7.9 Hz, 1H), 7.58 (s, 1H), 7.46 (t, J = 8.0 Hz, 1H), 7.36 (ddd, J = 8.1, 2.5, 1.1 Hz, 1H), 6.77 (s, 1H), 3.18 (s, 3H), 2.46 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.3, 149.1, 140.6, 139.4, 130.3, 126.1, 125.8, 124.1, 121.1, 37.7, 31.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{11}\text{H}_{12}\text{ClSO}_4[\text{M}+\text{H}]^+$: 275.0145, found: 275.0140.

Methyl (Z)-3-(1-chloro-3-oxobut-1-en-1-yl)benzoate (2k). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), yellow solid, 40.9 mg, 86% yield. Mp 44–45°C ^1H NMR (400 MHz, Chloroform-*d*) δ 8.31 (s, 1H), 8.08 (d, J = 7.7 Hz, 1H), 7.85 (d, J = 8.9 Hz, 1H), 7.49 (t, J = 7.8 Hz, 1H), 6.82 (s, 1H), 3.93 (s, 3H), 2.47 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.4, 166.1, 141.6, 137.6, 131.5, 130.7, 128.9, 128.2, 125.4, 52.4, 31.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{12}\text{H}_{12}\text{ClO}_3[\text{M}+\text{H}]^+$: 239.0475, found: 239.0470.

(Z)-4-(1-chloro-3-oxobut-1-en-1-yl)benzonitrile (2l). Purified by flash column chromatography (ethyl acetate / hexanes = 1:10), yellow solid, 33.2 mg, 81% yield. Mp 76–78°C. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.77 (d, J = 8.3 Hz, 2H), 7.70 (d, J = 8.3 Hz, 2H), 6.79 (s, 1H), 2.47 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 196.1, 141.4, 140.2, 132.4, 127.9, 126.8, 117.9, 114.1, 31.8. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{11}\text{H}_9\text{ClNO}[\text{M}+\text{H}]^+$: 206.0373, found: 206.0366.

(Z)-4-(4-acetylphenyl)-4-chlorobut-3-en-2-one (2m). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), yellow oil, 37.4 mg, 85% yield. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.97 (d, J = 8.5 Hz, 2H), 7.75 (d, J = 8.6 Hz, 2H), 6.81 (s, 1H), 2.61 (s, 3H), 2.48 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 197.1, 196.4, 141.3, 138.4, 128.5, 127.5, 126.2, 31.9, 26.7. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{12}\text{H}_{12}\text{ClO}_2[\text{M}+\text{H}]^+$: 223.0526, found: 223.0520.

(Z)-3-chloro-1,3-diphenylprop-2-en-1-one (2n). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow oil, 43 mg, 89% yield. ^1H NMR (500 MHz, Chloroform-*d*) δ 8.02 (d, J = 7.0 Hz, 2H), 7.83 – 7.72 (m, 2H), 7.60 (t,

$J = 7.3$ Hz, 1H), 7.55 – 7.42 (m, 5H), 7.37 (s, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 189.8, 143.3, 133.4, 130.6, 128.7, 128.7, 128.7, 127.2, 121.5. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{15}\text{H}_{12}\text{ClO}$ $[\text{M}+\text{H}]^+$: 243.0577, found: 243.0569.

(Z)-3-chloro-1-(4-nitrophenyl)-3-phenylprop-2-en-1-one (2o).^{3c} Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), yellow oil, 47 mg, 82% yield. ^1H NMR (500 MHz, Chloroform- d) δ 8.34 (d, $J = 8.9$ Hz, 2H), 8.14 (d, $J = 8.9$ Hz, 2H), 7.79 (dt, $J = 6.7, 1.6$ Hz, 2H), 7.61 – 7.41 (m, 3H), 7.37 (s, 1H). HRMS (ESI-FT-ICR) calcd. for $\text{C}_{15}\text{H}_{11}\text{ClNO}_3$ $[\text{M}+\text{H}]^+$: 288.0427, found: 288.0419.

4. Hydrobromination of ynones 1

To a solution of ynone **1** (0.2 mmol) in DCM (0.5 mL) was added DMPU/HBr (60 wt/wt%) (54 mg, 0.4 mmol) at 0 °C, then the mixture was warmed to room temperature and was stirred for 8 h. After completion, the solvent was evaporated under the reduced pressure and the residue was purified by silica gel column chromatography (hexanes/EtOAc as elute solvent) to afford the corresponding products **3**.

(Z)-4-bromo-4-phenylbut-3-en-2-one (3a).²³ Purified by flash column chromatography (ethyl acetate / hexanes = 1:25), pale yellow oil, 33.8 mg, 76% yield. ^1H NMR (400 MHz, Chloroform- d) δ 7.62 (dd, $J = 7.0, 2.6$ Hz, 2H), 7.45 – 7.29 (m, 3H), 6.99 (s, 1H), 2.43 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform- d) δ 196.68, 139.30, 133.99, 130.44, 128.55, 128.26, 128.14, 127.90, 31.77.

(Z)-1-bromo-4,4-dimethyl-1-phenylpent-1-en-3-one (3b). Purified by flash column chromatography (ethyl acetate / hexanes = 1:25), yellow oil, 43 mg, 81% yield. ^1H NMR (500 MHz, Chloroform- d) δ 7.73 – 7.54 (m, 2H), 7.43–7.40 (m, 3H), 7.24 (s, 1H), 1.24 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 204.10, 139.83, 134.18, 130.16, 128.52, 128.10, 124.01, 44.38, 26.27. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{13}\text{H}_{16}\text{BrO}$ $[\text{M}+\text{H}]^+$: 267.0385, found: 267.0378.

(Z)-3-bromo-1-cyclohexyl-3-phenylprop-2-en-1-one (3c). Purified by flash column chromatography (ethyl acetate / hexanes = 1:25), yellow oil, 46.1 mg, 79% yield. ^1H NMR (500 MHz, Chloroform- d) δ 7.74 – 7.57 (m, 2H), 7.52 – 7.30 (m, 3H), 7.07 (s, 1H), 2.57 – 2.52 (m, 1H), 2.06 – 1.90 (m, 2H), 1.85 – 1.82 (m, 2H), 1.74 – 1.63 (m, 1H), 1.54 – 1.11 (m, 5H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 201.9, 139.6, 133.6, 130.2, 128.5, 128.1, 126.4, 51.7, 28.3, 25.9, 25.6. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{15}\text{H}_{18}\text{BrO}$ $[\text{M}+\text{H}]^+$: 293.0541, found: 293.0534.

(Z)-1-bromo-1-phenyloct-1-en-3-one (3d). Purified by flash column chromatography (ethyl acetate / hexanes = 1:25), yellow oil, 44.8 mg, 80% yield. ^1H NMR (500 MHz, Chloroform- d) δ 7.80 – 7.54 (m, 2H), 7.46 – 7.31 (m, 3H), 7.01 (s, 1H), 2.66 (t, $J = 7.4$ Hz, 2H), 1.70 – 1.68 (m, 2H), 1.50 – 1.20 (m, 4H), 0.92 (t, $J = 7.0$ Hz, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 199.2, 139.5, 133.3, 130.3, 128.5, 128.1, 127.2, 44.4, 31.4, 23.6, 22.5, 13.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{14}\text{H}_{18}\text{BrO}$ $[\text{M}+\text{H}]^+$: 281.0541, found: 281.0535.

Methyl (Z)-3-(1-bromo-3-oxooct-1-en-1-yl)benzoate (3e). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), yellow oil, 54.7 mg, 81% yield. ^1H NMR (500 MHz, Chloroform- d) δ 8.26 (s, 1H), 8.06 (d, $J = 8.2$ Hz, 1H), 7.82 (d, $J = 7.8$ Hz, 1H), 7.48 (t, $J = 7.8$ Hz, 1H), 7.05 (s, 1H), 3.94 (s, 3H), 2.66 (t, $J = 7.3$ Hz, 2H), 1.86 – 1.52 (m, 2H), 1.34 (dt, $J = 7.7, 3.6$ Hz, 4H), 0.90 (t, $J = 6.8$ Hz, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 199.0, 166.2, 139.8, 132.5, 131.7, 131.1, 130.6, 128.9, 128.8, 128.0, 52.4, 44.4, 31.3, 23.5, 22.5, 13.9. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{16}\text{H}_{20}\text{BrO}_3$ $[\text{M}+\text{H}]^+$: 339.0596, found: 339.0589.

(Z)-4-(4-acetylphenyl)-4-bromobut-3-en-2-one (3f). Purified by flash column chromatography (ethyl acetate / hexanes = 1:10), yellow oil, 42.5 mg, 80% yield. ^1H NMR (500 MHz, Chloroform- d) δ 197.05, 196.47, 143.34, 138.15, 131.97, 129.30, 128.45, 128.36, 31.66, 26.73. $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 197.1, 196.5, 143.3, 138.2, 132.0, 129.3, 128.5, 128.4, 31.7, 26.7. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{12}\text{H}_{12}\text{BrO}_2$ $[\text{M}+\text{H}]^+$: 267.0012, found: 267.0014.

(Z)-4-bromo-4-(4-bromophenyl)but-3-en-2-one (3g). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow oil, 43.3 mg, 72% yield. ^1H NMR (500 MHz, Chloroform- d) δ 7.54 (d, $J = 8.3$ Hz, 2H), 7.50 (d, $J = 8.7$ Hz, 2H), 6.98 (s, 1H), 2.44 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 196.4, 138.2, 132.4, 131.8, 129.6, 128.2, 124.9, 31.7. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{10}\text{H}_9\text{Br}_2\text{O}$ $[\text{M}+\text{H}]^+$: 302.9015, found: 302.9013.

(Z)-4-bromo-4-(3-methoxyphenyl)but-3-en-2-one (3h). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), yellow oil, 24.9 mg, 49% yield. ^1H NMR (500 MHz, Chloroform- d) δ 7.30 (d, $J = 7.8$ Hz, 2H), 7.21 (d, $J = 7.7$ Hz, 2H), 7.16 (s, 1H), 7.00 (s, 1H), 6.96 (d, $J = 8.2$ Hz, 2H), 3.85 (s, 3H), 2.45 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 196.7, 159.5, 140.7, 133.6, 129.5, 128.1, 120.4, 116.0, 113.9, 55.4, 31.7. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{11}\text{H}_{12}\text{BrO}_2$ $[\text{M}+\text{H}]^+$: 255.0021, found: 255.0013.

Methyl (Z)-3-(1-bromo-3-oxobut-1-en-1-yl)benzoate (3i). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), yellow oil, 44.9 mg, 80% yield. ^1H NMR (500 MHz, Chloroform- d) δ 8.27 (s, 1H), 8.07 (d, $J = 7.8$ Hz, 1H), 7.83 (d, $J = 7.5$ Hz, 1H), 7.49 (t, $J = 7.8$ Hz, 1H), 7.05 (s, 1H), 3.95 (s, 3H), 2.45 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 196.4, 166.2, 139.7, 132.5, 132.4, 131.3, 130.6, 129.0, 128.8, 128.7, 52.4, 31.7. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{12}\text{H}_{12}\text{BrO}_3$ $[\text{M}+\text{H}]^+$: 282.9970, found: 282.9963.

(Z)-3-bromo-1-(4-nitrophenyl)-3-phenylprop-2-en-1-one (3j). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow oil, 51.5 mg, 78% yield. ^1H NMR (500 MHz, Chloroform- d) δ 8.35 (d, $J = 8.9$ Hz, 2H), 8.16 (d, $J = 8.9$ Hz, 2H), 7.79 – 7.68 (m, 2H), 7.55 (s, 1H), 7.51 – 7.43 (m, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz, Chloroform- d) δ 188.6, 150.4, 142.04, 138.79, 137.17, 130.91, 129.66, 128.11, 124.15, 123.98. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{15}\text{H}_{11}\text{BrNO}_3$ $[\text{M}+\text{H}]^+$: 331.9922, found: 331.9914.

5. Hydrobromination of ynamides 4

To a solution of ynamides **4** (0.2 mmol) in DCM (0.5 mL) was added DMPU/HBr (60 wt/wt%) (54 mg, 0.4 mmol) at 0 °C, then the mixture was warmed to room temperature and was

stirred for 8 h. After completion, the solvent was evaporated under the reduced pressure and the residue was purified by silica gel column chromatography (hexanes/EtOAc as elute solvent) to afford the corresponding products **5**.

(E)-*N*-(1-bromo-2-phenylvinyl)-*N*-4-methylbenzenesulfonamide (**5a**).¹ Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), white solid, 72.4 mg, 82% yield. Mp 142–143°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.86 (d, *J* = 8.1 Hz, 2H), 7.34 (d, *J* = 8.1 Hz, 2H), 7.33 – 7.27 (m, 2H), 7.24 (dd, *J* = 6.2, 1.9 Hz, 2H), 7.23 – 7.04 (m, 6H), 6.81 (s, 1H), 4.83 (d, *J* = 13.1 Hz, 1H), 3.94 (d, *J* = 13.1 Hz, 1H), 2.46 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.7, 139.7, 134.3, 133.6, 133.2, 129.9, 129.6, 129.2, 128.7, 128.6, 128.3, 128.1, 128.0, 119.6, 53.3, 21.7.

(E)-*N*-(1-bromo-2-phenylvinyl)-*N*-heptyl-4-methylbenzenesulfonamide (**5b**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:25), pale yellow oil, 85 mg, 86% yield. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.82 (d, *J* = 8.2 Hz, 2H), 7.76 – 7.61 (m, 2H), 7.66 – 7.23 (m, 5H), 6.98 (s, 1H), 3.62 – 3.59 (m, 1H), 2.89 – 2.87 (m, 1H), 2.44 (s, 3H), 1.77 – 1.35 (m, 2H), 1.35 – 0.93 (m, 8H), 0.82 (t, *J* = 7.0 Hz, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.6, 139.0, 134.0, 133.8, 129.4, 129.2, 129.0, 120.2, 49.7, 31.6, 28.8, 27.4, 26.9, 22.5, 21.7, 14.0. HRMS (ESI-FT-ICR) calcd. for C₂₂H₂₉BrNO₂S [M+H]⁺: 450.1102, found: 450.1099.

(E)-*N*-(1-bromo-2-phenylvinyl)-*N*-methylmethanesulfonamide (**5c**).¹⁰ Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), pale yellow solid, 49 mg, 85% yield. Mp 90–92°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.8 – 7.4 (m, 2H), 7.36 – 7.31 (m, 3H), 6.9 (s, 1H), 3.1 (s, 3H), 3.0 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 137.3, 133.4, 129.2, 128.8, 128.6, 120.5, 37.4, 36.5. HRMS (ESI-FT-ICR) calcd. for C₁₀H₁₃BrNO₂S [M+H]⁺: 289.9850, found: 289.9842.

(E)-*N*-(1-bromo-2-phenylvinyl)-*N*-phenylmethanesulfonamide (**5d**).¹⁰ White solid, 58 mg, 83% yield. Mp 136–137°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.03 – 7.36 (m, 4H), 7.54 – 7.14 (m, 6H), 7.06 (s, 1H), 3.14 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 138.9, 138.5, 133.3, 128.7, 128.6, 127.4, 123.7, 117.9, 39.0. HRMS (ESI-FT-ICR) calcd. for C₁₅H₁₅BrNO₂S [M+H]⁺: 352.0007, found: 352.0001.

(E)-*N*-(1-bromo-2-(2-chlorophenyl)vinyl)-*N*,4-dimethylbenzenesulfonamide (**5e**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow solid, 62 mg, 78% yield. Mp 82–83°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.80 (dd, *J* = 6.7, 2.8 Hz, 1H), 7.71 (dd, *J* = 8.4, 1.6 Hz, 2H), 7.43 – 7.29 (m, 1H), 7.26 (dd, *J* = 7.2, 3.4 Hz, 4H), 7.17 (s, 1H), 2.95 (s, 3H), 2.42 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.6, 133.5, 133.4, 133.3, 132.3, 129.6, 129.5, 126.9, 123.5, 36.9, 21.6. HRMS (ESI⁺) calcd. for C₁₆H₁₆BrClNO₂S [M+H]⁺: 399.9774, found: 399.9770.

(E)-*N*-(1-bromo-2-(3-chlorophenyl)vinyl)-*N*,4-dimethylbenzenesulfonamide (**5f**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), white solid, 65.2 mg, 82% yield. Mp 113–114°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.83 – 7.67 (m, 2H), 7.50 (s, 1H), 7.46 (s, 1H), 7.28 (t, *J* = 6.5 Hz, 4H), 6.79 (d, *J* = 2.0 Hz, 1H), 2.98 (s, 3H), 2.43 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.6, 133.5, 133.4, 133.5, 132.3, 129.6, 129.5, 126.9, 123.5, 36.9, 21.6.

HRMS (ESI-FT-ICR) calcd. for C₁₆H₁₆BrClNO₂S [M+H]⁺: 399.9774, found: 399.9766.

(E)-*N*-(1-bromo-2-(4-chlorophenyl)vinyl)-*N*,4-dimethylbenzenesulfonamide (**5g**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow solid, 62.8 mg, 79% yield. Mp 142–143°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.76 (d, *J* = 8.3 Hz, 2H), 7.52 (d, *J* = 8.6 Hz, 2H), 7.33 – 7.27 (m, 4H), 6.59 (s, 1H), 3.02 (s, 3H), 2.44 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.6, 134.5, 134.1, 131.6, 131.1, 129.9, 129.5, 128.9, 128.7, 35.7, 21.6. HRMS (ESI-FT-ICR) calcd. for C₁₆H₁₆BrClNO₂S [M+H]⁺: 399.9774, found: 399.9764.

(E)-*N*-(1-bromo-2-(4-fluorophenyl)vinyl)-*N*,4-dimethylbenzenesulfonamide (**5h**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), white solid, 62.6 mg, 82% yield. Mp 86–88°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.77 (d, *J* = 7.7 Hz, 2H), 7.67 – 7.51 (m, 2H), 7.30 (d, *J* = 7.9 Hz, 2H), 7.03 (t, *J* = 8.3 Hz, 2H), 6.81 (s, 1H), 2.97 (s, 3H), 2.43 (s, 3H). ¹⁹F NMR (376 MHz, Chloroform-*d*) δ -111.2 (dq, *J* = 9.0, 5.0, 4.3 Hz). ¹³C{¹H} NMR (100 MHz, Chloroform-*d*) δ 162.7 (d, *J* = 250.0 Hz), 144.7, 135.8, 133.5, 130.7, 130.6, 129.5, 129.0, 121.1, 115.7 (d, *J* = 21.7 Hz), 36.6, 21.6. HRMS (ESI-FT-ICR) calcd. for C₁₆H₁₆BrFNO₂S [M+H]⁺: 384.0069, found: 384.0061.

(E)-*N*-(1-bromo-2-(4-bromophenyl)vinyl)-*N*-phenylmethanesulfonamide (**5i**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), white solid, 69 mg, 81% yield. Mp 121–122°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.53 (d, *J* = 8.1 Hz, 2H), 7.44 (q, *J* = 8.4 Hz, 4H), 7.35 (t, *J* = 7.6 Hz, 2H), 7.29 (d, *J* = 6.6 Hz, 2H), 6.99 (s, 1H), 3.14 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 138.7, 137.3, 132.2, 131.9, 130.1, 129.6, 127.6, 123.6, 123.3, 118.6, 39.0. HRMS (ESI-FT-ICR) calcd. for C₁₅H₁₄Br₂NO₂S [M+H]⁺: 429.9107, found: 429.9100.

(E)-*N*-(1-bromo-2-(4-cyanophenyl)vinyl)-*N*,4-dimethylbenzenesulfonamide (**5j**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), colorless oil, 64 mg, 82% yield. Mp 131–134°C. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.75 (d, *J* = 7.7 Hz, 2H), 7.68 (d, *J* = 8.0 Hz, 2H), 7.62 (d, *J* = 7.9 Hz, 2H), 7.31 (d, *J* = 7.8 Hz, 2H), 6.87 (s, 1H), 2.98 (s, 3H), 2.44 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 145.1, 138.2, 135.1, 133.1, 132.4, 129.6, 129.1, 129.0, 124.9, 118.6, 112.0, 36.8, 21.7. HRMS (ESI-FT-ICR) calcd. for C₁₇H₁₆BrN₂O₂S [M+H]⁺: 391.0116, found: 391.0106.

Methyl (E)-3-(2-bromo-2-(*N*-phenylmethylsulfonamido)vinyl)benzoate (**5k**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), colorless oil, 52.4 mg, 64% yield. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.15 (s, 1H), 7.97 (d, *J* = 6.8 Hz, 1H), 7.85 (d, *J* = 8.1 Hz, 1H), 7.57 (d, *J* = 8.1 Hz, 2H), 7.43 (t, *J* = 7.8 Hz, 1H), 7.34 (t, *J* = 7.6 Hz, 2H), 7.28 (d, *J* = 7.1 Hz, 1H), 7.08 (s, 1H), 3.90 (s, 3H), 3.13 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 166.5, 138.9, 137.2, 133.6, 132.5, 130.5, 130.0, 129.9, 129.6, 128.9, 127.7, 124.0, 119.3, 52.2, 39.0. HRMS (ESI-FT-ICR) calcd. for C₁₇H₁₇BrNO₄S [M+H]⁺: 410.0062, found: 410.0054.

6. Hydrochlorination of ynamides **4**

To a solution of ynamides **4** (0.2 mmol) in DCM (0.5 mL) was added DMPU/HCl (43 wt/wt%) (34 mg, 0.4 mmol) at 0 °C,

then the mixture was warmed to room temperature and was stirred for 8 h. After completion, the solvent was evaporated under the reduced pressure and the residue was purified by silica gel column chromatography (hexanes/EtOAc as elute solvent) to afford the corresponding products **6**.

(E)-N-(1-chloro-2-phenylvinyl)-N-methylmethanesulfonamide (6a).¹⁰ Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), white solid, 46.5 mg, 95% yield. ¹H NMR (400 MHz, Chloroform-d) δ 7.56 (d, J = 7.0 Hz, 2H), 7.34 – 7.30 (m, 3H), 6.68 (s, 1H), 3.13 (s, 3H), 3.00 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 132.7, 132.5, 129.1, 128.8, 128.6, 38.1, 35.7. HRMS (ESI-FT-ICR) calcd. for C₁₀H₁₃ClNO₂S [M+H]⁺: 246.0356, found: 246.0350.

(E)-N-(1-chloro-2-phenylvinyl)-N-heptyl-4-methylbenzenesulfonamide (6b). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), colorless oil, 72.9 mg, 85% yield. ¹H NMR (400 MHz, Chloroform-d) δ 7.81 (d, J = 8.3 Hz, 2H), 7.70 (d, J = 7.1 Hz, 2H), 7.34 – 7.30 (m, 5H), 6.77 (s, 1H), 3.56 (s, 1H), 3.01 (s, 1H), 2.43 (s, 3H), 1.53 (bs, 2H), 1.55 – 0.94 (m, 8H), 0.82 (t, J = 7.0 Hz, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.4, 134.7, 134.4, 133.1, 129.5, 129.1, 129.0, 128.9, 128.5, 127.9, 48.6, 31.6, 28.8, 27.3, 26.8, 22.4, 21.6, 14.0. HRMS (ESI-FT-ICR) calcd. for C₂₂H₂₈ClNO₂S [M+H]⁺: 406.1608, found: 406.1599.

(E)-N-benzyl-N-(1-chloro-2-phenylvinyl)-4-methylbenzenesulfonamide (6c). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), white solid, 74.6 mg, 94% yield. Mp 127–129°C. ¹H NMR (400 MHz, Chloroform-d) δ 7.86 (d, J = 8.3 Hz, 2H), 7.42 – 7.28 (m, 4H), 7.24 – 7.00 (m, 8H), 6.60 (s, 1H), 4.80 (d, J = 12.3 Hz, 1H), 4.04 (d, J = 12.3 Hz, 1H), 2.46 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.6, 135.0, 134.9, 133.3, 132.8, 129.8, 129.6, 128.9, 128.7, 128.5, 128.3, 128.2, 128.0, 127.5, 52.4, 21.7. HRMS (ESI-FT-ICR) calcd. for C₂₂H₂₁ClNO₂S [M+H]⁺: 398.0982, found: 398.0980.

(E)-N-(1-chloro-2-phenylvinyl)-N-phenylmethanesulfonamide (6d).¹⁰ Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), pale yellow liquid, 60.1 mg, 95% yield. ¹H NMR (400 MHz, Chloroform-d) δ 7.83 – 7.41 (m, 4H), 7.46 – 7.07 (m, 6H), 6.84 (s, 1H), 3.13 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 138.8, 133.6, 132.6, 129.5, 129.1, 128.7, 128.6, 128.1, 127.6, 124.3, 39.6. HRMS (ESI-FT-ICR) calcd. for C₁₅H₁₅ClNO₂S [M+H]⁺: 308.0512, found: 308.0506.

(E)-N-allyl-N-(1-chloro-2-phenylvinyl)-4-methylbenzenesulfonamide (6e).¹⁰ Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), yellow solid, 66 mg, 95% yield. ¹H NMR (400 MHz, Chloroform-d) δ 7.81 (d, J = 8.3 Hz, 2H), 7.67 (d, J = 7.1 Hz, 2H), 7.50 – 7.18 (m, 5H), 6.74 (s, 1H), 5.72 – 5.71 (m, 1H), 5.24 – 5.23 (m, 1H), 5.12 – 5.10 (m, 1H), 4.28 (s, 1H), 3.64 (s, 1H), 2.44 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.5, 134.7, 134.4, 133.2, 130.5, 129.5, 129.0, 128.7, 128.8, 128.5, 127.6, 120.9, 51.5, 21.6. HRMS (ESI-FT-ICR) calcd. for C₁₈H₁₉ClNO₂S [M+H]⁺: 348.0825, found: 348.0817.

(E)-N-(1-chloro-2-phenylvinyl)-N-(2-(cyclohex-1-en-1-yl)ethyl)-4-methylbenzenesulfonamide (6f). Purified by flash column chromatography (ethyl acetate / hexanes = 1:25), yellow oil, 68 mg, 82% yield. ¹H NMR (400 MHz, Chloroform-d)

δ 7.81 (d, J = 8.3 Hz, 2H), 7.69 (d, J = 7.4 Hz, 2H), 7.43 – 7.26 (m, 5H), 6.76 (s, 1H), 5.33 (s, 1H), 3.71 (s, 1H), 3.05 (s, 1H), 2.43 (s, 3H), 2.18 – 2.15 (m, 2H), 1.86 – 1.83 (m, 4H), 1.52 – 1.49 (m, 4H). ¹³C{¹H} NMR (100 MHz,) δ 144.4, 134.7, 134.5, 133.4, 133.1, 129.5, 129.3, 128.8, 128.4, 127.6, 123.9, 47.1, 35.6, 28.1, 25.1, 22.7, 22.1, 21.6. HRMS (ESI⁺) calcd. for C₂₃H₂₇ClNO₂S [M+H]⁺: 416.1446, found: 416.1441.

(E)-3-(1-chloro-2-phenylvinyl)oxazolidin-2-one (6g). Purified by flash column chromatography (ethyl acetate / hexanes = 1:10), white solid, 41 mg, 92% yield. Mp 124–126°C. ¹H NMR (400 MHz, Chloroform-d) δ 7.31 (s, 5H), 6.73 (s, 1H), 4.42 (t, J = 7.8 Hz, 2H), 3.76 (t, J = 7.9 Hz, 2H). ¹³C{¹H} NMR (100 MHz,) δ 155.3, 132.9, 130.4, 128.8, 128.7, 127.9, 125.9, 62.8, 44.5. HRMS (ESI-FT-ICR) calcd. for C₁₁H₁₁ClNO₂ [M]⁺: 224.0448, found: 224.0441.

(E)-N-(1-chloro-2-(naphthalen-1-yl)vinyl)-N-methylmethanesulfonamide (6h). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), pale yellow liquid, 53 mg, 90% yield. ¹H NMR (400 MHz, Chloroform-d) δ 7.94 (d, J = 7.6 Hz, 1H), 7.84 (t, J = 9.2 Hz, 2H), 7.76 (d, J = 7.2 Hz, 1H), 7.62 – 7.43 (m, 3H), 7.32 (s, 1H), 3.02 (s, 3H), 2.81 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 133.4, 131.5, 131.2, 130.1, 130.0, 129.1, 128.7, 126.6, 126.5, 126.1, 125.6, 123.8, 38.5, 36.3. HRMS (ESI-FT-ICR) calcd. for C₁₄H₁₅ClNO₂S [M+H]⁺: 296.0512, found: 296.0506.

(E)-N-(1-chloro-2-(4-fluorophenyl)vinyl)-N,4-dimethylbenzenesulfonamide (6i).²⁴ Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), pale yellow solid, 63.7 mg, 94% yield. ¹H NMR (400 MHz, Chloroform-d) δ 7.77 (d, J = 8.1 Hz, 2H), 7.64 – 7.54 (m, 2H), 7.30 (d, J = 7.9 Hz, 2H), 7.09 – 6.99 (m, 2H), 6.60 (s, 1H), 3.01 (s, 3H), 2.43 (s, 3H). ¹⁹F NMR (376 MHz, Chloroform-d) δ -111.5 – -111.6 (m). ¹³C{¹H} NMR (100 MHz, Chloroform-d) δ 162.7 (d, J = 249.7 Hz), 144.6, 134.1, 131.1, 130.6 (d, J = 8.3 Hz), 129.5, 129.3, 128.8, 115.7 (d, J = 21.7 Hz), 35.7, 21.6.

(E)-N-(1-chloro-2-(4-chlorophenyl)vinyl)-N,4-dimethylbenzenesulfonamide (6j). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), white solid, 61.7 mg, 87% yield. Mp 108–110°C. ¹H NMR (400 MHz, Chloroform-d) δ 7.76 (d, J = 7.9 Hz, 2H), 7.52 (d, J = 7.5 Hz, 2H), 7.30 (d, J = 7.6 Hz, 4H), 6.80 (s, 1H), 2.97 (s, 3H), 2.44 (s, 3H). ¹³C{¹H} NMR (100 MHz,) δ 144.8, 135.7, 134.6, 133.5, 132.3, 129.9, 129.5, 129.0, 128.9, 122.0, 36.7, 21.6. HRMS (ESI-FT-ICR) calcd. for C₁₆H₁₆Cl₂NO₂S [M+H]⁺: 356.0279, found: 356.0272.

(E)-N-(1-chloro-2-(4-methoxyphenyl)vinyl)-N-methylmethanesulfonamide (6k). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), white solid, 50.6 mg, 92% yield. Mp 125–127°C. ¹H NMR (500 MHz, Chloroform-d) δ 7.54 (d, J = 8.7 Hz, 2H), 6.89 (d, J = 8.7 Hz, 2H), 6.62 (s, 1H), 3.81 (s, 3H), 3.14 (s, 3H), 3.03 (s, 3H). ¹³C{¹H} NMR (125 MHz,) δ 160.1, 132.0, 130.2, 126.9, 125.4, 114.1, 55.3, 38.0, 35.6. HRMS (ESI-FT-ICR) calcd. for C₁₁H₁₅ClNO₃S [M+H]⁺: 276.0461, found: 276.0453.

(E)-N-(1-chloro-2-(4-nitrophenyl)vinyl)-N,4-dimethylbenzenesulfonamide (6l). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), white solid, 64.4 mg, 88% yield. Mp 154–157°C. ¹H NMR (500 MHz,

Chloroform-*d*) δ 8.21 (d, J = 8.9 Hz, 2H), 7.77 (d, J = 7.5 Hz, 4H), 7.33 (d, J = 8.4 Hz, 2H), 3.06 (s, 3H), 2.46 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz,) δ 147.3, 145.0, 139.6, 133.7, 133.6, 130.1, 129.7, 129.4, 128.7, 123.9, 35.8, 21.6. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{16}\text{H}_{16}\text{ClN}_2\text{O}_4\text{S}$ $[\text{M}+\text{H}]^+$: 367.0519, found 367.0512.

(*E*)-*N*-(2-(4-acetylphenyl)-1-chlorovinyl)-*N*,4-dimethylbenzenesulfonamide (**6m**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:10), white solid, 66 mg, 91% yield. Mp 142–143°C. ^1H NMR (400 MHz, Chloroform-*d*) δ 7.93 (d, J = 8.4 Hz, 2H), 7.75 (d, J = 8.1 Hz, 2H), 7.67 (d, J = 7.9 Hz, 2H), 7.30 (d, J = 7.8 Hz, 2H), 6.68 (s, 1H), 3.03 (s, 3H), 2.59 (s, 3H), 2.42 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz,) δ 197.3, 144.7, 137.7, 136.7, 133.9, 132.1, 131.2, 129.6, 128.8, 128.7, 128.6, 35.6, 26.6, 21.6. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{18}\text{H}_{19}\text{ClNO}_3\text{S}$ $[\text{M}+\text{H}]^+$: 364.0774, found: 364.0766.

(*E*)-*N*-(1-chloro-2-(thiophen-3-yl)vinyl)-*N*,4-dimethylbenzenesulfonamide (**6n**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:15), white solid, 58.8 mg, 90% yield. Mp 139–141°C. ^1H NMR (500 MHz, Chloroform-*d*) δ 7.84 (d, J = 6.8 Hz, 2H), 7.38 (d, J = 4.4 Hz, 1H), 7.34 (d, J = 7.7 Hz, 2H), 7.15 (d, J = 3.5 Hz, 1H), 7.00 (t, J = 3.9 Hz, 1H), 6.88 (s, 1H), 3.05 (s, 3H), 2.46 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz,) δ 144.6, 135.9, 134.1, 129.9, 129.5, 129.1, 128.3, 127.0, 126.7, 126.4, 35.5, 21.7. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{14}\text{H}_{15}\text{ClNO}_2\text{S}_2$ $[\text{M}+\text{H}]^+$: 328.0233, found: 328.0225.

(3*S*,10*R*,13*R*,17*R*)-10,13-dimethyl-17-((*R*)-6-methylheptan-2-yl)-2,3,4,7,8,9,10,11,12,13,14,15,16,17-tetradecahydro-1*H*-cyclopenta[*a*]phenanthren-3-yl (E)-6-chloro-6-((*N*,4-dimethylphenyl)sulfonamido)hex-5-enoate (**6o**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:20), white solid, 114 mg, 82% yield. Mp 88–89°C. ^1H NMR (500 MHz, Chloroform-*d*) δ 7.76 (d, J = 8.2 Hz, 2H), 7.32 (d, J = 8.0 Hz, 2H), 5.79 (t, J = 7.5 Hz, 1H), 5.38 (d, J = 5.1 Hz, 1H), 4.64 (dq, J = 11.9, 7.0, 5.6 Hz, 1H), 2.92 (s, 3H), 2.44 (s, 3H), 2.36 – 2.32 (m, 6H), 2.05 – 1.93 (m, 2H), 1.92 – 1.79 (m, 3H), 1.68 – 1.06 (m, 23H), 1.03 (s, 3H), 0.92 (d, J = 6.5 Hz, 3H), 0.88 (d, J = 2.3 Hz, 3H), 0.87 (d, J = 2.3 Hz, 3H), 0.68 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz,) δ 172.6, 144.3, 139.7, 134.3, 134.0, 129.5, 129.0, 128.5, 122.6, 74.0, 56.7, 56.1, 50.0, 42.3, 39.7, 39.5, 38.1, 37.0, 36.6, 36.2, 35.9, 35.8, 34.1, 31.9, 31.9, 28.6, 28.2, 28.0, 27.8, 24.3, 23.9, 23.8, 22.8, 22.6, 21.6, 21.0, 19.3, 18.7. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{41}\text{H}_{63}\text{ClNO}_4\text{S}$ $[\text{M}+\text{H}]^+$: 700.4166, found: 700.4158.

7. Further transformations of halogenated alkene products

A mixture of **2a** (0.2 mmol), phenylboronic acid (36.5 mg, 0.3 mmol), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (7 mg, 5 mol%) and Cs_2CO_3 (652 mg, 2 mmol) was added 1,4-dioxane/ H_2O (4/1, 2 mL) heated at 60 °C for 8 h. After the completion of the reaction, the reaction mixture extracted with EtOAc, the combined organic layers dried over Na_2SO_4 and removal of the solvent by rotary evaporation the resulting residue was purified by chromatography on silica gel to give the desired products.

4,4-Diphenylbut-3-en-2-one (**7a**). 25 ^1H NMR (400 MHz, Chloroform-*d*) δ 7.50 – 7.36 (m, 3H), 7.39 – 7.27 (m, 5H), 7.23 –

7.18 (m, 1H), 6.58 (s, 1H), 1.88 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, Chloroform-*d*) δ 200.2, 154.0, 140.8, 138.9, 129.6, 129.5, 128.8, 128.4, 128.4, 127.7, 30.3.

A mixture of **4a** (73.5 mg, 0.3 mmol), phenylboronic acid (51.2 mg, 4.2 mmol), $\text{Pd}(\text{OAc})_2$ (7 mg, 10 mol%) and K_2CO_3 (82.8 mg, 0.6 mmol) in EtOH/ H_2O (4/1) (2 mL) was heated at 80 °C for 8 h. After the completion of the reaction, the reaction mixture was filtered through a pad of celite and removal of the solvent in vacuum to give the crude product. The resulting residue was purified by silica gel chromatography to give the desired products.

(*Z*)-*N*-(1,2-diphenylvinyl)-*N*-methylmethanesulfonamide (**Z-8a**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:40), colorless oil, 40.2 mg, 47% yield. ^1H NMR (500 MHz, Chloroform-*d*) δ 7.57 (d, J = 8.8 Hz, 2H), 7.52 (d, J = 7.2 Hz, 2H), 7.45 – 7.41 (m, 5H), 7.33 (d, J = 7.4 Hz, 1H), 6.83 (s, 1H), 3.12 (s, 3H), 2.50 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz,) δ 139.6, 137.7, 135.7, 128.9, 128.8, 128.7, 128.7, 128.1, 127.0, 40.0, 37.1. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{16}\text{H}_{17}\text{NO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 288.1058, found: 288.1050.

(*E*)-*N*-(1,2-diphenylvinyl)-*N*-methylmethanesulfonamide (**E-8a**). Purified by flash column chromatography (ethyl acetate / hexanes = 1:40), white solid, 37.2 mg, 43% yield. Mp 113–115°C ^1H NMR (500 MHz, Chloroform-*d*) δ 7.37 (dd, J = 7.1, 2.8 Hz, 2H), 7.32 (dd, J = 5.2, 1.9 Hz, 3H), 7.16 – 7.12 (m, 3H), 7.06 – 6.98 (m, 2H), 6.75 (s, 1H), 3.13 (s, 3H), 2.99 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz,) δ 140.0, 135.4, 135.2, 129.8, 129.2, 128.8, 128.7, 128.1, 127.5, 127.3, 39.2, 37.1. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{16}\text{H}_{18}\text{NO}_2\text{S}$ $[\text{M}+\text{H}]^+$: 288.1058, found: 288.1051.

A mixture of **5c** (87 mg, 0.3 mmol), ethynyltrimethylsilane (41.2 mg, 4.2 mmol), $\text{Pd}(\text{PPh}_3)\text{Cl}_2$ (10 mg, 5 mol%), and CuI (5 mg, 8 mol%) in $\text{Et}_3\text{N}/\text{THF}$ (1/1; 1.5 mL) was stirred at room temperature for 8 h. After the completion of the reaction, the reaction mixture was filtered through celite and removal of the solvent in vacuum gave the crude product. The crude product was purified by silica gel chromatography (ethyl acetate / hexanes = 1:30) to give the desired product **9a** as a pale-yellow oil (87 mg, 95% yield). Major isomer: ^1H NMR (400 MHz, Chloroform-*d*) δ 7.84 (dd, J = 7.6, 2.0 Hz, 2H), 7.42 – 7.27 (m, 3H), 7.05 (s, 1H), 3.12 (s, 3H), 2.94 (s, 3H), 0.26 (s, 9H). $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz,) δ 140.3, 133.5, 129.5, 129.3, 128.3, 118.4, 104.3, 99.0, 35.7, -0.4. Minor isomer: ^1H NMR (400 MHz, Chloroform-*d*) δ 7.70 (d, J = 8.1 Hz, 2H), 6.80 (s, 1H), 3.05 (s, 3H), 3.01 (s, 3H), 0.23 (s, 8H). $^{13}\text{C}\{^1\text{H}\}$ NMR (126 MHz,) δ 139.0, 133.6, 129.9, 128.7, 119.9, 101.0, 97.1, 36.4, -0.3. HRMS (ESI-FT-ICR) calcd. for $\text{C}_{15}\text{H}_{22}\text{NO}_2\text{SSi}$ $[\text{M}+\text{H}]^+$: 308.1141, found: 308.1133.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/

Copies of NMR spectra (PDF).

AUTHOR INFORMATION

Corresponding Author

* Email: bo.xu@dhru.edu.cn;
gb.hammond@louisville.edu

ACKNOWLEDGMENT

We are grateful to the National Science Foundation of China for financial support (NSFC-21672035) and China Recruitment Program of Global Experts for financial support. G.B.H. is grateful to the National Institutes of Health (1R01GM121660). X.Z is grateful to the China Scholarship Council for financial support.

REFERENCES

- (a) Johansson Seechurn, C. C. C.; Kitching, M. O.; Colacot, T. J.; Snieckus, V. *Angew. Chem. Int. Ed.* **2012**, *51*, 5062-5085; (b) Iwai, T.; Fujihara, T.; Terao, J.; Tsuji, Y. *J. Am. Chem. Soc.* **2009**, *131*, 6668-6669; (c) Petrone, D. A.; Franzoni, I.; Ye, J.; Rodriguez, J. F.; Poblador-Bahamonde, A. I.; Lautens, M. J. *Am. Chem. Soc.* **2017**, *139*, 3546-3557; (d) Uehling, M. R.; Rucker, R. P.; Lalic, G. *J. Am. Chem. Soc.* **2014**, *136*, 8799-8803; (e) Yamaguchi, R.; Kawasaki, H.; Kawanisi, M. *Synth. Commun.* **1982**, *12*, 1027-1037; (f) Zhu, G.; Chen, D.; Wang, Y.; Zheng, R. *Chem. Commun.* **2012**, *48*, 5796-5798; (g) Dérien, S.; Klein, H.; Bruneau, C. *Angew. Chem. Int. Ed.* **2015**, *54*, 12112-12115.
- (a) Baran, P. S.; Shenvi, R. A.; Mitsos, C. A. *Angew. Chem. Int. Ed.* **2005**, *44*, 3714-3717; (b) Chevolut, L.; Chevolut, A. M.; Gajhede, M.; Larsen, C.; Anthoni, U.; Christophersen, C. *J. Am. Chem. Soc.* **1985**, *107*, 4542-4543; (c) Sun, C.; Camp, J. E.; Weinreb, S. M. *Org. Lett.* **2006**, *8*, 1779-1781.
- (a) Weiss, H. M. *J. Chem. Educ.* **1995**, *72*, 848; (b) Podhajsky, S. M.; Sigman, M. S. *Organometallics* **2007**, *26*, 5680-5686; (c) Xu, C.-X.; Ma, C.-H.; Xiao, F.-R.; Chen, H.-W.; Dai, B. *Chin. Chem. Lett.* **2016**, *27*, 1683-1685; (d) Oliver-Meseguer, J.; Doménech-Carbó, A.; Boronat, M.; Leyva-Pérez, A.; Corma, A. *Angew. Chem. Int. Ed.* **2017**, *56*, 6435-6439; (e) Wang, B.; Wang, S.; Li, P.; Wang, L. *Chem. Commun.* **2010**, *46*, 5891-5893; (f) Iwai, T.; Fujihara, T.; Terao, J.; Tsuji, Y. *J. Am. Chem. Soc.* **2012**, *134*, 1268-1274; (g) Schevenels, F. T.; Shen, M.; Snyder, S. A. *J. Am. Chem. Soc.* **2017**, *139*, 6329-6337.
- (a) Chelpanova, L. F.; Mashlyakovskii, L. N. *Zh. Obshch. Khim.* **1966**, *2*, 602-7; (b) Kerr, P. J.; Pyke, S. M.; Ward, A. D. *Aust. J. Chem.* **2008**, *61*, 350-358; (c) Shchukin, A. O.; Vasilyev, A. V. *Appl. Catal., A* **2008**, *336*, 140-147; (d) Semenova, I. S.; Yarovenko, V. N.; Levchenko, K. S.; Krayushkin, M. M. *Russ. Chem. Bull.* **2013**, *62*, 1022-1025; (e) Yan, W.; Li, Z.; Kishi, Y. *J. Am. Chem. Soc.* **2015**, *137*, 6219-6225.
- (a) Kundu, N. G.; Choudhuri, L. N.; Das, B. *J. Chem. Soc., Perkin Trans. 1* **1989**, 1165-6; (b) Kundu, N. G.; Chaudhuri, L. N. *J. Chem. Soc., Perkin Trans. 1* **1991**, 1677-82.
- Yang, F.; Ji, K.-G.; Ali, S.; Liang, Y.-M. *J. Org. Chem.* **2011**, *76*, 8329-8335.
- Mulder, J. A.; Kurtz, K. C. M.; Hsung, R. P.; Coverdale, H.; Frederick, M. O.; Shen, L.; Zifcick, C. A. *Org. Lett.* **2003**, *5*, 1547-1550.
- (a) Ohashi, K.; Mihara, S.; Sato, A. H.; Ide, M.; Iwasawa, T. *Tetrahedron Lett.* **2014**, *55*, 632-635; (b) Sato, A. H.; Ohashi, K.; Iwasawa, T. *Tetrahedron Lett.* **2013**, *54*, 1309-1311.
- Prabagar, B.; Nayak, S.; Mallick, R. K.; Prasad, R.; Sahoo, A. K. *Org. Chem. Front.* **2016**, *3*, 110-115.
- Kim, S. W.; Um, T.-W.; Shin, S. *Chem. Commun.* **2017**, 53, 2733-2736.
- (a) Compain, G.; Jouvin, K.; Martin-Mingot, A.; Evano, G.; Marrot, J.; Thibaudeau, S. *Chem. Commun.* **2012**, *48*, 5196-5198; (b) Métayer, B.; Compain, G.; Jouvin, K.; Martin-Mingot, A.; Bachmann, C.; Marrot, J.; Evano, G.; Thibaudeau, S. *J. Org. Chem.* **2015**, *80*, 3397-3410.
- Zeng, X.; Liu, S.; Shi, Z.; Xu, B. *Org. Lett.* **2016**, *18*, 4770-4773.
- (a) Okoromoba, O. E.; Hammond, G. B.; Xu, B. *Org. Lett.* **2015**, *17*, 3975-3977; (b) Okoromoba, O. E.; Han, J.; Hammond, G. B.; Xu, B. *J. Am. Chem. Soc.* **2014**, *136*, 14381-14384; (c) Okoromoba, O. E.; Li, Z.; Robertson, N.; Mashuta, M. S.; Couto, U. R.; Tormena, C. F.; Xu, B.; Hammond, G. B. *Chem. Commun.* **2016**, *52*, 13353-13356.
- Laurence, C.; Brameld, K. A.; Graton, J.; Le Questel, J.-Y.; Renault, E. *J. Med. Chem.* **2009**, *52*, 4073-4086.
- (a) Liang, S.; Ebule, R.; Hammond, G. B.; Xu, B. *Org. Lett.* **2017**, *19*, 4524-4527; (b) Li, Z.; Ebule, R.; Kostyo, J.; Hammond, G. B.; Xu, B. *Chem. Eur. J.* **2017**, *23*, 12739-12743.
- (a) DeKorver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. *Chem. Rev.* **2010**, *110*, 5064-5106; (b) Evano, G.; Coste, A.; Jouvin, K. *Angew. Chem. Int. Ed.* **2010**, *49*, 2840-2859; (c) Wang, X.-N.; Yeom, H.-S.; Fang, L.-C.; He, S.; Ma, Z.-X.; Kedrowski, B. L.; Hsung, R. P. *Acc. Chem. Res.* **2014**, *47*, 560-578; (d) Hu, L.; Xu, S.; Zhao, Z.; Yang, Y.; Peng, Z.; Yang, M.; Wang, C.; Zhao, J. *J. Am. Chem. Soc.* **2016**, *138*, 13135-13138; (e) Xu, S.; Liu, J.; Hu, D.; Bi, X. *Green Chem.* **2015**, *17*, 184-187; (f) Zhang, Y. *Tetrahedron* **2006**, *62*, 3917-3927; (g) Wu, W.; Jiang, H. *Acc. Chem. Res.* **2014**, *47*, 2483-2504; (h) Li, J.; Yang, W.; Yang, S.; Huang, L.; Wu, W.; Sun, Y.; Jiang, H. *Angew. Chem. Int. Ed.* **2014**, *53*, 7219-7222; (i) Li, Y.; Liu, X.; Jiang, H.; Liu, B.; Chen, Z.; Zhou, P. *Angew. Chem. Int. Ed.* **2011**, *50*, 6341-6345; (j) Li, J.; Yang, S.; Huang, L.; Chen, H.; Jiang, H. *RSC Adv.* **2013**, *3*, 11529-11532.
- (a) Stang, P. J.; Summerville, R. *J. Am. Chem. Soc.* **1969**, *91*, 4600-1; (b) Zhang, S.-L.; Wan, H.-X.; Deng, Z.-Q. *Org. Biomol. Chem.* **2017**, *15*, 6367-6374.
- (a) Carey, F. A.; Sundberg, R. J., *Advanced Organic Chemistry, Part A: Structure and Mechanisms*. In *Advanced Organic Chemistry, Part A: Structure and Mechanisms*, 5th ed.; Springer: 2007; Vol. A, pp 536-540; (b) Ashtekar, K. D.; Veticatt, M.; Yousefi, R.; Jackson, J. E.; Borhan, B. *J. Am. Chem. Soc.* **2016**, *138*, 8114-8119.
- (a) Weiss, H. M.; Touchette, K. M.; Andersen, F.; Iskhakov, D. *Org. Biomol. Chem.* **2003**, *1*, 2148; (b) Weiss, H. M.; Touchette, K. M.; Angell, S.; Khan, J. *Org. Biomol. Chem.* **2003**, *1*, 2152; (c) Weiss, H. M.; Touchette, K. M. *J. Chem. Soc., Perkin Trans. 2* **1998**, 1523-1528.
- Zeng, X.; Lu, Z.; Liu, S.; Hammond, G. B.; Xu, B. *Adv. Synth. Catal.*, n/a-n/a.
- Doyle, M. P.; Devia, A. H.; Bassett, K. E.; Terpstra, J. W.; Mahapatro, S. N. *J. Org. Chem.* **1987**, *52*, 1619-1621.
- Mona, H.-S.; Zahra, M. *Bull. Chem. Soc. Jpn.* **2011**, *84*, 778-782.
- Shchukin, A. O.; Vasilyev, A. V. *Appl. Catal., A* **2008**, *336*, 140-147.
- Prabagar, B.; Nayak, S.; Mallick, R. K.; Prasad, R.; Sahoo, A. K. *Organic Chemistry Frontiers* **2016**, *3*, 110-115.
- Tani, M.; Sakaguchi, S.; Ishii, Y. *J. Org. Chem.* **2004**, *69*, 1221-1226.

TOC

