

Subscriber access provided by NORTH CAROLINA STATE UNIV

Article

Traceless silulation of #-C(sp3)–H bonds of alcohols via perfluorinated acetals

Ala Bunescu, Trevor W Butcher, and John F. Hartwig

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.7b12150 • Publication Date (Web): 28 Dec 2017

Downloaded from http://pubs.acs.org on December 28, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

1 2

Traceless silulation of β -C(sp³)–H bonds of alcohols via perfluorinated acetals

Ala Bunescu, Trevor W. Butcher, John F. Hartwig*

Department of Chemistry, University of California, Berkeley, California 94720, USA

Supporting Information Placeholder

ABSTRACT: We report the silvlation of primary C–H bonds located β to secondary and tertiary alcohols by exploiting perfluorinated esters as traceless directing groups. The conversion of a secondary or tertiary alcohol to a perfluoroalkyl ester and conversion of the ester to the corresponding silvl acetals by hydrosilvlation allows for selective β -C(sp³)–H silvlation catalyzed by the combination of [Ir(cod)OMe]₂ and Me₄Phen (3,4,7,8-tetramethyl-1,10-phenanthroline) to form 6membered oxasilolanes. Tamao-Fleming oxidation of these oxasilolanes leads to 1,2 diols. The developed sequence was applied to a series of natural products containing hydroxyl groups.

Introduction.

Site-selective functionalization of specific C–H bonds in the presence of an array of C–H bonds and functional groups is transforming the logic by which organic molecules are synthesized and modified.¹ Among existing C–H functionalization reactions, the borylation² and silylation³ of C–H bonds are particularly valuable because the silicon and boron-based products undergo a range of cross-coupling and oxidative functionalization reactions.

The hydroxyl group is the most commonly encountered functional group in natural compounds, making alcoholdirected C–H functionalization reactions particularly important for the selective, direct modification of complex molecules.⁴ The C–H bonds located α to the hydroxyl group are the most reactive and are cleaved during radical and oxidative processes.⁵ The C–H bonds δ to an alcohol undergo functionalization by 1,5-hydrogen atom abstraction of highly energetic oxygen-centered radicals.⁶ The functionalization of alcohols at the γ and β positions are more limited.⁷

Previously, our group developed Ir-catalyzed silylations of unactivated γ -C(sp³)–H bonds in alcohols to form 1,3 diols (see Scheme 1a).⁸ In this approach, (hydrido)silyl ethers obtained by the dehydrogenative coupling of the alcohol with diethylsilane undergo functionalization of primary C(sp³)–H bonds at the position γ to the alcohol. Subsequent Tamao-Fleming oxidation of the resulting 5-membered oxasilolanes furnishes the 1,3-diol product.^{8b} However, this approach, without modification, is not applicable to the functionalization of C–H bonds located β to the alcohol because the reaction would require the formation of strained 4-membered ring oxasilolanes (see Scheme 1a).

Dong recently reported an alternative strategy that led to the β functionalization of alcohols. Pd-catalyzed β -acetoxylation⁹ and β -tosylation¹⁰ reactions of alcohols occur with an *O*-alkyl oxime as directing group (see Scheme 1b). Although providing novel selectivity, the multi-step installation and reductive removal of this directing group, along with the moderate selectivity for monofunctionalization of the many identical C–H bonds, are drawbacks of this method, particularly when functionalizing complex molecules containing an array of functional groups. Thus, we have sought alternative strategies for the silvlation of C–H bonds located β to an alcohol.

Scheme 1. Alcohol-directed C-H functionalization

a) Previous: γ -C(sp³)-H silylation of alcohols $R_{1}^{0} \xrightarrow{OH}_{R^{2}} H \xrightarrow{Ir}_{Et_{2}SIH_{2}} R_{1}^{1} \xrightarrow{O}_{R^{2}} H \xrightarrow{Ir}_{R^{2}} \xrightarrow{O}_{R^{2}} \xrightarrow{OH}_{R^{2}} \xrightarrow{$

60

To functionalize the β -C(sp³)–H bonds of alcohols, we envisioned a strategy by which an ester, converted to a silyl acetal, would serve as a readily cleaved directing group. Jeon recently used silyl acetals as directing groups for the ortho-silylation of phenols.¹¹ Our proposed strategy for the functionalization of β -C(sp³)–H bonds is outlined in Scheme 1c. Ester **2** would be accessed by a simple acylation of alcohol **1**. Hydrosilylation of ester **2** would generate the silylacetal **3**, which could undergo C(sp³)–H functionalization at the β position to form the 6membered dioxasilinane **4**. Subsequent Tamao-Fleming oxidation of the oxasilolane **4** would provide the corresponding 1,2diol.

We report the successful implementation of this strategy by which the silylation of primary C–H bonds located β to secondary and tertiary alcohols occurs with perfluorinated esters as traceless directing groups. The hydrosilylation of a perfluoroalkyl ester, subsequent *in situ* silylation of a C(sp³)–H bond, and subsequent Tamao-Fleming oxidation at the Si-C bond lead to the introduction of a hydroxyl group at the primary carbon located in the β position to an alcohol. This sequence is suitable for the derivatization of C–H bonds located β to both secondary and tertiary alcohols, including those in natural products.

Results and Discussion.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

To achieve the silylation of C–H bonds β to alcohols by this strategy (Scheme 1c), several challenges needed to be addressed. First, the silylacetal intermediate **3** must be stable during the C(sp³)–H silylation. Although an silylacetal was applied by Jeon for the functionalization of aromatic C–H bonds ortho to a phenol, an acetal that is stable to the conditions for cleavage of an unactivated, aliphatic C–H bonds is needed.¹¹ Second, a system that forms a 7-membered metalacycle intermediate by C(sp³)–H activation is needed.¹² Third, the catalyst used for ester reduction must be compatible with the catalysts used for C–H silylation if the process is to be run without isolation of intermediates. By virtue of the ester intermediate and silane reagent, the stability and reactivity of the intermediate silylacetal can be tuned by the acyl group that will be removed *in situ* during the Tamao-Fleming oxidation.

Effect of the silylacetal substituent on the C-H silylation reaction. The feasibility of the planned reaction sequence was assessed by conducting reactions with several catalysts and esters derived from 6-methylheptan-2-ol. A full list of results with esters is given in the supporting information. A selection of the results is shown in Scheme 1. The choice of ester was important for the success of the envisaged transformation; we investigated formate 2a, acetate 2b, pivalate 2c and trifluoroacetate 2d. The electronic and steric properties of these esters vary among this series. The hydrosilylation of all four esters with diethylsilane proceeded in the presence of catalytic (1,5cyclooctadiene)(methoxy)iridium(I) dimer ([Ir(cod)OMe]2 cod =1,5-cyclooctadiene) to complete conversion (by ¹H NMR spectroscopy) to form the intermediate silvlacetals 3a-3d. The crude silvlacetals were then used for the C-H bond activation step after evaporation of the solvent and other volatile materials.

The C–H silylation reactions of silylacetals **3a-3d** were conducted with Ir and Rh based catalysts. The reaction of trifluoromethylacetal **3d** in the presence of $[Ir(cod)OMe]_2$, 3,4,7,8-tetramethyl-1,10-phenanthroline (Me₄Phen) as catalyst and norbornene (nbe) as hydrogen acceptor provided 45% yield (by ¹H NMR spectroscopy) of the desired dioxasilinane **4d** over two steps. Side-products resulting from the disproportionation of silylacetals **3d** accounted for 12% of the total conversion. The same procedure with 6-methylheptan-2-yl formate **2c** formed the 6-membered silinane **4a** in 24% yield over the two steps. The analogous reactions with methylacetal **3b** and *tert*-butylacetal **3c** furnished a complex mixture without the formation of expected dioxasilinane **4b** and **4c**, respectively.

We tested rhodium catalysts that had been reported for the silylation of aromatic C–H bonds. The reactions of trifluoromethylacetal **3d** with chlorobis(cyclooctene)rhodium(I) dimer ([Rh(coe)₂Cl]₂, coe = cyclooctene) and 5,5'-Bis[di(3,5-di-*tert*butyl-4-methoxyphenyl)phosphino]-4,4'-bi-1,3-benzodioxole (DTBM-SEGPHOS) furnished the dioxasilinane **4d** in 28% yield, as determined by ¹H NMR spectroscopy. In the case of the rhodium-catalyzed reaction, a side product resulting from the hydrosilylation of norbornene with the trifluoromethylacetal **3d** formed in 17% yield.

Scheme 2. Initial evaluation of directing groups for β -C(sp³)–H silylation



^{*a*}[Ir(cod)OMe]₂ (1.0 mol%), Et₂SiH₂ (4.0 equiv), THF (0.5 M), rt or 50 °C, 24 h, N₂; ^{*b*}[Rh(coe)₂Cl]₂ (2.0 mol%), DTBM-SEGPHOS (6.0 mol%), nbe (1.5 equiv), THF (0.1 M), 100 °C, 16 h, N₂. ^{*c*}[Ir(cod)OMe]₂ (2.0 mol%), Me₄Phen (6.0 mol%), nbe (1.5 equiv), THF (0.1 M), 100 °C, 16 h, N₂.

The observed reactivity of the series of acetals **3a-3d** can be rationalized in terms of the stability and steric properties of the directing groups. The stability of silylacetal intermediates **3a-3d** is revealed by comparing the hydration constants of formaldehyde, acetaldehyde, pivalaldehyde, and trifluoroacetal-dehyde (see Scheme 2).¹³ Generally, the presence of electron-withdrawing groups adjacent to the carbonyl group of an aldehyde increase the stability of the hydrate relative to the free aldehyde. Consequently, formaldehyde and trifluoroacetal-hyde exist primarily as a geminal diol in aqueous solution, while pivalaldehyde exists primarily as a free aldehyde in aqueous solution. Consistent with this greater stability of these

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36 37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

acetals, trifluoroacetaldehyde silylacetal **3d** and formaldehyde silylacetal **3a** furnished the desired dioxasilinane **4**, but the substrates corresponding to less stable acetals did not. Although the silylacetals of formaldehyde and trifluoroacetaldehyde are stable (K_{hydration} > 10²), the trifluoroacetaldehyde silylacetal **3b** furnished the product of C–H silylation in higher yield than the formaldehyde silylacetal **3c** (45% vs 24%, see Scheme 2). This difference in reactivity parallels the greater steric hindrance of the trifluoroacetate than of the hydrogen (A-Value (–H) = 0 kcal/mol) of the formate (see Scheme 2).¹⁴ The more hindered trifluoromethyl group will impart a greater Thorpe-Ingold effect on the cyclization reaction.¹⁵

Because the reaction conducted with the combination of $[Ir(cod)OMe]_2$ and Me_4Phen furnished the desired dioxasilinane in higher yield than the reactions conducted with combination of $[Rh(coe)_2Cl]_2$ and DTBM-SEGPHOS, we tested a series of reaction conditions for the dehydrogenative silylation of trifluoroacetal **3d** catalyzed by $[Ir(cod)OMe]_2$. Evaluation of the ligand, hydrogen acceptor, solvent, concentration, and temperature, did not result in an increase in reaction yield (refer to the Supporting Information). Thus, we modified the substituent on the acetal to achieve high yields for the dehydrogenative silylation step.

Considering the initially observed effect of the stability and the steric hindrance of silylacetal on the silylation step, we evaluated a series of perfluorinated esters of 6-methylheptan-2-ol with varied steric properties (**2e-2h**, Table 1). The reduction of perfluoropropanoate **2e** with diethylsilane (4.0 equiv) in the presence of [Ir(cod)OMe]₂ (2.0 mol%) in heptane at 60 °C led to complete formation of the silylacetal (see entry 2, Table 1). Intramolecular C–H silylation of perfluoroethyl silylacetal **3e** in the presence of [Ir(cod)OMe]₂ (2.0 mol%), Me₄Phen (6.0 mol%) and nbe (1.5 equiv) in THF (0.1 M) at 100 °C formed the desired 6-membered dioxasilinane **4e** in 68% yield (by ¹H NMR spectroscopy) over two steps.

Table 1. Evaluation of fluorinated directing group

Me 2e , R _f 2f , R _f 2g , R _f 2h , R _f	$R_{f} = 1) [Ir(cod)C$ $H = C_{2}F_{5}$ $= C_{3}F_{7}$ $= CF_{2}Ph$ $= C_{7}F_{15}$	Me] ₂ , Et ₂ SiH ₂ ie, 60 °C ^a Me] ₂ , Me₄Phen IF, 100 °C ^b Me	R_{f} Me O SiEt ₂ 4e, $R_{f} = C_{2}F_{5}$ 4f, $R_{f} = C_{3}F_{7}$ 4g, $R_{f} = C_{7}F_{15}$
Entry	Ester	Conversion ^c	Yield ^d
1 ^{<i>c</i>}	2d , $R_f = CF_3$	100%	45%
2	$2e, R_f = C_2 F_5$	100%	68%
3	$2f, R_f = C_3 F_7$	96%	78%
4	$2g, R_f = CF_2Ph$	98%	75%
5	2h , $R_f = C_7 F_{15}$	95%	84%(75%) ^e

^{*a*}Conditions for hydrosilylation of ester: $[Ir(cod)OMe]_2$ (1.0 mol%), Et₂SiH₂ (4.0 equiv), heptane (0.5 M), 60 °C, 24-48 h, N₂; ^{*b*}Conditions for β -C(sp³)–H silylation: $[Ir(cod)OMe]_2$ (2.0 mol%), Me₄Phen (6.0 mol%), nbe (1.5 equiv), THF (0.1 M), 100 °C, 16 h, N₂; ^{*c*}Conversion for the hydrosilylation step determined by ¹H NMR spectroscopy; ^{*d*}Overall yield for the two step determined by ¹H NMR spectroscopy using CH₂Br₂ as internal standard; ^{*e*}Isolated yield.

This positive effect of the larger perfluoroalkyl group on the yield led us to test the perfluorobutanoate **2f** under the same reaction conditions (see entry 3, Table 1). The hydrosilylation of the ester occurred in 96% conversion, and the overall yield for the formation of dioxasilinane **4f** from the perfluorobutanoate **2f** was a 78%. Similar results were obtained with the 2,2-difluoro-2-phenylacetate **4g** (75% yield, entry 4, Table 1); only trace amounts of product resulting from C(sp²)–H bond silylation of the phenyl ring of the acetal was observed. Finally, subjecting the perfluorooctanoate ester **2h** to the sequence of hydrosilylation and β -C(sp³)–H silylation yielded the dioxasilinane **4h** in 84% yield over the two steps.

After identifying conditions for the hydrosilylation of the esters and dehydrogenative silylation of the $C(sp^3)$ –H bonds, we tested if dioxasilinane **4h** would undergo Tamao-Fleming oxidation to furnish 1,2-diols. After evaluating a series of conditions for the Tamao-Fleming oxidation we found that isolated silinane **4h** converts to the corresponding 1,2-diol in 72% yield (¹H NMR) in the presence of hydrogen peroxide, potassium bicarbonate and potassium fluoride in methanol at 60 °C (Scheme 3). Moreover, we found that these conditions for oxidation can be used without isolating the silinane **4h**. The crude silinane **4h** formed 6-methylheptane-1,2-diol **5a** in 59% isolated yield over the three steps, without isolation of any intermediates.

Scheme 3. Tamao-Fleming oxidation



^{*a*}Hydrosilylation of ester: $[Ir(cod)OMe]_2$ (1.0 mol%), Et₂SiH₂ (4.0 equiv), heptane (0.5 M), 60 °C, 48 h, N₂; ^{*b*}C–H silylation step: $[Ir(cod)OMe]_2$ (2.0 mol%), Me₄Phen (6.0 mol%), nbe (1.5 equiv), THF (0.1 M), 100 °C, 16 h, N₂; ^{*c*}Oxidation step: KHCO₃ (4.0 equiv), KF(4.0 equiv), 50% aqueous H₂O₂(10 equiv), MeOH, 60 °C, 2h; ^{*d*}Determined by ¹H NMR spectroscopy using CH₂Br₂ as internal standard; ^{*e*}Overall isolated yield for the steps 1-3.

Scope of the β -C–H silylation reaction. Studies to evaluate the selectivity and functional-group tolerance of the developed sequence comprising hydrosilylation and C–H silylation are summarized in Scheme 4. Substrates containing both primary and either secondary (4h), tertiary (4i, 4j) and benzylic (4q, 4r) β -C(sp³)–H bonds underwent silylation exclusively at the primary C–H bonds. Moreover, products resulting from potential intermolecular silylation of aromatic C–H bonds were not observed. Esters containing a TBDPS-protected alcohol (4n), an alkyl halide (4p) or a trisubstituted alkenes (4s) underwent the hydrosilylation, C–H bond silylation sequence without interference of the auxiliary functional groups. In addition, substrates bearing aryl chloride (4m), bromide (4r) and fluoride (4q) moieties underwent the same sequence to furnish the six-membered ring dioxasilinanes in 73-76% yield. Meth-

Page 4 of 7

ylenedioxy protected catechol (40) and methyl phenol ether (41) were also compatible with the reaction conditions. Esters, amides, ketones, and nitroarenes were not tolerated; these groups were reduced under the conditions for hydrosilylation. Substrates containing coordinating groups, such as nitriles and thiols, underwent the hydrosilylation step in low conversion.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

The hydrosilylation of perfluorooctanoates of tertiary alcohols under the developed conditions did not furnish the desired silylacetals, presumably due to the high steric hindrance. Thus, we examined a series of esters as directing groups for the sequence of hydrosilylation and C-H silylation of tertiary alcohols (refer to the Supporting Information for details). Among the investigated esters, only the hydrosilylation of trifluoroacetate and 2,2-difluoro-2-phenylacetate proceeded to high conversion. The silvlacetal derived from a tertiary 2,2difluoro-2-phenylacetate provided substantial amounts of products resulting from silvlation of the $C(sp^2)$ -H bond of the phenyl group in the difluorobenzyl substituent, leading to low yield of the C(sp³)–H silvlation product. However, the hydrosilylation of the trifluoroacetate of 2-methyl-4-phenylbutan-2ol 1t and subsequent dehydrogenative silvlation occurred to form the dioxasilinane 4t in 72% yield (by ¹H NMR spectroscopy) over the two steps (Scheme 3). Due to the instability of the dioxasilinane bearing a trifluoromethyl group, we isolated the 1,2-diol after the Tamao-Fleming oxidation in 52% yield over 3 steps. Thus, simple modification of the substituent on the directing group can match the steric and electronic requirements to achieve oxidation of the β C–H bond in high yield for both secondary and tertiary alcohols.

Scheme 4. Scope of the C-H silylation reaction



^{*a*}Hydrosilylation of ester: $[Ir(cod)OMe]_2$ (1.0 mol%), Et₂SiH₂ (4.0 equiv), heptane (0.5 M), 60 °C, 48 h, N₂; ^{*b*}C–H silylation step: $[Ir(cod)OMe]_2$ (2.0 mol%), Me₄Phen (6.0 mol %), nbe (1.5 equiv), THF (0.1 M), 100 °C, 16 h, N₂; ^{*c*}Determined by ¹H NMR spectroscopy using CH₂Br₂ as internal standard; ^{*d*}Isolated yield for the 1,2 diols over the threes step, oxidation step: KHCO₃ (4.0

equiv), KF (4.0 equiv), 50% aqueous $\rm H_2O_2$ (10 equiv), MeOH, 60 °C, 2h.

Hydroxylation of natural products derivatives. The developed sequence of hydrosilylation, C-H silylation, and oxidation was applicable to the selective hydroxylation of a series of natural products derivatives containing tertiary and secondary alcohols (Scheme 5). The perfluorooctanoate ($R_f = {}^{n}C_7F_{15}$) substituent was a suitable traceless directing group for reactions of natural products containing secondary 2-methyl alcohols. The developed sequence led to the introduction of a second hydroxyl group in the β position of the alcohols derived from geranylacetone (5aa), dihydro- β -ionone (5ab), dihydro- α -ionone (5ac) and the 27-nor-25-ketocholesterol (5ad). The combination of hydrosilylation and C-H silvlation occurred in 69-76% yield (determined by ¹H NMR spectroscopy) over the two steps. The resulting dioxasilinane was subjected to Tamao-Fleming oxidation to furnish the final 1,2diols in 41-54% isolated yield over the three-step sequence.





^{*a*}Hydrosilylation of ester: $[Ir(cod)OMe]_2$ (1.0 mol%), Et_2SiH_2 (4.0 equiv), heptane (0.5 M), 60 °C, 48 h, N₂; ^{*b*}C-H silylation step: $[Ir(cod)OMe]_2$ (2.0 mol%), Me₄Phen (6.0 mol %), nbe (1.5

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

equiv), THF (0.1 M), 100 °C, 16 h, N₂; ^cOxidation step: KHCO₃ (4.0 equiv), KF(4.0 equiv), 50% aqueous $H_2O_2(10 \text{ equiv})$, MeOH, 60 °C, 2h; ^dIsolated yield for the 1,2 diols over the three steps ^eYield for the oxasilolane determined by ¹H NMR spectroscopy using CH₂Br₂ as internal standard; ^fThe starting dihydro- α -ionone was purchased as the mixture of cyclohex-2-ene and cyclohex-3-ene derivative in 4:1 ratio.

Natural products containing tertiary alcohols underwent the sequence with a trifluoromethyl substituent containing removable directing group. The developed strategy led to hydroxylation of the primary β C–H bonds of (+)-cedrol (**5ab**), ledol (**5ag**), (-)- α -bisabolol (**5af**) and α -terpinol (**5ae**) to form the coresponding 1,2-diols. The diols were isolated in 21-36% yield over the three-step sequence. Although occurring in modest yield this sequence leads to the formation of readily isolable amounts of the modified products. The initial introduction of the directing group (TFAA, pyridine, DCM) did not require isolation by column chromatography.

The complementarity of our developed method to functionalize alcohols is well illustrated in the case of (+)-cedrol (5ah), a sesquiterpenoid used in the perfume industry extracted from Texas cedarwood oil. Direct chemical or microbial¹⁶ oxidation of cedrol has been studied extensively. Generally, the oxidation reactions of cedrol with *m*-chlorobenzoic acid and ruthenium or iron catalysts lead to the introduction of a hydroxyl group at the tertiary C(2) carbon (5ak).¹⁷ The oxidation of cedrol with the "Gif IV system" resulted in the introduction of ketone functionality at the C(10) position (**5ai**).¹⁸ Finally, methods based on the generation of oxygen-centered radicals, followed by 1,5-hydrogen atom transfer, lead to etherification of the distal methyl group (C(14) position, 5aj).¹⁹ None of the developed methods have led to the selective oxidation of primary C–H bonds located β to the alcohol. Thus, the method we report offers new opportunities for selective alcohol oxidation.

Conclusion.

We have developed an Ir-catalyzed functionalization of C– H bonds located β to hydroxyl functionality by incorporating traceless perfluorinated ester directing groups. Diethyl silane plays a dual role in the overall process, acting as the reductant for the hydrosilylation of the ester and as the reagent for silylation of the C–H bond, in this case to obtain six-membered dioxasilinanes. The silicon-containing intermediates undergo oxidation to 1,2-diols, leading to a net alcohol directed β -C(sp³)–H hydroxylation. The iridium catalysts react preferentially with primary β -C(sp³)–H bonds over secondary, benzylic or tertiary C–H bonds, leading to a selectivity that complements the selectivity of previously developed oxidation methods. Thus, this functionalization of an alkyl C–H bond enriches the currently-available toolbox of C–H bond functionalization reaction suitable for diversification of complex structures.

ASSOCIATED CONTENT

Supporting Information

Experimental details, synthetic procedures and spectral data.

AUTHOR INFORMATION

Corresponding Author

*jhartwig@berkeley.edu

ORCID

John F. Hartwig: 0000-0002-4157-468X.

Ala Bunescu: 0000-0002-3880-8182 Trevor W. Butcher: 0000-0001-5006-692X

Funding Sources

This work was supported by the NIH (GM 115812) and The Dow Chemical Company.

ACKNOWLEDGMENT

We thank the NIH (GM 115812) and The Dow Chemical Company for support of this work. AB thanks the Swiss National Science Foundation for a postdoctoral fellowship. TWB gratefully acknowledges the National Science Foundation for a predoctoral fellowship.

REFERENCES

1. (a) Jazzar, R.; Hitce, J.; Renaudat, A.; Sofack Kreutzer, J.; Baudoin, O., *Chemistry - A European Journal* **2010**, *16*, 2654-2672; (b) Gutekunst, W. R.; Baran, P. S., *Chem. Soc. Rev.* **2011**, *40*, 1976-1991; (c) McMurray, L.; O'Hara, F.; Gaunt, M. J., *Chem. Soc. Rev.* **2011**, *40*, 1885-1898; (d) Brückl, T.; Baxter, R. D.; Ishihara, Y.; Baran, P. S., *Acc. Chem. Res.* **2012**, *45*, 826-839; (e) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K., *Angew. Chem. Int. Ed.* **2012**, *51*, 8960-9009; (f) He, J.; Wasa, M.; Chan, K. S. L.; Shao, Q.; Yu, J.-Q., *Chem. Rev.* **2017**, *117*, 8754-8786.

2. (a) Hartwig, J. F., *Chem. Soc. Rev.* **2011**, *40*, 1992-2002; (b) Mkhalid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F., *Chem. Rev.* **2010**, *110*, 890-931.

3. (a) Ghavtadze, N.; Melkonyan, F. S.; Gulevich, A. V.; Huang, C.; Gevorgyan, V., *Nat. Chem.* **2014**, *6*, 122; (b) Cheng, C.; Hartwig, J. F., *Chem. Rev.* **2015**, *115*, 8946-8975; (c) Kon, K.; Suzuki, H.; Takada, K.; Kohari, Y.; Namikoshi, T.; Watanabe, S.; Murata, M., *ChemCatChem* **2016**, *8*, 2202-2205; (d) Fukumoto, Y.; Hirano, M.; Chatani, N., *ACS Catal.* **2017**, *7*, 3152-3156.

4. Henkel, T.; Brunne, R. M.; Müller, H.; Reichel, F., Angew. Chem. Int. Ed. 1999, 38, 643-647.

5. (a) Zhang, S.-Y.; Zhang, F.-M.; Tu, Y.-Q., *Chem. Soc. Rev.* **2011**, *40*, 1937-1949; (b) Ketcham, J. M.; Shin, I.; Montgomery, T. P.; Krische, M. J., *Angew. Chem. Int. Ed.* **2014**, *53*, 9142-9150.

6. (a) Barton, D. H. R.; Beaton, J. M.; Geller, L. E.; Pechet, M. M., *J. Am. Chem. Soc.* **1960**, *82*, 2640-2641; (b) Hartung, J.; Gottwald, T.; Špehar, K., *Synthesis* **2002**, *2002*, 1469-1498.

7. (a) Doyle, M. P.; Kalinin, A. V.; Ene, D. G., *J. Am. Chem. Soc.* **1996**, *118*, 8837-8846; (b) Espino, C. G.; Wehn, P. M.; Chow, J.; Du Bois, J., *J. Am. Chem. Soc.* **2001**, *123*, 6935-6936; (c) Chen, K.; Richter, J. M.; Baran, P. S., *J. Am. Chem. Soc.* **2008**, *130*, 7247-7249.

8. (a) Simmons, E. M.; Hartwig, J. F., J. Am. Chem. Soc. 2010, 132, 17092-17095; (b) Simmons, E. M.; Hartwig, J. F., Nature 2012, 483, 70-73; (c) Li, B.; Driess, M.; Hartwig, J. F., J. Am. Chem. Soc. 2014, 136, 6586-6589; (d) Lee, T.; Wilson, T. W.; Berg, R.; Ryberg, P.; Hartwig, J. F., J. Am. Chem. Soc. 2015, 137, 6742-6745; (e) Lee, T.; Hartwig, J. F., Angew. Chem. Int. Ed. 2016, 55, 8723-8727; (f) Su, B.; Zhou, T. G.; Li, X. W.; Shao, X. R.; Xu, P. L.; Wu, W. L.; Hartwig, J. F.; Shi, Z. J., Angew. Chem. Int. Ed. 2017, 56, 1092-1096.

9. Ren, Z.; Mo, F.; Dong, G., J. Am. Chem. Soc. 2012, 134, 16991-16994.

10. Xu, Y.; Yan, G.; Ren, Z.; Dong, G., Nat Chem 2015, 7, 829-834.

11. Hua, Y.; Asgari, P.; Avullala, T.; Jeon, J., J. Am. Chem. Soc. 2016, 138, 7982-7991.

12. (a) Saget, T.; Cramer, N., Angew. Chem. Int. Ed. 2012, 51, 12842-12845; (b) Piou, T.; Bunescu, A.; Wang, Q.; Neuville, L.; Zhu, J., Angew. Chem. Int. Ed. 2013, 52, 12385-12389.

13. Guthrie, J. P., Can. J. Chem. 1975, 53, 898-906.

14. Hirsch, J. A., Table of Conformational Energies—1967. In *Top. Stereochem.*, John Wiley & Sons, Inc.: 2007; pp 199-222.

15. Jung, M. E.; Piizzi, G., Chem. Rev. 2005, 105, 1735-1766.

16. Fraga, B. M.; Guillermo, R.; Hanson, J. R.; Truneh, A., *Phytochemistry* **1996**, *42*, 1583-1586.

17. (a) Tenaglia, A.; Terranova, E.; Waegell, B., J. Org. Chem. **1992**, *57*, 5523-5528; (b) Gómez, L.; Canta, M.; Font, D.; Prat, I.; Ribas, X.; Costas, M., J. Org. Chem. **2013**, *78*, 1421-1433; (c) Jana, S.; Ghosh, M.; Ambule, M.; Sen Gupta, S., Org. Lett. **2017**, *19*, 746-749; (d) Mack, J. B.; Gipson, J. D.; Du Bois, J.; Sigman, M. S., J. Am. Chem. Soc. **2017**, *139*, 9503-9506. 18. Barton, D. H. R.; Beloeil, J. C.; Billion, A.; Boivin, J.; Lallemand, J. Y.; Lelandais, P.; Mergui, S., *Helv. Chim. Acta* , *70*, 2187-2200.

19. (a) Brun, P.; Waegell, B., *Tetrahedron* **1976**, *32*, 1137-1145; (b) Dorta, R. L.; Francisco, C. G.; Freire, R.; Suárez, E., *Tetrahedron Lett.* **1988**, *29*, 5429-5432; (c) Hung, K.; Condakes, M. L.; Morikawa, T.; Maimone, T. J., *J. Am. Chem. Soc.* **2016**, *138*, 16616-16619.

