


Modeling the Claus Reaction: Preparation of *trans*-Pt(PPh₃)₂(phthalimido)S(O)₂SR

Alan Shaver,* Hélène Boily, and Anne-Marie Lebuis


Department of Chemistry, McGill University, 801 Sherbrooke Street West, Montreal, Quebec, Canada H3A 2K6

Received May 24, 1996

The removal of sulfur from crude oil is accomplished by hydrodesulfurization (HDS)¹ followed by the Claus process,² which converts the H₂S produced by HDS to sulfur and water (eq 1). Natural gas contaminated by H₂S is purified by direct




application of the Claus process. The Claus reaction is conducted at 300 °C using alumina as a catalyst although other materials also catalyze the reaction. The mechanism is unknown, but the reaction must involve the formation of sulfur–sulfur bonds and oxygen transfer, a suite of reactions with few precedents in homogeneous catalysis. Three simple conceptual models of the Claus reaction can be described. Model A depicts



attack by H₂S on adsorbed SO₂, model B represents attack by SO₂ on adsorbed H₂S,³ and in C both the H₂S and the SO₂ are adsorbed before reaction.⁴ Studies of the sequential adsorption and reactions of SO₂ and H₂S on alumina⁵ are for the most part inconclusive; however, preadsorbed SO₂ is reactive toward H₂S.^{5d} Therefore, the complex Pt(PPh₃)₃SO₂,⁶ chosen as an example of preadsorbed SO₂, was treated with a source of RS⁺ as a simulation of model A.

Treatment of Pt(PPh₃)₃SO₂ with RSphth⁷ gave pale yellow complexes of the type *trans*-Pt(PPh₃)₂(phth)S(O)₂SR, **1a–f**, where phth = phthalimido (eq 2). The IR and NMR (¹H and



1a, R = CH₃
b, R = CH₂CH₃
c, R = n-C₃H₇
d, R = CHMe₂
e, R = 4-C₆H₄Me
f, R = CH₂Ph

³¹P) spectra of **1a–f** are consistent with their formulation,⁸ and the structure of **1f**, shown in Figure 1, confirms the presence of the PtS(O)₂SCH₂Ph moiety.⁹ Interestingly, while organic thio-sulfonates (RS(O)₂SR') are known,¹⁰ few structures have been reported;¹¹ complexes **1a–f** are the first to contain an RSS(O)₂[−] ligand.¹² Sulfito groups are the parent ligands of this new class.^{3k,13}

Attachment of the RS⁺ residue to the SO₂ ligand of Pt(PPh₃)₃SO₂ and oxidation of the metal center result in a much shorter Pt–S distance (2.261(3) Å) in **1f** than in the precursor (2.368(3) Å).¹⁴ The Pt–S distance in **1f** is also shorter than those in *cis*-Pt(PPh₃)₂(phth)SSCHMe₂ (2.353(3) Å),¹⁵ [Pt(PPh₃)₂(S(O)–

* Author to whom correspondence should be addressed.

- (1) (a) Satterfield, C. N. *Heterogeneous Catalysis in Industrial Practice*, 2nd ed.; McGraw-Hill: New York, 1991; pp 378–383. (b) Gates, B. C.; Katzer, J. R.; Schuit, G. C. A. *Chemistry of Catalytic Processes*; McGraw Hill: New York, 1979; pp 390–445.
- (2) (a) Grancher, P. *Hydrocarbon Process.* **1978**, 57, 155. (b) Grancher, P. *Hydrocarbon Process.* **1978**, 57, 257. (c) George, Z. M.; Tower, R. W. *Can. J. Chem. Eng.* **1985**, 63, 618 and references therein.
- (3) For examples of model B involving soluble metal thiolates and sulfides see: (a) Eller, P. G.; Kubas, G. J. *J. Am. Chem. Soc.* **1977**, 99, 4346. (b) Shaver, A.; Plouffe, P.-Y. *Inorg. Chem.* **1992**, 31, 1823. (c) Schenk, W. A.; Dombrowski, E.; Reuther, I.; Stur, T. Z. *Naturforsch., B: Chem. Sci.* **1992**, 47B, 732. (d) Darendsbourg, M. Y.; Tuntulani, T.; Reibenspies, J. H. *Inorg. Chem.* **1994**, 33, 611. (e) Kubas, G. J.; Ryan, R. R. *J. Am. Chem. Soc.* **1985**, 107, 6138. (f) Kubas, G. J.; Wasserman, H. J.; Ryan, R. R. *Organometallics* **1985**, 4, 419. (g) Brunner, H.; Klement, U.; Pfauntsch, J.; Wachter, J. *Angew. Chem., Int. Ed. Engl.* **1987**, 26, 230. (h) Kubas, G. J.; Ryan, R. R. *Inorg. Chem.* **1984**, 23, 3181. (i) Toupadakis, A.; Kubas, G. J.; Burns, C. J. *Inorg. Chem.* **1992**, 31, 3810. (j) Kubas, G. J.; Ryan, R. R.; Kubat-Martin, K. A. *J. Am. Chem. Soc.* **1989**, 111, 7823. (k) Kubat-Martin, K. A.; Kubas, G. J.; Ryan, R. R. *Organometallics* **1989**, 8, 1910. (l) Yamanari, K.; Mori, M.; Dogi, S.; Fuyuhiko, A. *Inorg. Chem.* **1994**, 33, 4807. (m) Heyke, O.; Neher, A.; Lorenz, I.-P. *Z. Anorg. Allg. Chem.* **1992**, 608, 23.
- (4) For examples of model C involving soluble complexes containing both thiolato and SO₂ ligands see refs 3b and 3c and: Kubas, G. J. *Inorg. Chem.* **1979**, 18, 182.
- (5) (a) Datta, A.; Cavell, R. G.; Tower, T. W.; George, Z. M. *J. Phys. Chem.* **1985**, 89, 443. (b) Datta, A.; Cavell, R. G. *J. Phys. Chem.* **1985**, 89, 450. (c) Datta, A.; Cavell, R. G. *J. Phys. Chem.* **1985**, 89, 454. (d) Karge, H. G.; Dalla Lana, I. G. *J. Phys. Chem.* **1984**, 88, 1958.
- (6) Levison, J. J.; Robinson, S. D. *J. Chem. Soc., Dalton Trans.* **1972**, 2013.
- (7) (a) Behforouz, M.; Kerwood, J. E. *J. Org. Chem.* **1969**, 34, 51. (b) Buchel, K. H.; Conte, A. *Chem. Ber.* **1967**, 100, 1248. (c) Harpp, D. N.; Ash, D. K.; Back, T. G.; Gleason, J. G.; Orwig, B. A.; van Horn, W. F.; Snyder, J. P. *Tetrahedron Lett.* **1970**, 3551.
- (8) Preparative procedures and characterization data are given in the Supporting Information.
- (9) Crystal data for *trans*-Pt(PPh₃)₂(phth)S(O)₂SCH₂Ph·0.75CH₂Cl₂ **1f**: monoclinic *P2₁/c* (No. 14), *a* = 21.212(4) Å, *b* = 13.996(2) Å, *c* = 17.789(3) Å, *V* = 4868.9(14) Å³, β = 112.79(1)°, *Z* = 4, *D_c* = 1.524 g/cm³, μ(Mo Kα) = 3.190 mm^{−1}, λ(Mo Kα) = 0.709 30 Å (graphite monochromated); 2θ_{max} = 50°, 32 688 measured reflections (8585 unique, *R*(int) = 0.133); *R* = 0.063, *R_w* = 0.108, GOF = 0.887 (4409 reflections with *I* > 2.00σ(*I*)). A complete structural report is included in the Supporting Information.
- (10) (a) Block, E. *Angew. Chem., Int. Ed. Engl.* **1992**, 31, 1135. (b) Freeman, F. *Chem. Rev.* **1984**, 84, 117. (c) Oae, S. *Organic Chemistry of Sulfur*; Plenum Press: New York, 1977. (d) Reid, E. E. *Organic Chemistry of Bivalent Chemistry*; Chemical Publishing Co.: New York, 1958; Vol. 1.
- (11) (a) Wahl, G. H.; Bordner, J.; Harpp, D. N.; Gleason, J. G. *Acta Crystallogr., Sect. B* **1973**, B29, 2272. (b) Noordik, J. H.; Vos, A. *Recl. Trav. Chim. Pays-Bas* **1967**, 86, 156. (c) Block, E.; O'Connor, J. J. *Am. Chem. Soc.* **1974**, 96, 3921.
- (12) The RSS(O)₂[−] ligands in **1a–f** are isomeric with those in (a) CpRu(PPh₃)(CO)SS(O)₂-4-C₆H₄Me (Shaver, A.; Plouffe, P. Y. *J. Am. Chem. Soc.* **1991**, 113, 7780) and (b) CpRu(CO)₂SS(O)₂Ph (Trojansek, D. Ph.D. Thesis, McGill University, 1995).
- (13) (a) Michelin, R. A.; Napoli, M.; Ros, R. *J. Organomet. Chem.* **1979**, 175, 239. (b) Barlex, D. M.; Kemmitt, R. D. W. *J. Chem. Soc., Dalton Trans.* **1972**, 1437. (c) Graziani, M.; Ros, R.; Carturna, G. *J. Organomet. Chem.* **1971**, 27, C19. (d) Schenk, W. A.; Karl, U. Z. *Naturforsch., B: Chem. Sci.* **1989**, 44B, 993. (e) Schenk, W. A.; Urban, P.; Stahrfeldt, T.; Dombrowski, E. A. *Z. Naturforsch., B: Chem. Sci.* **1992**, 47B, 1493.
- (14) Ellen, P. G.; Ryan, R. R.; Moody, D. C. *Inorg. Chem.* **1976**, 15, 2442.
- (15) Shaver, A.; Hartgerink, J.; Lai, R. D.; Bird, P. *Organometallics* **1983**, 2, 938.

