

Communication

Redox Non-Innocent Behavior of Tris(2-PyridyImethyI)amine Bound to a Lewis Acidic Rh(III) Ion Induced by C-H Deprotonation

Hiroaki Kotani, Takumi Sugiyama, Tomoya Ishizuka, Yoshihito Shiota, Kazunari Yoshizawa, and Takahiko Kojima

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.5b06237 • Publication Date (Web): 24 Aug 2015 Downloaded from http://pubs.acs.org on August 25, 2015

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Redox Non-Innocent Behavior of Tris(2-Pyridylmethyl)amine Bound to a Lewis Acidic Rh(III) Ion Induced by C-H Deprotonation

Hiroaki Kotani,⁺ Takumi Sugiyama,⁺ Tomoya Ishizuka,⁺ Yoshihito Shiota,[‡] Kazunari Yoshizawa,^{‡,§} and Takahiko Kojima^{*,†}

[†]Department of Chemistry, Graduate School of Pure and Applied Sciences, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8571, Japan [‡]Institute for Materials Chemistry and Engineering, Kyushu University, Motooka, Nishi-Ku, Fukuoka 819-0395, Japan

[§]Elements Strategy Initiative for Catalysts & Batteries, Kyoto University, Nishikyo-ku, Kyoto 615-8520, Japan

Supporting Information Placeholder

ABSTRACT: Rh(III) complexes having tris(2-pyridylmethyl)amine (TPA) and its derivative as tetradentate ligands showed reversible deprotonation at a methylene moiety of the TPA ligands upon addition of a strong base as confirmed by spectroscopic measurements and X-ray crystallography. The deprotonation selectively occurred at the axial methylene moiety rather than equatorial counterparts because of the thermodynamic stability of corresponding deprotonated complexes. One-electron oxidation of the deprotonated Rh(III)-TPA complex afforded a unique TPA radical bound to the Rh(III) center by a ligand-centered oxidation. This is the first example to demonstrate emergence of the redox-noninnocent character of the TPA ligand.

Metal-ligand cooperation has been recognized as one of new methods to develop novel and striking functionality of metal complexes, which often derives from pyridine-based ligands.¹ Milstein and co-workers have reported that a novel Ru catalyst having a pyridine-based PNN pincer ligand is capable of proceeding the C– H bond activation of alcohols and amines, triggered by deprotonation and synergetic dearomatization processes of the ligand by the metal-ligand cooperation.² The deprotonated complex is produced as a key intermediate and stabilized by the contribution of a dearomatized resonance structure among plausible resonance structures as shown in Scheme 1,³ although the free pyridylmethyl group is not acidic without metal coordination. In fact, the contribution of dearomatized resonance structures was suggested by crystal structures of deprotonated complexes in the light of bond



lengths in the ligand.³

On the other hand, the ligand deprotonation has been known to induce redox-non-innocent character in an originally redoxinnocent ligand.⁴ For instance, Grützmacher and co-workers have reported successful isolation of a Rh(I) complex with an aminyl radical by one-electron oxidation of a deprotonated Rh(I)-amido complex.⁵ Also, de Bruin and co-workers have reported ligand oxidation of a deprotonated Ir(I)-BPA complex (BPA = bis(2picolyl)amine) to form a radical ligand.⁶ However, these sequential deprotonation and oxidation processes of deprotonated ligands have been limited to the case of metal complexes having amino (NH) groups in the ligand as deprotonation sites. In order to expand the scope of applicable ligands without NH groups as the deprotonation site, a combination of metal-ligand cooperation and subsequent ligand oxidation is required because a variety of radical intermediate-bound metal complexes $([M^{n+}(L_{\bullet})]^{n+})$ have been investigated as, for example, models of galactose oxidase to realize the importance of the radical character of the ligand.⁷

The concept of metal-ligand cooperation is expected to be expanded to metal complexes having versatile and well-known chelating ligands without NH groups. In contrast to the aforementioned BPA ligand, tripodal pyridylamine ligands such as TPA (= tris(2pyridylmethyl)amine) have no easily removable protons and only methylene protons can be deprotonated.^{8,9} So far, a large number of metal-TPA complexes have been reported to investigate their reactivity in oxidation reactions and in O₂ activation chemistry.¹⁰ However, there is no report on the observation of a metal complex having deprotonated TPA as a ligand and redox non-innocent behavior



Figure 1. (a, b) Chemical structures of Rh(III) complexes used in this study. (c) An ORTEP drawing of $[Rh^{III}(Cl)_2(L_2-H6)]^+$.



Figure 2. ¹H NMR spectra of (a) $[Rh^{III}(Cl)_2(L_1-H6)]^+$ (2.1 mM) and (b) $[Rh^{III}(L_1-H5)(Cl)_2]$ obtained upon addition of KOH (5.0 mM) in DMSO-*d*₆ at 298 K under Ar. Dotted lines denote the chemical shift of the axial pyridine moiety before and after deprotonation process.

of any metal-bound TPA derivatives.

We report herein unique deprotonation from the TPA ligand bound to a Lewis-acidic Rh(III) center via metal-ligand cooperation as shown in Scheme 1 and subsequent one-electron oxidation of a deprotonated complex allows us to investigate redox-noninnocent behavior of the TPA ligand. The Rh(III) ion was selected to stabilize the deprotonated species¹¹ toward further chemical manipulation based on the Lewis acidity and to perform oxidation of the deprotonated ligand based on the redox-inactive character, which fixes the oxidation state to be the Rh(III) state due to the instability of the Rh(II) and Rh(IV) states in the TPA environment.^{10c}

Synthesis of $[Rh^{III}(Cl)_2(L_1-H6)](Cl)^{12a}$ and $[Rh^{III}(Cl)_2(L_2-H6)](Cl)^{12a}$ H6](PF₆) (L₂-H6: TPACOOMe = tris(5-methoxycarbonyl-2pyridylmethyl)amine^{12b}) was accomplished by synthetic routes described in Supporting Information (SI). The methoxycarbonyl (COOMe) groups were introduced to the 5-positions of all pyridine rings to stabilize the deprotonated complex (vide infra). The crystal structure of [Rh^{III}(Cl)₂(L₂-H6)](PF₆) was determined by Xray crystallography as shown in Figure 1c. It should be noted that the impact of introduction of the COOMe groups on the structural parameters of $[Rh^{III}(Cl)_2(L_2-H6)]^+$ is negligible; there is no difference (< 0.01 Å) of bond lengths of Rh–Nx (x = 1-4) between $[Rh^{III}(Cl)_2(L_1-H6)]^+$ and $[Rh^{III}(Cl)_2(L_2-H6)]^+$ (see Table S2 in SI). Upon addition of a strong base such as potassium tert-butoxide (KO'Bu) or KOH into a DMSO-d₆ solution containing $[Rh^{III}(Cl)_2(L_1-H6)]^+$, drastic change of the ¹H NMR spectrum was observed as shown in Figure 2.13 The 1H NMR signals of deproto-



Figure 3. A schematic description of $[Rh^{\rm III}(L_1\text{-}H5)(Cl)_2]$ and an ORTEP drawing of the crystal structure.



Figure 4. UV-vis spectra observed in titration of $[Rh^{III}(Cl)_2(L_1-H6)]^+$ (0.10 mM) with (a) KO'Bu or (b) DBU in CH₃CN at 294 K under Ar. Inset: Plots of Abs at 293 nm vs initial concentration of bases. A dotted line denotes UV-vis spectrum of DBU (15 mM) in CH₃CN.

nated complexes were assigned by ¹H-¹H COSY spectroscopy (Figure S1 in SI). The ¹H NMR signals due to the axial pyridine moiety of $[Rh^{III}(L_1-H5)(Cl)_2]$ (L_1 -H5: deprotonated TPA) were revealed to show larger change of chemical shifts in comparison with those assigned to the equatorial pyridine moieties by the deprotonation. In addition, peak integration of the singlet signal at 4.15 ppm due to the methylene moiety of the axial pyridylmethyl arm in $[Rh^{III}(L_1-H5)(Cl)_2]$ decreased to about 50% of the original value for the singlet peak at 5.33 ppm observed for $[Rh(Cl)_2(L_1-H6)]^+$ (Figure 2). These results indicate that the deprotonation of $[Rh^{III}(Cl)_2(L_1-H6)]^+$ occurs at the axial pyridylmethyl group rather than at the equatorial pyridylmethyl groups.

In order to clarify the reason for the selectivity of the deprotonation at the axial methylene moiety, we evaluated the thermodynamic stability of $[Rh^{III}(L_1-H5)(Cl)_2]$ (deprotonation from the methylene moiety of the axial pyridylmethyl arm) and $[Rh^{III}(L_1-H5eq)(Cl)_2]$ (deprotonation from the methylene moiety of one of the equatorial pyridylmethyl arms), using density functional theory (DFT) calculations. As a result, $[Rh^{III}(L_1-H5eq)(Cl)_2]$ was indicated to be more stable by 9.8 kcal/mol than $[Rh^{III}(L_1-H5eq)(Cl)_2]$. This difference should be derived from the conformational distortion of $[Rh^{III}(L_1-H5eq)(Cl)_2]$ as depicted in Figure S3 in SI: The distortion of $[Rh^{III}(L_1-H5eq)(Cl)_2]$ would be derived from the rotation around an N-Rh axis due to the sp² character of the deprotonated equatorial methylene carbon.

The structure of $[Rh^{III}(L_1-H5)(Cl)_2]$ was further confirmed by X-ray crystallography as shown in Figure 3.¹⁴ The bond length of C1–C2 in $[Rh^{III}(L_1-H5)(Cl)_2]$ (1.364(8) Å) is much shorter than that of $[Rh^{III}(Cl)_2(L_1-H6)]^+$ (1.480(9) Å^{12a}), as described in Table S3 and Figure S4 in the SI. Judging from the structural change, difference of bond lengths was mainly observed on the axial pyridine moiety in comparison with those of equatorial pyridine moieties even in the crystal. This result is consistent with those obtained from the spectroscopic measurements. The structure of $[Rh^{III}(L_1-H5)(Cl)_2]$ also indicate that the dearomatized resonance structures of **B**, **C**, and **D** (Scheme 1) of the axial pyridylmethyl moiety 1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 60 should be dominant in the actual structure of $[Rh^{III}(L_1-H5)(Cl)_2]$ compared to the aromatic structure of **A** as shown in Scheme 1. In addition, the coordination of the pyridine N atom to a highly Lewis-acidic Rh(III) center should stabilize the dearomatized resonance structure **D** (Scheme 1) of $[Rh^{III}(L_1-H5)(Cl)_2]$ by trapping the negative charge on the N atom.

When we titrated $[Rh^{III}(Cl)_2(L_1-H6)]^+$ with KO'Bu by using UV-vis spectroscopy in CH₃CN at 294 K, one equivalent of KO'Bu was consumed to complete the spectral change, which should be derived from the deprotonation of $[Rh^{III}(Cl)_2(L_1-H6)]^+$, as shown in Figure 4a. This result indicates that the further deprotonation of $[Rh^{III}(L_1-H5)(Cl)_2]$ is negligible. Upon addition of 1,8diazabicyclo [5.4.0] undec-7-ene (DBU) as a weaker base (pK_a = 24.3 in CH₃CN)¹⁵ than KO'Bu, saturation behavior was observed in a titration plot as shown in Figure 4b. The pK_a value of the axial methylene proton of [Rh^{III}(Cl)₂(L₁-H6)]⁺ was thus determined to be 27.3 in CH₃CN at 294 K, according to the equation of the acidbase equilibrium (Experimental Section in SI). Similarly, the pK_{a} value of $[Rh^{III}(Cl)_2(L_1-H6)]^+$ in DMSO was also determined to be 16.1 by ¹H NMR titration with potassium acetylacetonate ($pK_a =$ 15.1 in DMSO)¹⁶ (Figure S5 in SI). This pK_a value in DMSO was significantly smaller than that of 2-benzylpyridine as a reference compound without a Rh(III) ion ($pK_a = 28.8$ in DMSO).¹⁶ The large decrease of the pK_a value is derived from the stabilization of $[Rh^{III}(L_1-H5)(Cl)_2]$ by coordination of deprotonated axial pyridine to a highly Lewis-acidic Rh(III) center as evidenced by the analysis on the crystal structure shown in Figure 3.

If the localized negative charge of the dearomatized resonance structure **B** or **C** depicted in Scheme 1 would be further stabilized by electron-withdrawing substituents introduced to the corresponding positions, the deprotonated complex should be more stabilized. According to this hypothesis, COOMe groups as electron-withdrawing substituents were introduced to the 5-positions of all pyridyl groups of TPA as described in Figure 1b. Similar spectral change was detected for the ¹H NMR spectrum of $[Rh(Cl)_2(L_2-$ H6)]⁺ as observed for $[Rh(Cl)_2(L_1-H6)]^+$, as depicted in Figure S6 (SI). This result indicates that one methylene proton of the axial pyridylmethyl arm in $[Rh(Cl)_2(L_2-H6)]^+$ is removed to afford $[Rh(Cl)_2(L_2-H5)]^+$. The pK_a value of $[Rh^{III}(Cl)_2(L_2-H6)]^+$ was determined to be 20.9 in CH₃CN at 296 K by UV-vis titration with triethyl amine (p K_a = 18.8 in CH₃CN),¹⁵ as shown in Figure S7 in SI. The lower pK_a value indicates that stabilization of the deprotonated complex is made through delocalization of the negative charge at the 5-position (dearomatized structure B in Scheme 1) into the methoxycarbonyl group. On the basis of the pK_a values of $[Rh^{\rm III}(Cl)_2(L_1\text{-}H6)]^{\scriptscriptstyle +}$ (27.3) and $[Rh^{\rm III}(Cl)_2(L_2\text{-}H6)]^{\scriptscriptstyle +}$ (20.9) in CH₃CN, the deprotonated species derived from the latter is more stable by 8.7 kcal mol $^{-1}(\Delta\Delta G^{\circ})$ than that derived from the former.

In order to develop reactivity of the Rh(III) complex having the deprotonated TPA ligand, we examined a reaction of the deprotonated Rh(III)-TPA complex with methyl iodide (MeI). Upon addition of MeI into the DMF solution of $[Rh^{III}(L_1-HS)(Cl)_2]$, methylation of $[Rh^{III}(L_1-HS)(Cl)_2]$ occurred at 298 K. The structure of the isolated methylated product $([Rh^{III}(Cl)(L_1-HSMe)]^+$ (L_1 -HSMe = methylated TPA) was determined by X-ray crystallography to reveal that the methylation occurred at the deprotonated methylene moiety as shown in Figure 5. Since the methylation induces the chirality at the methyne carbon, in the monoclinic crystal, both enantiomers (the *R*- and *S*-form) existed as a racemic mixture. The ¹H NMR spectrum of $[Rh^{III}(Cl)_2(L_1-HSMe)]^+$ showed



Figure 5. Methylation of $[Rh^{III}(L_1-H5)(Cl)_2]$ and an ORTEP drawing of $[Rh^{III}(Cl)_2(L_1-H5Me)]^+$.

doublets due to the methyl group and the methyne proton of the reacted axial pyridylmethyl arm with the same coupling constant (6.8 Hz), together with two AX doublets (15.3 and 16.5 Hz) assigned to the equatorial methylene protons (Figure S8 in SI). These results indicate that the methylation of $[Rh^{III}(L_1-H5)(Cl)_2]$ only occurs at the axial methylene position, although the negative charge should delocalize on the pyridine moiety as well.¹⁷ The site selectivity of the methylation reaction mainly depends on the stability of the methylated product rather than the population of the negative charge, because the methylation at the 3 or 5-position of the axial pyridine ring produces an unstable dearomatized product. The selective methylation of the axial methylene moiety is expected to embark a new method to provide a chiral TPA derivative by using a chiral reagent toward the development of TPA-based chiral catalysts for asymmetric transformation.

Redox properties of the deprotonated Rh(III)-TPA complexes were examined by cyclic voltammogram (CV) measurements in CH₃CN. The CV trace of $[Rh^{III}(L_1-H5)(Cl)_2]$ exhibited only one irreversible anodic wave at 0.08 V vs SCE even at 233 K under Ar as shown in Figure S10 in SI. In sharp contrast, a pseudo-reversible redox wave was observed for $[Rh^{III}(L_2-H5)(Cl)_2]$ with a peak separation $(\Delta E_{\rm P} = 75 \text{ mV})$ as shown in Figure 6a.¹⁸ The one-electron oxidation potential $(E_{1/2})$ of $[Rh^{III}(L_2-H5)(Cl)_2]$ was determined to be 0.40 V vs SCE in CH₃CN at 233 K. The reversibility was drastically improved by the introduction of the electron-withdrawing COOMe groups to the 5-positions of pyridine rings to lower the SOMO level, because the SOMO of one-electron oxidized $[Rh^{III}(L_2-H5)(Cl)_2]$ ($[Rh^{III}(Cl)_2(L_2-H5\bullet)]^+$) was indicated to delocalize over the axial methylene moiety and the 3- and 5-position of the pyridine ring by DFT calculations as shown in Figure S12 in SI. This reversibility allows us to elucidate the oxidation reaction as discussed below. In order to confirm that the oxidation of [Rh^{III}(L₂- $H5)(Cl)_2$ occurs at the metal center or the deprotonated TPA ligand, ESR measurements were performed on the solution of $[Rh^{III}(L_2-H5)(Cl)_2]$ with addition of 1 equiv of one-electron oxidant such as $[Ru^{III}(bpy)_3]^{3+}$ (bpy = 2,2'-bipyridyl) at 100 K. As a result, a featureless ESR signal was observed at g = 2.0043 as shown



Figure 6. (a) CV of $[Rh^{III}(L_2-H5)(Cl)_2]$ in CH₃CN at 233 K and (b) ESR spectrum of $[Rh^{III}(Cl)_2(L_2-H5\bullet)]^+$ measured at 100 K after the reaction of $[Ru^{III}(bpy)_3]^{3+}$ and $[Rh^{III}(L_2-H5)(Cl)_2]$.

in Figure 6b. Judging from the *g* value, the one-electron oxidation of $[Rh^{III}(L_2-H5)(Cl)_2]$ takes place on the ligand to form a Rh(III) complex with a carbon-centered ligand radical ($[Rh^{III}(Cl)_2(L_2-H5\cdot)]^+$) rather than at the Rh(III) center, in the light of the *g* value reported for Rh(IV)Cl₆ (g = 2.08).¹⁹ The carbon-centered radical was further confirmed by a reaction with a nitroxyl radical (2,2,6,6tetramethylpiperidine 1-oxyl = TEMPO) in CH₃CN at 233 K; the spin adduct, $[Rh(Cl)_2(L_2-H5-TEMPO)]^+$, was characterized by ESI-MS and ¹H NMR measurements (Figure S13 in the SI).^{20,21} Thus, this is the first observation of redox non-innocent behavior of a metal-bound TPA ligand, after deprotonation of a methylene proton of the ligand.

In summary, we have successfully demonstrated selective and reversible deprotonation of the methylene moiety of the axially coordinated pyridylmethyl arm of TPA ligands and the reactivity of the deprotonated TPA ligands in the Rh(III) coordination sphere. The deprotonation is governed by the stabilization of the dearomatized resonance structures (Scheme 1): For this reason, the coordination of a Lewis-acidic metal ion and the introduction of electronwithdrawing groups to the position where the negative charge can be delivered in the resonance structures are effective. The deprotonation of the TPA ligand allows us to encounter unprecedented redox-non-innocent characteristics of the versatile tetradentate ligands involving a carbon-centered radical of metal-bound TPA. The results described in this paper will provide a basis toward the development of new functionality of transition metal-TPA complexes.

ASSOCIATED CONTENT

Supporting Information. Figures S1–S13, Tables S1⁻S3 and crystallographic data for $[Rh^{III}(Cl)_2(L_2-H6)]^+$, $[Rh^{III}(L_1-H5)(Cl)_2]$, and $[Rh^{III}(Cl)_2(L_1-H5Me)]^+$ in the CIF format. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

kojima@chem.tsukuba.ac.jp

ACKNOWLEDGMENT

This work was partially supported by a Grant-in-Aid (No 24245011) from the Japan Society of Promotion of Science (JSPS, MEXT) of Japan, and the Cooperative Research Program of "Network Joint Research Center for Materials and Devices". We appreciate Dr. Motoo Shiro (Rigaku Corp., Japan) for his kind help in Xray analysis and Prof. S. Yamago (Kyoto University) for his helpful suggestion on the radical trapping.

REFERENCES

- (a) Grützmacher, H. Angew. Chem., Int. Ed. 2008, 47, 1814– 1818. (b) Gunanathan, C.; Milstein, D. Acc. Chem. Res. 2011, 44, 588–602. (c) Gunanathan, C.; Milstein, D. Chem. Rev. 2014, 114, 12024–12087.
- Gunanathan, C.; Ben-David, Y.; Milstein, D. Science 2007, 317, 790–792.
- (a) Ben-Ari, E.; Leitus, G.; Shimon, L. J. W.; Milstein, D. J. Am. Chem. Soc. 2006, 128, 15390–15391. (b) Zhang, J.; Leitus, G.; Ben-David, Y.; Milstein, D. Angew. Chem., Int. Ed. 2006, 45, 1113–1115.
- (4) (a) Lu, C. C.; Bill, E.; Weyhermüller, T.; Bothe, E.; Wieghardt,
 K. J. Am. Chem. Soc. 2008, 130, 3181–3197. (b) Shaffer, D.

W.; Ryken, S. A.; Zarkesh, R. A.; Heyduk, A. F. *Inorg. Chem.* **2011**, *50*, 13–21.

- (5) (a) Büttner, T.; Geier, J.; Frison, G.; Harmer, J.; Calle, C.; Schweiger, A.; Schönberg, H.; Grützmacher, H. Science 2005, 307, 235–238. (b) Donati, N.; Stein, D.; Büttner, T.; Schönberg, H.; Harmer, J.; Anadaram, S.; Grützmacher H. Eur. J. Inorg. Chem. 2008, 4691–4703.
- Tejel, C.; Ciriano, M. A.; del Río, M. P.; Hetterscheid, D. G. H.; Tsichlis i Spithas, N.; Smits, J. M. M.; de Bruin B. *Chem.-Eur. J.* 2008, *14*, 10932–10936.
- (7) (a) Jazdzewski, B. A.; Tolman, W. B. Coord. Chem. Rev. 2000, 200-202, 633-685. (b) Shimazaki, Y.; Tajima, T.; Tani, F.; Karasawa, S.; Fukui, K.; Naruta, Y.; Yamauchi, O. J. Am. Chem. Soc. 2007, 129, 2559-2568. (c) Storr, T.; Verma, R.; Pratt, R. C.; Wasinger, E. C.; Shimazaki, Y.; Stack, T. D. P. J. Am. Chem. Soc. 2008, 130, 15448-15459. (d) Pratt, R. C.; Lyons, C. T.; Wasinger, E. C.; Stack, T. D. P. J. Am. Chem. Soc. 2012, 134, 7367-7377. (e) Lyons, C. T.; Stack, T. D. P. Coord. Chem. Rev. 2013, 257, 528-540.
- (8) Anderegg, G.; Wenk, F. Helv. Chim. Acta 1967, 50, 2330– 2332.
- (9) (a) Jacobson, R. R.; Tyeklar, Z.; Farooq, A.; Karlin, K. D.; Liu, S.; Zubieta, J. J. Am. Chem. Soc. 1988, 110, 3690–3692. (b) Lim, M. H.; Rohde, J.; Stubna, A.; Bukowski, M. R.; Costas, M.; Ho, R. Y. N.; Münck, E.; Nam, W.; Que, L. Jr., Proc. Nat. Acad. Sci. U.S.A. 2003, 100, 3665–3670. (c) Hirai, Y.; Kojima, T.; Mizutani, Y.; Shiota, Y.; Yoshizawa, K.; Fukuzumi, S. Angew. Chem., Int. Ed. 2008, 47, 5772–5776.
- (10) (a) Que, L., Jr.; Ho, R. Y. N. *Chem. Rev.* 1996, *96*, 2607–2624.
 (b) Mirica, L. M.; Ottenwaelder, X.; Stack, T. D. P. *Chem. Rev.* 2004, *104*, 1013–1045. (c) de Bruin, B.; Budzelaar, P. H. M.; Gal, A. W. *Angew. Chem., Int. Ed.* 2004, *43*, 4142–4157.
- (11) Investigation on deprotonation of other metal-TPA complexes is currently ongoing in our laboratory.
- (12) (a) Lonnon, D. G.; Craig, D. C.; Colbran, S. B. Dalton Trans.
 2006, 3785–3797. (b) Ru^{II} complexes of L₂-H6 have been reported: Yamaguchi, M.; Kousaka, H.; Izawa, S.; Ichii, Y.; Kumano, T.; Masui, D.; Yamagishi, T.; Inorg. Chem. 2006, 45, 8342–8354.
- (13) Similar ¹H NMR spectral change due to the deprotonation of $[Rh^{III}(Cl)_2(L_1-H6)]^+$ was also observed in CD₃CN, although the line broadening of the signals assigned to the axial pyridyl-methyl moiety occurred by fast proton exchange between water and the highly basic deprotonated complex.
- (14) One of three independent $[Rh^{III}(L_1-H5)(Cl)_2]$ structures in the asymmetric unit was selected. The ORTEP drawing containing the other $[Rh^{III}(L_1-H5)(Cl)_2]$ structures also revealed a deprotonated form because there is no counter anions in the asymmetric unit.
- (15) Kaljurand, I.; Kütt, A.; Sooväli, L.; Rodima, T.; Mäemets, V.; Leito, I.; Koppel, I. A. *J. Org. Chem.* **2005**, *70*, 1019–1028.
- (16) Bordwell, F. G. Acc. Chem. Res. **1988**, 21, 456–463.
- (17) The HOMO of $[Rh^{III}(L_1-H5)(Cl)_2]$ was shown to delocalize on the methylene carbon, the 3- and 5-positions as well as the N-atom of the axial pyridine ring by DFT calculations (Figure S9 in SI).
- (18) The lifetime of $[Rh(Cl)_2(L_2-H5\bullet)]$ was estimated to be 8.3 s on the basis of CV measurements (Figure S12 in the SI).
- (19) Ellison, I. J.; Gillard, R. D. *Polyhedron* **1996**, *15*, 339–348.
- Beckwith, A. L. J.; Bowry, V. W.; Ingold, K. U. J. Am. Chem. Soc. 1992, 114, 4983–4992.
- (21) [RhCl₂(L₂-H5-TEMPO)]⁺: ESI-MS, m/z = 792.00; ¹H NMR
 (270 MHz, CD₃CN), 6.32 ppm (s, CH), 4.98 and 5.72 ppm (2 ABq, 16 Hz, CH₂)).

